

# Assessing the knowledge-base for commercially exploited marine fishes and invertebrates with a new global database of stock assessments

Alternative Title 1: A new global stock assessment database for exploited marine species

Alternative Title 2: Understanding marine population dynamics using a new global database

Suggested Running Title: A new global stock assessment database

Daniel Ricard<sup>a,\*</sup>, C  il  n Minto<sup>a,1</sup>, Julia Baum<sup>b,2</sup>, Olaf Jensen<sup>c,3</sup>

<sup>a</sup>*Department of Biology, Dalhousie University, Halifax, NS B3H 4J1, Canada*

<sup>b</sup>*Scripps Institution of Oceanography, UCSD, 9500 Gilman Drive, La Jolla, CA 92093-0202, USA*

<sup>c</sup>*School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195-5020, USA*

---

## Abstract

Data used to assess the status of individual fish stocks varies from very little information on many of the world's artisanal fisheries, to commercial landings at various levels of geographic and taxonomic aggregation, research surveys, and sophisticated population dynamics models that integrate many sources of information. Previous evaluations of the state of global fisheries have used catch or landings data, which may be poor proxies for fish stock abundances. A global compilation of stock assessment data in the mid-1990s enabled substantial syntheses of stock status; however its focus was on stock-recruitment relationships and it is now 15 years out of date. To facilitate contemporary

---

\*Corresponding author: Tel: 902-494-2146, Fax: 902-494-3736

*Email address:* ricardd@mathstat.dal.ca (Daniel Ricard)

<sup>1</sup>Current Address: Marine and Freshwater Research Centre, Galway-Mayo Institute of Technology, Dublin Road, Galway, Ireland

<sup>2</sup>Current Address: National Center for Ecological Analysis and Synthesis, UCSB, 735 State St. Suite 300, Santa Barbara, CA 93101, USA

<sup>3</sup>Current Address: Institute of Marine and Coastal Sciences, Rutgers University, 71 Dudley Road, New Brunswick, NJ 08901-8525, USA

syntheses, we have assembled a database of the most intensively studied commercially exploited marine fish stocks, including time series of: total biomass, spawner biomass, recruits, fishing mortality, and catch; reference points; and ancillary information on the life history, management, and assessment methods for each stock. Here, we present the first overview of this database and use it to evaluate the knowledge-base for assessed marine fishes. Globally, stock assessments were found for 324 stocks (288 species of fishes representing 45 families and 36 species of invertebrates representing 12 families), from 18 national and international management institutions. Assessments are available for only XX percent of global marine fisheries catches by weight and XX percent by value. There is substantial spatial variation in availability of assessed stocks, with XX percent coming from north temperate regions (North Atlantic, North Pacific). Geographic differences in assessment methods show that Statistical Catch at Age (SCA) models are widely used by the west coast of the U.S. (XX percent of assessments), regional fishery management organizations in the Pacific (XX percent of assessments), and New Zealand (XX percent of assessments); the east coast of the U.S. is transitioning from Virtual Population Analysis (VPA) to SCA (XX percent of assessments conducted since 2000 have used SCA); while VPA is still the dominant assessment technique in western Europe (XX percent of assessments).

*Keywords:* marine fisheries, meta-analysis, population dynamics models, relational database, stock assessment, synthesis

---

## Introduction

Marine wild capture fisheries provide more than 80 million tons of fisheries products (both food and industrial) per year and employ 43.5 million people (wild capture and aquaculture, (FAO, 2009b)). At the same time, fishing has been recognized as having one of the most widespread human impacts in the world's oceans (Halpern *et al.*, 2008), and the Food and Agricultural Organization of the United Nations (FAO) estimates that two-thirds of fish stocks globally are fully exploited or overexploited (FAO, 2009b).

While many fisheries have reduced exploitation rates to levels that should in theory promote recovery, overfishing continues to be a serious global problem (Worm *et al.*, 2009). Fishery managers are asked to address multiple competing objectives, including maximizing yields, ensuring profitability, reducing bycatch, and minimizing the risk of overfishing. Given the enormous social and economic costs (Rice *et al.*, 2003) and ecosystems consequences (Frank *et al.*, 2005; Myers *et al.*, 2007) of collapsed fisheries, it is imperative that we are able to quickly learn from successful and failed fisheries from around the world.

Effective management of exploited fish populations generally requires an understanding of where the current size and harvest rate lie in relation to the size and rate which maximize fishery benefits or limit the risk of overfishing. This process of quantitative determination of stock status and estimation of reference points is called stock assessment. Some fisheries in developing countries have apparently provided sustainable yields for long periods of time without formal stock assessment (e.g. many community-managed fisheries in Oceania, (Johannes, 2002)). This has been achieved by limiting harvest rates, often through gear restrictions or seasonal or area closures. In modern industrialized fisheries, however, where fishing capacity exceeds the productivity of fished stocks stock assessment is an integral component of responsible management (Hilborn and Walters, 1992).

The global databases of fishery landings compiled by FAO (FAO, 2009a) and synthesized by the Sea Around Us project (Watson *et al.*, 2004) have proven to be valuable resources for understanding the status of fisheries worldwide; however, catch data alone can be misleading when used as a proxy for stock size. Many papers have used these data to examine changes in fishery status (Worm *et al.*, 2006; Costello *et al.*, 2008), including changes in trophic level (Pauly *et al.*, 1998; Essington *et al.*, 2006; Newton *et al.*, 2007). Most of these analyses rely (either explicitly or implicitly) on the assumption that catch or landings is a reliable index of stock size. Critics have pointed out that catch can change for a number of reasons unrelated to stock size, including changes in targeting, fishing restrictions, or market preferences (de Mutsert *et al.*, 2008; Mu-

rawski *et al.*, 2007; Hilborn, 2007). Standardizing catch by the amount of fishing effort (catch-per-unit-of-effort, CPUE) is an improvement, particularly when these data are modeled to account for spatial, temporal, and operational factors affecting the CPUE, but CPUE can still be an unreliable index of relative abundance since it is difficult to account for all relevant factors (Hutchings and Myers, 1994; Harley *et al.*, 2001; Walters, 2003; Polacheck, 2006).

Stock assessments consider time series of catch along with other sources of information such as: natural mortality rates, changes in size or age composition, stock-recruitment relationships, and CPUE of different sectors or of fishery-independent surveys. Because they integrate across multiple sources of information, stock assessment models are thought to provide a more accurate picture of changes in abundance than catch data alone (Sibert *et al.*, 2006). Yet, without a current and comprehensive database of stock assessments, scientists wishing to conduct comparative analyses of marine fish population dynamics and fishery status have little choice but to use problematic catch data.

The first global database of stock assessment information, the Myers Stock Recruitment Database, was developed by Ransom Myers and colleagues in the mid-1990s (Myers *et al.*, 1995b). While the database was primarily known for its time series of stock and recruitment, it did contain time series of fishing mortality rates for many stocks; biological reference points were, however, largely absent. The original release version of the Myers database (Myers *et al.*, 1995b) included spawning stock size and recruitment time series for 274 stocks representing 92 species as well as fishing mortality rates time series for 144 stocks. The number of entered stocks grew to approximately 652 stocks (507 with at least one SR pair) over the period from 1995-2005. It was used to: 1) decisively answer the question of whether recruitment shows any relationship to spawning stock size (Myers and Barrowman, 1996), 2) investigate potential depensation in stock-recruitment relationships (Myers *et al.*, 1995a; Liermann and Hilborn, 1997; Garvey *et al.*, 2009), 3) discover generalities in the annual reproductive rates of fishes (Myers *et al.*, 1999, 2002b), 4) investigate density-dependence in juvenile mor-

tality (Myers, 2001; Minto *et al.*, 2008), 5) develop informative Bayesian priors on steepness (Myers *et al.*, 1999, 2002a; Dorn, 2002), and 6) examine patterns of collapse and recovery in exploited fish populations (Hilborn, 1997; Hutchings, 2000, 2001) .

Although the original Myers database (Myers *et al.*, 1995b) was critical for motivating comparative analyses in fisheries science, most of the stocks are now 15 years out of date. For stocks that were depleted in 1995, the past 15 years include valuable observations at low stock size or of a recovering population, both of which are critical for estimating population dynamics parameters such as the behaviour of the stock-recruitment relationship near the origin. In addition, there have been numerous improvements in stock assessment methodologies (including important advances in statistical catch-at-age or catch-at-length models) and assessments have been conducted for the first time for many species.

Meta-analyses of fishery status also have been hampered by the lack of a global assessment database containing biological reference points (BRPs, e.g., the total/spawning biomass and fishing mortality rate that produce Maximum Sustainable Yield (MSY),  $B_{MSY}$  and  $F_{MSY}$ ). Knowledge of BRPs is important if stocks are to be managed for high yields that can be sustained over time (Mace, 1994). Without information on reference points, previous analyses of stock assessments or catch data have been forced to use non-biological thresholds to define fishery status, such as the greatest 15-year decline (Hutchings and Reynolds, 2004) or 10 percent of maximum catch (Worm *et al.*, 2006). Ad hoc reference points based on some fraction of the maximum of a time series also have undesirable statistical properties and can result in false collapses when applied to inherently variable time series of catch or abundance (Wilberg and Miller, 2007; Branch, 2008). Complicating comparisons of fishery status is the fact that different BRPs are used in different parts of the world and even the same BRP can be used in a different manner, for example, as a target or a limit.

Here we present a new global database of stock assessments for commercially exploited marine fish populations. The database is an update and extension of that developed by Ransom Myers, and is named the RAM Legacy database in honor of his

pioneering contribution. This effort is the first global stock assessment database to:

1. Use a formal relational database structure;
2. Use source control software to organise release versions;
3. Include metadata related to the geographic location of the stock, the type of assessment model used, and the original source document for the assessment data;
4. Include biological reference points and stock-specific life history information.

We use the new RAM Legacy database (Version 1.0, 2010) to evaluate the knowledge-base for commercially exploited marine populations in terms of institutional contributions, geography, taxonomy, ecology, timespan, stock assessment methodologies and BRPs, and to compare it to that of global fisheries catches. We then evaluate the status of assessed stocks globally, and by management body, referencing all stocks to a comparable benchmark. Finally, we discuss biases in the knowledge base for assessed marine species, highlight potential applications of the database, point out its caveats and limitations, and outline directions for future development.

## Methods

### *The RAM Legacy database*

The RAM Legacy database is a global relational database designed and developed to store data from all current and accessible population dynamics model-based fisheries stock assessments for marine fish and invertebrate populations. Time series of spawning stock biomass (SSB), total biomass (TB), recruits (R), total catch (TC) or landings (TL), and fishing mortality (F) from individual stock assessments form the core of the database. Apart from catch/landings, these time series are not raw data, but rather the output of population dynamics models; depending on the type of assessment model not all of these time series were available for every stock. The database also contains details about the time series data, including the age and sex of spawners, age of recruits, and the ages used to compute the fishing mortality, as well as BRPs and some life history information (e.g. growth parameters, age and length at 50% maturity, natural mortality). Metadata for each stock assessment consists of taxonomic information about the species and the geographic location of the stock (detailed in “Links to related databases”), the management body that conducted the assessment, the assessment methodology, the reference for the stock assessment document, the name of the recorder entering the assessment data, and the date the assessment was entered. Some assessments, particularly those for invertebrates, were based only on CPUE time series rather than population dynamics models. While we included these in the database (n=26), the database description and analyses presented here focus on those stocks assessed using population dynamics models.

Over the past two and a half years, we have employed a variety of search methods in an attempt to obtain all recent fisheries stock assessments. Publically available stock assessment reports available from the internet were the primary data source. These reports were obtained either from the website of the relevant management agency or directly from stock assessment scientists. Other assessments were obtained from the primary literature and through personal contacts at fisheries management agencies.

Significant contributions also were made by the other members of the National Center for Ecological Analysis and Synthesis (NCEAS) working group “Finding common ground in marine conservation and management”. Relevant assessment data were first transferred into a standardized spreadsheet template by a number of recorders, including ourselves, assessment authors, our NCEAS collaborators, and associated graduate students and postdoctoral researchers, and then uploaded into the relational database management system by the first author.

#### *Database structure and advantages*

The database is implemented in the Open Source PostgreSQL relational database management system (RDBMS) (PostgreSQL Global Development Group, 2009), and includes tables for the assessment metadata, time series values, time series units, and biometrics (a catch-all term for data, such as life history characteristics or BRPs, that are not part of a time series). The entity relationship diagram of the database and its component tables can be found in the Supplementary materials.

RDBMSs form the server back-end to a great many applications of interest to ecologists, including web-clients and GIS software, and have a number of advantages over spreadsheet or flat text file data compilations. First, housing stock assessments in an RDBMS allows multiple users to concurrently access and extract subsets of data in an efficient and reproducible manner. Second, with the development of Application Programming Interfaces (APIs) that allow analytical softwares to directly communicate and extract data from the database, a common data environment is established, independent of one’s choice of analytical software (e.g., SAS:SAS ACCESS, Matlab: Matlab/Database, R:RDBI/RODBC, Perl:DBI, etc.). Users familiar with Structured Query Language (SQL) can also query the database directly from their analytical software of choice and the same SQL query will extract the same data through each of these applications. Third, data products tailored to specific projects can be generated and stored as dynamic (i.e. continually updated) “views” within the database. These are typically rectangular, spreadsheet-like results of an expansive query of the relevant



tables that can be readily read into all commonly-used analytical software. The use of views is, however, greatly advantageous over manipulating spreadsheets or flat text files for importing into a specific analytic software, which runs the risk of losing data integrity (e.g. multiple copies) and becomes impractical with large, non-rectangular datasets and multiple users.

#### *Data integrity and quality control*

We have employed several mechanisms to ensure that the database is of high quality. During the data recording process, assessment authors often were contacted to clarify aspects of the assessment or to obtain more detailed data. Time series data presented only in assessment report figures were, for example, only entered into the database if the exact numbers could be obtained from the assessment or its authors. The only exception to this is Indian Ocean bigeye tuna, a stock assessed by the Indian Ocean Tuna Commission (IOTC) for which we could not obtain the model output from the assessment authors. In cases where multiple models were presented in an assessment without a preferred or best model being denoted, we attempted to ascertain which model was preferred by the stock assessment scientist, but included all model results whenever this was not possible. Once uploaded into the database all stock assessments underwent an additional Quality Assurance/Quality Control (QA/QC) step, to ensure that the entered data replicated that of the original assessment document exactly. This process consisted of creating a QA/QC summary document for each assessment, containing summary details of the stock, a selection of biometrics and ratios for comparison (e.g. current status relative to BRP), and time series plots of the biomass, recruitment, and exploitation trajectories. QA/QC documents were then returned to assessment recorders and an electronic trail of subsequent correspondence was captured using a bug tracking system. Recorders were responsible for checking, and where necessary correcting, their QA/QC documents, after which all corrections were transmitted back to the operational database and a quality controlled flag was inserted to signify the assessment had passed the check. Only assessments that have passed this QA/QC step

are available for subsequent analyses.

#### *Links to related databases*

To facilitate integration of the RAM Legacy database with related databases, such as Fishbase (Froese and Pauly, 2009) and the Sea Around Us Project’s (SAUP) global landings database (Watson *et al.*, 2004), each species present in the RAM Legacy database was assigned a matching FishBase species name and species code, a matching SAUP taxon code, and taxonomic information from the Integrated Taxonomic Information System (ITIS) (<http://www.itis.gov>). Additionally, each stock was assigned to a primary (and in some cases secondary and tertiary) Large Marine Ecosystem (LME) (NOAA, 1998). LMEs encompass the continental shelves of the world’s oceans and represent the most productive areas of the oceans. Open ocean areas beyond the continental shelves are, however, not included. Large, highly migratory oceanic species such as tuna were therefore assigned to new categories “Atlantic High Seas”, “Pacific High Seas”, “Indian High Seas”, and “Subantarctic High Seas”.

#### *Assessing the knowledge-base for commercially exploited stocks*

We assess the knowledge-base for commercially exploited stocks, as represented by the RAM Legacy database, using a variety of metrics. To evaluate the taxonomic scope of the database, we compare the taxonomy of assessed stocks with the diversity of i) all marine fishes (as represented by FishBase), and ii) marine fishes in global fisheries catches (as represented by the SAUP database), and discuss taxonomic biases in species included in catch data and in populations assessed using stock assessments. We evaluate the ecological scope of assessed stocks in terms of age at sexual maturity as reported in the assessments, and trophic level of those assessed stocks as reported in FishBase. We overview the types of assessment models used, and BRPs estimated, for all stock assessments and by each management body; we also evaluate how the frequencies of different assessment model types and BRPs has changed over time. To determine what fraction of world wild-capture fisheries landings come from assessed stocks, we used the SAUP’s average global fisheries catches from the most recent ten years of available

data (1995-2004); we also discuss limitations to obtaining assessments for some of the world's major fisheries.

### *Assessing the status of commercially exploited marine stocks*

We evaluate the status of assessed stocks overall and by management body, using standard reference points so that all stocks are referenced to a comparable benchmark. Following Froese and Proelß (2010) and Worm *et al.* (2009), we compare the current biomass and exploitation rate of stocks relative to their MSY reference points,  $B_{MSY}$  and  $u_{MSY}$ , respectively.

We do not advocate the use of MSY targets for management, but still report MSY-related BRPs because they are the most commonly estimated BRP and can be used to compare multiple stocks. For those assessments that did not contain MSY reference points, but did include total catch ( $TC_{i,s}$ ,  $i \in 1, \dots, n_s$ ) and total biomass ( $TB_{i,s}$ ,  $i \in 1, \dots, n_s$ ) time series data, we used a Schaefer surplus production model to estimate total biomass and exploitation rate at MSY ( $TB_{MSY_s}$  and  $u_{MSY}$ , respectively). Surplus production of stock  $s$  in year  $t$ ,  $P_{s,t}$ , is a commonly used measure of stock productivity, representing the amount of catch that can be taken while maintaining the biomass at a constant size, and can be calculated as:

$$P_{s,t} = TB_{s,t+1} - TB_{s,t} + TC_{s,t} \quad (1)$$

where,

$TB_{s,t}$  is the total biomass of stock  $s$  in year  $t$

$TC_{s,t}$  is the total catch of stock  $s$  in year  $t$

We fit a Schaefer surplus-production model, which is based on a logistic model of population growth to the catch and total biomass time series data. The predicted surplus production in each year in the Schaefer model is given by:

$$\hat{P}_{s,t} = \frac{4mTB_{s,t}}{K} - 4m \left( \frac{TB_{s,t}}{K} \right)^2 \quad (2)$$

where,

$m$  is the maximum sustainable yield, equal to  $rK/4$

$K$  is the carrying capacity or equilibrium total biomass in the absence of fishing (Hilborn and Walters, 1992)

We estimated the model parameters ( $m$  and  $K$ ) using maximum likelihood in AD Model Builder (ADMB Project, 2009) assuming that the residuals  $\epsilon_{s,t} = P_{s,t} - \hat{P}_{s,t}$  were normally distributed. For the Schaefer model,  $B_{MSY}$  is simply  $0.5K$ , and the harvest rate that results in maximum sustainable yield,  $u_{MSY}$ , is  $m/B_{MSY}$ . Carrying capacity was constrained to be less than twice the maximum observed total biomass.

Finally, we discuss potential applications of the database, point out its limitations and caveats about its use, and outline directions for future development.

Statistical analyses and plot generation were conducted with the R Environment for Statistics and Graphics (Development Core Team, 2009) using the packages RODBC (Ripley *et al.*, 2009), KernSmooth (original by Matt Wand. R port by Brian Ripley., 2009), gregmisc (?), xtable (Dahl, 2009), ape (Paradis *et al.*, 2004), gsubfn (Grothendieck, 2010), IDPmisc (Ruckstuhl *et al.*, 2009), doBy (with contributions from Kevin Wright and Leidi., 2010), and beanplot (Kampstra, 2008). Figure 1 was generated using the Generic Mapping Tools (Wessel and Smith, 1991).

## Results

### *The knowledge-base for commercially-exploited marine stocks*

In total, 324 recent stock assessments for 288 marine fish and 36 invertebrate populations are included in the RAM Legacy database (Version 1.0, 2010; Table S1). Together these comprise time series of catch/landings were available for 308 stocks (95%), of SSB for 271 stocks (84%), and of recruitment for 269 stocks (83%) (Table S1).

### *Management bodies and geography*

Stock assessments are derived from fisheries management bodies in Europe, the United States, Canada, New Zealand, Australia, Russia, South Africa and Argentina (Table 1). Also included are assessments conducted by eight Regional Fisheries Management Organizations (RFMOs), in the Northwest Atlantic, Atlantic, Pacific and Indian Ocean (Table 1). Assessments from the United States comprise by far the most stocks of any country or region (n=139); assessments from the European Union's management body ICES comprise the the second greatest number of stocks (n=63). Whereas nations are responsible for managing all populations within their EEZs, RFMOs typically focus on a certain type of species (e.g. halibut, tunas) or fisheries (e.g. pelagic high seas) within a given area and hence assess a smaller number of stocks.

Most assessments come from North America, Europe, Australia, New Zealand and the High Seas, while few are entered from regions such as Southeast Asia, South America, and the Indian Ocean (outside Australian waters) (Figure 1). Assessments were available for 31 LMEs, with the greatest number of assessed stocks coming from Northeast U.S. Continental Shelf (n=58), California Current (n=35), New Zealand Shelf (n=29), Gulf of Alaska (n=26), Celtic-Biscay Shelf (n=26), East Bering Sea (n=22) and Southeast U.S. Continental Shelf (n=20) (Figure 1).

### *Taxonomy*

157 species from 57 families and 20 orders are included in the database (Figure 2). Five taxonomic orders (Gadiformes (n=67), Perciformes (n=62), Pleuronectiformes

(n=53), Scorpaeniformes (n=41) and Clupeiformes (n=36)) account for 80% of available stock assessments. Of these, Perciformes, the most speciose Order of marine fishes are in fact underrepresented in the database (46.04% of all marine fish species vs. 19% of all marine fish assessments), while it is the other four orders that are taxonomically overrepresented: Clupeiformes (2.1% of marine fishes vs. 11% in the database), Gadiformes (3.3% of marine fishes vs. 21% in the database), Pleuronectiformes (4.5% of marine fishes vs. 17% in the database), Scorpaeniformes (8.5% of marine fishes vs. 12% in the database) (Figure 3).

Assessed marine fish stocks in the RAM Legacy database comprise a relatively small proportion of harvested taxa (24% of fish species from the SAUP database) and an even smaller proportion of marine fish biodiversity (1% of fish species in FishBase; Figure 3). In turn, catches from the SAUP database, which come from 649 species and 36 orders (Figure 3), represent only 5% of the 12339 species and 67% of the 54 different orders present in FishBase (Figure 3).

#### *Timespan*

Of the 324 stock assessments, time series data of catch/landings were available for 308 stocks (95%), of SSB for 271 stocks (84%), and of recruitment for 269 stocks (83%). The median lengths of catch/landings, SSB, and recruitment timeseries were 38, 34, and 33 years, respectively. The time period covered by 90% of assessments is: catch/landings (1967-2007), SSB (1972-2007), recruitment (1971-2006), while that covered by 50% of assessments is: catch/landings (1983-2004), SSB (1985-2005), recruitment (1984-2003) (Figure 5).

#### *Stock assessment methodologies and BRPs*

In addition to the 324 assessments in the database, indices of relative abundance from scientific surveys are available for an additional 26 stocks. The three most common assessment methods were Statistical catch-at-age/length models (n=164), Virtual Population Analyses (n=90) and Biomass dynamics model (n=45).. .....Need to add a sentence here about the regional differences.... .

The total number and percentage of assessments that reported biomass- or exploitation-based reference points of any sort was 257 (81%) and 222 (69%), respectively. From these assessments, 66 report both a biomass-based and an exploitation-based BRP and appear as solid dots on Figures 6 and 7. Schaefer-derived BRPs add an additional 175 assessments, for a total of 241 assessments used to generate Figure 6. Overall, 58% of assessed stocks are below their biomass-related MSY BRP and 30% are above their exploitation-related MSY BRP. Different management bodies have different overall status of current biomass to BRPs (Figure 7).

### *Ecology*

Assessed species span a range of ecological traits. In terms of their trophic level, we see XXX (Figure 4). The total number and percentage of assessments that reported any life-history information (growth, maturity, fecundity) was 288.

## Discussion

The RAM Legacy Database of available stock assessments for the world's marine fisheries, although non-exhaustive, provides a basis to evaluate the existing knowledge-base of exploited marine populations. The stocks comprising the database are predominantly from developed nations with properly identified fisheries management bodies and tend to cover a recent time period. Few assessments extend back beyond 30 years from present. The taxonomic makeup of available assessments is a very limited subset of the accepted taxonomic coverage of marine species worldwide.

### *Biases*

Even in developed countries, however, not all stocks are assessed. For example, in 2007, of the 528 fish and invertebrate stocks recognized by the National Marine Fisheries Service (NMFS), only 179, or slightly over one-third, were fully assessed (National Marine Fisheries Service, 2008). An assessment by the European Environment Agency (EEA) in 2006 indicated that the percentage of commercial landings obtained from assessed stocks ranged between 66-97 percent in northern European waters and 30-77 percent in the Mediterranean (European Environment Agency, 2009). The New Zealand Ministry of Fisheries reports the status of 117 stocks or sub-stocks out of a total of 628 stocks managed under New Zealand's Quota Management System (New Zealand Ministry of Fisheries, 2009). In Australia, 98 federally managed stocks have been assessed (Wilson *et al.*, 2009) out of an unknown total. The extent to which stocks are assessed elsewhere in the world is currently unknown.

### *Geographic bias*

There are important geographic biases in the amount of assessments entered per LME

The question of geographic bias relates to whether: 1) an assessment is conducted on a stock; 2) it is possible to access the assessment; and 3) the non-exhaustive collation we undertook may have overlooked the assessment. Whether an assessment is conducted



for a given stock depends upon a multitude of factors, including the economic value of the stock and availability of fiscal resources to collect the data required for an assessment. How accessible assessments are for entry depends upon the transparency and access policies of the relevant management agencies, which varies geographically. Similarly, the legal context where fisheries are prosecuted will strongly influence the requirement for conducting stock assessments. In the United States, the Magnuson-Stevens Act defines what stocks are to be monitored and managed, hence a large number of the assessments in the RAM Legacy database are under the jurisdiction of the US National Marine Fisheries Services. Our incomplete search for assessments could also give rise to geographic biases, as concerted collation efforts have only been conducted in those assessment-rich regions of It is hoped that readers of this article can assist in correcting these biases by participating in future updates of the RAM Legacy database.

Note Also: from Figure 4 SOFIA 2008 (FAO 2009) - The top 10 wild-caught marine fisheries producer countries are China, Peru, U.S.A., Indonesia, Japan, Chile, India, Russia, Thailand, Philippines. We only have assessments for U.S.A. and Russia (and only 2 for Russia!). Most major fish producing countries do not do assessments or do not make them accessible, still to be decided how to phrase this.

#### *Taxonomic bias*

Taxonomic biases in those species that are assessed include ...

#### *Caveats and limitations*

Assessment outputs e.g. biomass timeseries, are model estimates, not raw data. The uncertainty associated with these estimates should be carried forth in subsequent analyses. The RAM Legacy database structure allows for estimates of uncertainty (standard errors, 95% credible/confidence intervals), however these estimates are often missing from assessments either because they aren't produced by the assessment model (e.g. non-bootstrapped VPA assessments) or the focus of the assessment document was on central tendency (e.g. mean biomass), not the associated uncertainty. Note that this view is changing with the advent of MCMC approaches to Bayesian inference

for assessments, bootstrap methods, statistical catch-at-age models (ADMB Project, 2009) and a general focus on uncertainty (Walters and Maguire, 1996). As with any analysis, clearer inference on the strength of a signal is available when all uncertainty in the data is carried forth. This represents a difficulty for large-scale analyses of fisheries data in that in an ideal world one would access the raw data per sub-unit (e.g. stock) and carry forth the uncertainty at all levels of the analysis. In the case of assessments, the raw data is typically catch-at-age matrices and potentially survey indices. To understand the fleet characteristics and survey stratification schema for each stock in a potentially global meta-analysis would be extremely time consuming and error-prone. So, the expert opinion of those researchers most familiar with the data, stock assessment authors, is used but without accompanying uncertainty estimates the strength of conclusions drawn may be weakened.

The original database developed by Dr. Myers was used to address a variety of ecological question derived from stock-recruit relationships. This was possible because of the timeseries of stock and recruitment generated in assessment using VPA-type models. With the increasing use of statistical catch-at-age/length models, an underlying stock-recruit relationship is assumed to exist. Estimating the parameters of a stock-recruit relationship using timeseries from such models does not constitute a valid statistical procedure. More generally, the increasing use of Bayesian methods that incorporate prior information poses a challenge for meta-analysis of such model outputs.

Point data are stored in the database with an associated unit, value and year. We expect to also include age-varying and length-varying data such as maturity ogive and age-disaggregated natural mortality in subsequent releases of the RAM Legacy database. In addition to the initial aim of providing reliable access to timeseries information about stocks, we hope to also stimulate research in the relationships of life-history characteristics and their relation to exploitation.

### *Future Development*

It is anticipated that the RAM Legacy database will continue to grow with hitherto unentered stocks and updated assessments for already included stocks. The ultimate goal is to provide a data standard for researchers to use results from multiple regions to assist in their own applied and fundamental research in population ecology, fisheries science, and conservation biology. The development of a standard for assessment reporting would assist in realizing this goal. For example, ICES assessments have a very regular standard, including agreed-upon reference points and regular estimate reporting. This makes the process of data collation much more routine than unstandardized documents where the recorder trawls through a report for information. Certainly different stocks and regions require different formats but the basic output tables, consisting of total and spawning biomass, recruitment, catch/landings, estimated fishing mortality over vulnerable age groups, associated measures of uncertainty, and commonly-used reference points would streamline the process immensely. A process whereby the assessment spreadsheets are filled out at each assessment meeting would be the least error prone method. In return, the assessment scientists can access results for a global collation of assessments to further their own research initiatives in population assessment and management. Other products include management-agency-level reports containing summaries of all stocks within their remit. Future versions to the database will also include timelines of management actions per stock. The RAM Legacy database contains the corresponding species codes to the Sea Around Us Project and FishBase, thus facilitating researchers's use of a global fisheries data "toolkit" to address questions on the relationships between life history attributes and resulting population dynamics in an exploited setting.

## **Availability of the database**

Contributions or corrections to the existing database, as well as requests to use the database (subject to standard “Fair Use” policies), should be directed to the corresponding author.

## **Acknowledgments**

We sincerely thank all of the fisheries scientists around the world for conducting the stock assessments that form the basis of this new global database. We are also grateful for the database contributions, advice, and support of Trevor Branch, Jeremy Collie, Laurence Fauconnet, Mike Fogarty, Rainer Froese, Ray Hilborn, Jeff Hutchings, Simon Jennings, Heike Lotze, Pamela Mace, Michael Melnychuk, Ana Parma, Renée Préfontaine, Reg Watson, Boris Worm, Dirk Zeller, and the financial support of the National Science Foundation through an NCEAS Working Group, the Natural Sciences and Engineering Research Council (NSERC) of Canada, the Canadian Foundation for Innovation, the David H. Smith Conservation Research Fellowship, the Schmidt Research Vessel Institute, and the Census of Marine Life (CoML/FMAP).

- ADMB Project (2009). AD Model Builder: automatic differentiation model builder. Developed by David Fournier and freely available from [admb-project.org](http://admb-project.org).
- Branch, T. (2008). Not all fisheries will be collapsed in 2048. *Marine Policy* 32, 38–39.
- Costello, C., Gaines, S.D. and Lynham, J. (2008). Can catch shares prevent fisheries collapse? *Science (Washington)* 321(5896), 1678–1681.
- Dahl, D.B. (2009). *xtable: Export tables to LaTeX or HTML*. <http://CRAN.R-project.org/package=xtable>, r package version 1.5-6.  
**URL:** <http://CRAN.R-project.org/package=xtable>
- de Mutsert, K., Cowan, Jr., J.H., Essington, T.E. and Hilborn, R. (2008). Reanalyses of Gulf of Mexico fisheries data: Landings can be misleading in assessments of fisheries and fisheries ecosystems. *Proceedings of the National Academy of Sciences* 105(7), 2740–2744.
- Development Core Team, R. (2009). R: a Language and Environment for Statistical Computing.
- Dorn, M. (2002). Advice on West Coast rockfish harvest rates from Bayesian meta-analysis of stock-recruit relationships. *North American Journal of Fisheries Management* 22, 280–300.
- Essington, T.E., Beaudreau, A.H. and Wiedenmann, J. (2006). Fishing through marine food webs. *Proceedings of the National Academy of Science* 103(9), 3171–3175.
- European Environment Agency (2009).
- FAO (2009a). FISHSTAT-PC: Data retrieval, graphical and analytical software for microcomputers.
- FAO (2009b). SOFIA report 2008.

- Frank, K., Petrie, B., Choi, J. and Leggett, W. (2005). Trophic cascades in a formerly cod-dominated ecosystem. *Science (Washington)* 308(5728), 1621–1623.
- Froese, R. and Proelß, A. (2010). Rebuilding fish stocks no later than 2015: will Europe meet the deadline? *Fish and Fisheries* (in press).
- Froese, R. and Pauly, D. (2009). FishBase [www.fishbase.org](http://www.fishbase.org), version (10/2009). World Wide Web electronic publication.
- Garvey, J., Wright, R. and Marschall, E. (2009). Searching for threshold shifts in spawner-recruit data. *Canadian Journal of Fisheries and Aquatic Sciences* 66, 312–320.
- Grothendieck, G. (2010). *gsubfn: Utilities for strings and function arguments*. **http:** [//CRAN.R-project.org/package=gsubfn](http://CRAN.R-project.org/package=gsubfn), r package version 0.5-2.  
**URL:** <http://CRAN.R-project.org/package=gsubfn>
- Halpern, B., Walbridge, S., Selkoe, K. *et al.* (2008). A global map of human impact on marine ecosystems. *Science (Washington)* 319(5865), 948–952.
- Harley, S., Myers, R. and Dunn, A. (2001). Is catch-per-unit-effort proportional to abundance? *Canadian Journal of Fisheries and Aquatic Sciences* 58, 1705–1772.
- Hilborn, R., ed. (1997). *The frequency and severity of fish stock declines and increases*. Developing and sustaining world fisheries resources. Proceedings of the 2nd World Fisheries Congress. CSIRO Publishing, Victoria, Australia.
- Hilborn, R. and Walters, C.J. (1992). *Quantitative Fisheries Stock Assessment: Choice, Dynamics and Uncertainty*. Kluwer Academic Publishers.
- Hilborn, R. (2007). Biodiversity loss in the ocean: how bad is it? *Science (Washington)* 316(5829), 1281.
- Hutchings, J. (2000). Collapse and recovery of marine fishes. *Nature (London)* 406, 882–885.

- Hutchings, J. (2001). Influence of population decline, fishing, and spawner variability on the recovery of marine fishes. *Journal of Fish Biology* Suppl. A, 306–322.
- Hutchings, J. and Myers, R. (1994). What can be learned from the collapse of a renewable resource? Atlantic cod, *Gadus morhua*, of Newfoundland and Labrador. *Canadian Journal of Fisheries and Aquatic Sciences* 51, 2126–2146.
- Hutchings, J. and Reynolds, J. (2004). Marine fish population collapses: Consequences for recovery and extinction risk. *BioScience* 54, 297–309.
- Johannes, R. (2002). The renaissance of community-based marine resource management in Oceania. *Annual Review of Ecology, Evolution, and Systematics* 33, 317–340.
- Kampstra, P. (2008). Beanplot: A Boxplot Alternative for Visual Comparison of Distributions. *Journal of Statistical Software, Code Snippets* 28(1), 1–9. **http:** [//www.jstatsoft.org/v28/c01/](http://www.jstatsoft.org/v28/c01/).  
**URL:** <http://www.jstatsoft.org/v28/c01/>
- Liermann, M. and Hilborn, R. (1997). Depensation in fish stocks: a hierarchic Bayesian meta-analysis. *Canadian Journal of Fisheries and Aquatic Sciences* 54, 1976–1984.
- Mace, P.M. (1994). Relationships between Common Biological Reference Points Used as Thresholds and Targets of Fisheries Management Strategies. *Canadian Journal of Fisheries and Aquatic Sciences* 51(1), 110–122.
- Minto, C., Myers, R.A. and Blanchard, W. (2008). Survival variability and population density in fish populations. *Nature (London)* 452, 344–347.
- Murawski, S., Methot, R. and Tromble, G. (2007). Letter to the editors of Science. *Science (Washington)* 316, 1281.
- Myers, R.A. (2001). Stock and recruitment: Generalizations about maximum reproductive rate, density dependence and variability using meta-analytic approaches. *ICES Journal of Marine Science* 58, 937–951.

- Myers, R. and Barrowman, N. (1996). Is fish recruitment related to spawner abundance? *Fishery Bulletin* 94, 707–724.
- Myers, R., Barrowman, N., Hilborn, R. and Kehler, D. (2002a). Inferring bayesian priors with limited direct data: applications to risk analysis. *North American Journal of Fisheries Management* 22, 351–364.
- Myers, R., Barrowman, N., Hutchings, J. and Rosenberg, A. (1995a). Population dynamics of exploited fish stocks at low population levels. *Science (Washington)* 269, 1106–1108.
- Myers, R., Baum, J., Shepherd, T., Powers, S. and Peterson, C. (2007). Cascading Effects of the Loss of Apex Predatory Sharks from a Coastal Ocean. *Science* 315(5820), 1846–1850.
- Myers, R., Bowen, K. and Barrowman, N. (1999). Maximum reproductive rate of fish at low population sizes. *Canadian Journal of Fisheries and Aquatic Sciences* 56, 2404–2419.
- Myers, R., MacKenzie, B., Bowen, B. and Barrowman, N. (2002b). What is the carrying capacity for fish in the ocean? A meta-analysis of population dynamics of North Atlantic cod. *Canadian Journal of Fisheries and Aquatic Sciences* 58, 1464–1476.
- Myers, R.A., Bridson, J. and N.J., B. (1995b). Summary of Worldwide Spawner and Recruitment Data. *Canadian Technical Report of Fisheries and Aquatic Sciences, No. 2020* p. 327.
- National Marine Fisheries Service (2008). Status of US Fisheries 2007. Tech. rep., NMFS. [Http://www.nmfs.noaa.gov/sfa/statusoffisheries/SOSmain.htm](http://www.nmfs.noaa.gov/sfa/statusoffisheries/SOSmain.htm).
- New Zealand Ministry of Fisheries (2009). Stock Status <http://fs.fish.govt.nz/Page.aspx?pk=16>.



- Newton, K., Côté, I., Pilling, G., Jennings, S. and Dulvy, N. (2007). Current and Future Sustainability of Island Coral Reef Fisheries. *Current Biology* 17(7), 655–658.
- NOAA (1998). Large Marine Ecosystems. Tech. rep., National Oceanic and Atmospheric Administration.
- original by Matt Wand. R port by Brian Ripley., S. (2009). *KernSmooth: Functions for kernel smoothing for Wand Jones (1995)*. <http://CRAN.R-project.org/package=KernSmooth>, r package version 2.23-2.  
**URL:** <http://CRAN.R-project.org/package=KernSmooth>
- Paradis, E., Claude, J. and Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289–290.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R. and Torres, Jr., F. (1998). Fishing Down Marine Food Webs. *Science (Washington)* 279(5352), 860–863.
- Polacheck, T. (2006). Tuna longline catch rates in the Indian Ocean: Did industrial fishing result in a 90% rapid decline in the abundance of large predatory species. *Marine Policy* 30, 470–482.
- PostgreSQL Global Development Group (2009). PostgreSQL, version 8.3.8. <http://www.postgresql.org>.
- Rice, J., Shelton, P., Rivard, D., Chouinard, G. and Fréchet, A. (2003). Recovering Canadian Atlantic cod stocks: the shape of things to come. *ICES CM* 2003/U 06, 23pp.
- Ripley, B., and from 1999 to Oct 2002 Michael Lapsley (2009). *RODBC: ODBC Database Access*. <http://CRAN.R-project.org/package=RODBC>, r package version 1.3-1.  
**URL:** <http://CRAN.R-project.org/package=RODBC>

- Ruckstuhl, A., Unternaehrer, T. and Locher, R. (2009). *IDPmisc: Utilities of Institute of Data Analyses and Process Design (www.idp.zhaw.ch)*. <http://CRAN.R-project.org/package=IDPmisc>, r package version 1.1.06.  
**URL:** <http://CRAN.R-project.org/package=IDPmisc>
- Sibert, J., Hampton, J., Kleiber, P. and Maunder, M. (2006). Biomass, Size, and Trophic Status of Top Predators in the Pacific Ocean. *Science (Washington)* 314(5806), 1773–1776.
- Walters, C. and Maguire, J. (1996). Lessons for stock assessment from the northern cod collapse. *Reviews in Fish Biology and Fisheries* 6(2), 125–137.
- Walters, C. (2003). Folly and fantasy in the analysis of spatial catch rate data. *Canadian Journal of Fisheries and Aquatic Sciences* 60(12), 1433–1436.
- Warnes., G.R. (????). *gregmisc: Greg’s Miscellaneous Functions*. R package version 2.1.1.
- Watson, R., Kitchingman, A., Gelchu, A. and Pauly, D. (2004). Mapping global fisheries: sharpening our focus. *Fish and Fisheries* 5, 168–177.
- Wessel, P. and Smith, W.H.F. (1991). Free software helps map and display data. *EOS Transactions of the American Geophysical Union* 72, 441.
- Wilberg, M. and Miller, T. (2007). Comment on ”Impacts of biodiversity loss on ocean ecosystem services”. *Science (Washington)* 316, 1285b.
- Wilson, D., Curtotti, R., Begg, G. and Phillips, K., eds. (2009). *Fishery status reports 2008: status of fish stocks and fisheries managed by the Australian Government*. Bureau of Rural Sciences and Australian Bureau of Agricultural and Resource Economics, Canberra.
- with contributions from Kevin Wright, S.H. and Leidi., A.A. (2010). *doBy: Groupwise computations of summary statistics, general linear contrasts and other utilities*. **http:**

`//CRAN.R-project.org/package=doBy`, r package version 4.0.6.

**URL:** *http://CRAN.R-project.org/package=doBy*

Worm, B., Barbier, E., Beaumont, N. *et al.* (2006). Impacts of Biodiversity Loss on Ocean Ecosystem Services. *Science (Washington)* 314, 787–790.

Worm, B., Hilborn, R., Baum, J.K. *et al.* (2009). Rebuilding global fisheries. *Science (Washington)* 325, 578–585.

## Tables

Table 1: Number of assessments included in the RAM Legacy database

<i>Country/Ocean</i>	<i>Management Body</i>	<i>Acronym</i>	<i>No. stocks</i>
Australia	Australian Fisheries Management Authority	AFMA	16
Multinational	Commission for the Conservation of Antarctic Marine Living Resources	CCAMLR	1
Argentina	Consejo Federal Pesquero	CFP	6
South Africa	South African national management	DETMCM	14
Canada	Department of Fisheries and Oceans	DFO	22
Multinational	Inter-American Tropical Tuna Commission	IATTC	2
Multinational	International Commission for the Conservation of Atlantic Tunas	ICCAT	10
Multinational	International Council for the Exploration of the Sea	ICES	63
Multinational	Indian Ocean Tuna Commission	IOTC	1
Multinational	International Pacific Halibut Commission	IPHC	1
New Zealand	Ministry of Fisheries	MFish	29
Multinational	Northwest Atlantic Fisheries Organization	NAFO	9
USA	National Marine Fisheries Service	NMFS	139
Russia	Russian Federal Fisheries Agency	RFFA	2
Multinational	South Pacific Regional Fisheries Management Organi-	SPRFMO	1

<i>Stock Rank</i>	<i>Stock Number</i>	<i>Species (Common name, Latin name) or higher taxonomic unit</i>	<i>LME</i>	<i>In Database?</i>	<i>Reason if not included</i>
1	1	Peruvian anchoveta, <i>Engraulis ringens</i>	Humboldt Current	no	3
	2	Marine fishes not identified	South China Sea	no	1
	3	Marine fishes not identified	Bay of Bengal	no	1
2	4	Alaska pollock, <i>Theragra chalcogramma</i>	Okhotsk Sea	yes	
3	5	<i>Ammodytes</i>	North Sea	yes	
4	6	Atlantic herring, <i>Clupea harengus</i>	Norwegian Sea	yes	
5	7	Alaska pollock, <i>Theragra chalcogramma</i>	East Bering Sea	yes	
6	8	Capelin, <i>Mallotus villosus</i>	Iceland Shelf/Sea	yes	
7	9	European pilchard, <i>Sardina pilchardus</i>	Canary Current	yes	
8	10	Japanese anchovy, <i>Engraulis japonicus</i>	East China Sea	no	3

<i>Stock Rank</i>	<i>Stock Number</i>	<i>Species (Common name, Latin name) or higher taxonomic unit</i>	<i>LME</i>	<i>In Database?</i>	<i>Reason if not included</i>
9	11	Inca scad, <i>Trachurus murphyi</i>	Humboldt Current	yes	
	12	Marine fishes not identified	East China Sea	no	1
10	13	Gulf menhaden, <i>Brevoortia patronus</i>	Gulf of Mexico	yes	
	14	Marine fishes not identified	Yellow Sea	no	1
	15	Marine fishes not identified	Indonesian Sea	no	1
11	16	Alaska pollock, <i>Theragra chalcogramma</i>	Gulf of Alaska	yes	
12	17	Argentinean short-finned squid, <i>Illex argentinus</i>	Patagonian Shelf	no	1
13	18	Argentine hake, <i>Merluccius hubbsi</i>	Patagonian Shelf	yes	
14	19	Japanese anchovy, <i>Engraulis japonicus</i>	South China Sea	no	1
15	20	Araucanian herring, <i>Strangomera bentincki</i>	Humboldt Current	no	

<i>Stock Rank</i>	<i>Stock Number</i>	<i>Species (Common name, Latin name) or higher taxonomic unit</i>	<i>LME</i>	<i>In Database?</i>	<i>Reason if not included</i>
16	21	Atlantic cod, <i>Gadus morhua</i>	Barents Sea	no	
17	22	European sprat, <i>Sprattus sprattus</i>	Baltic Sea	yes	
18	23	Atlantic herring, <i>Clupea harengus</i>	North Sea	yes	
19	24	Alaska pollock, <i>Theragra chalcogramma</i>	Arctic Ocean	no	
20	25	Marine fishes not identified	Gulf of Thailand	no	
	26	Atlantic herring, <i>Clupea harengus</i>	Baltic Sea	yes	
21	27	Cape horse mackerel, <i>Trachurus capensis</i>	Benguela Current	yes	
22	28	Largehead hairtail, <i>Trichiurus lepturus</i>	East China Sea	no	
23	29	Japanese anchovy, <i>Engraulis japonicus</i>	Yellow Sea	no	

<i>Stock Rank</i>	<i>Stock Number</i>	<i>Species (Common name, Latin name) or higher taxonomic unit</i>	<i>LME</i>	<i>In Database?</i>	<i>Reason if not included</i>
24	30	European anchovy, <i>Engraulis encrasicolus</i>	Black Sea	no	
25	31	Chub mackerel, <i>Scomber japonicus</i>	East China Sea	no	
26	32	Indian oil sardine, <i>Sardinella longiceps</i>	Arabian Sea	no	1
27	33	<i>Decapterus</i>	South China Sea	no	
	34	<i>Sciaenidae</i>	Arabian Sea	no	
	35	Atlantic mackerel, <i>Scomber scombrus</i>	North Sea	yes	
28	36	Largehead hairtail, <i>Trichiurus lepturus</i>	Yellow Sea	no	
29	37	<i>Merluccius</i>	Benguela Current	yes	
	38	Marine fishes not identified	Kuroshio Current	no	
	39	Alaska pollock, <i>Theragra chalcogramma</i>	Sea of Japan	no	



<i>Stock Rank</i>	<i>Stock Number</i>	<i>Species (Common name, Latin name) or higher taxonomic unit</i>	<i>LME</i>	<i>In Database?</i>	<i>Reason if not included</i>
30	40	Round sardinella, <i>Sardinella aurita</i>	Canary Current	no	

Table 2: The world’s forty largest wild-caught fisheries (comprising less than 41% of total global catches, based on average catches 1995-2004 in SAUP database), and the thirty largest fisheries of individual stocks (i.e. fisheries identified to the species level; comprising more than 32% of total global catches), including their LME, whether or not stock assessments for them are included in the RAM Legacy database, and the reason if not included (e.g. 1= no known assessment, 2=assessment is not based on a population dynamics model, 3=assessment inaccessible).

## Figures

### *Figure legends*

Figure 1. Global map of the Large Marine Ecosystems (LMEs) showing the number of stock assessments present in the database for each LME or High Seas area.

Figure 2. Taxonomic coverage of assessed marine species present in the RAM Legacy database. The circle located near the middle of the circular dendrogram represents kingdom Animalia and each subsequent branching represents a different taxonomic group (Kingdom to Phylum to Class to Order to Family to Genus to Species). The width of each line is proportional to the square root of the number of assessments in the database. The outermost lines represent species and the number of lines is the number of assessments for each species. The names of multi-assessment species are not repeated on the outermost portion of the dendrogram but continue counter-clockwise from the first entry. Note that branch lengths are chosen for graphical purposes and do not convey phylogenetic distance.

Figure 3. Comparison of the taxonomic diversity of marine species as provided by FishBase (top panel), the coverage of catch data as provided by the Sea Around Us Project (SAUP) database (middle panel) and the RAM Legacy database (bottom panel). To facilitate the identification of the taxonomic groups that are not presented in the catch and assessment data, the FishBase branching pattern of the spoked dendrogram is maintained to generate other two dendrograms.

Figure 4. Mean trophic level of assessed species for overexploited and non-overexploited stocks. Overexploited stocks are those for which the current biomass is below the MSY BRP.

Figure 5. Temporal coverage of (A) catch/landings, (B) spawning stock biomass and

(C) recruitment. The temporal coverage for individual assessments is represented by thin alternating black and grey horizontal lines in the main panels. Thick horizontal lines at the base of each main panel represent the time periods which are present in 90% (black) and 50% (grey) of all series for that data type. Subfigure histograms contain the frequency of occurrence of the various timespans without reference to time period. Solid and long-dash vertical lines within the subfigures represent the median, 2.5% and 97.5% quantiles, respectively.

Figure 6. Current exploitation rate versus biomass for XXX individual stocks. Exploitation is scaled relative to that which should allow maximum sustainable yield (uMSY); biomass is scaled relative to BMSY. Colors indicate probability of occurrence as revealed by a kernel density smooth function. Solid circles indicate BMSY and uMSY that were obtained directly from assessments; open circles indicate that they were estimated from surplus production models.

Figure 7. Current exploitation rate versus biomass for individual stocks grouped by management unit. Plot details as in Figure 6.





Figure 1:


*Figures*



FishBase



SAUP



## RAM Legacy




Figure 3:




Figure 4:




Figure 5:



all assessments (n=241)




Figure 6:




Figure 7: