# STATUS OF THE BLACK ROCKFISH RESOURCE NORTH OF CAPE FALCON, OREGON TO THE U.S.-CANADIAN BORDER IN 2006

Farron R. Wallace<sup>1</sup>, Yuk Wing Cheng<sup>2</sup>, and Tien-Shui Tsou<sup>2</sup>

Washington Department of Fish and Wildlife
 48 Devonshire Road
 Montesano, Washington 98563

Washington Department of Fish and Wildlife
 600 Capitol Way NE
 Olympia, Washington 98501

# **Executive Summary**

#### Stock

This assessment applies to the Northern portion of the black rockfish (*Sebastes melanops*) stock found between Cape Falcon, Oregon and the U.S. border with Canada. This assessment treats these fish as a separate unit stock and the stock found South of Cape Falcon, Oregon is treated as another unit stock in a different assessment document. Black rockfish are not subjected to a targeted fishery in Canadian coastal waters and are not assessed.

#### **Catches**

Little information exists on the historical landings of black rockfish prior to the early 1960's. Landings of "rockfish" peaked at nearly 25,000 mt in 1945 in support of the war effort, however, there is no known species composition estimates for these catches. Due to the nearshore habitat of this species it is likely that very little of this catch was black rockfish. Predominate harvesters of black rockfish between 1963 and 1983 were commercial line and trawl fishers. Black rockfish trawl landings typically came from directed tows on nearshore rocky reefs and shipwrecks with few landings incidental to other targeted fisheries. Peak landings in the trawl fishery reached 350 mt in 1976 and declined to less than 10 mt in recent years. Black rockfish comprised less than 1% of total rockfish landings by the trawl fishery during this time period.


The "non-trawl" fishery is composed of three distinct line fisheries and each differs in target species. Oregon and Washington fish receiving tickets show nominal rockfish catch as early as 1970 in the salmon troll fishery, during 1973 in the jig fishery and during 1979 in the bottomfish troll fishery. Black rockfish are generally caught as bycatch in the commercial salmon troll fishery; landings peaked in the late 70's (151 mt) and steadily decreased coincident with losses in fishing opportunities for coastal salmon. The bottomfish troll fishery generally targeted lingcod; rockfish landings are small and estimated black rockfish catch never exceeded 2 mt. The jig fishery is primarily composed of small vessels less than 26 feet in length that generally fish near their port of access. Black rockfish were targeted in nearshore areas and are a significant fraction of the nominal rockfish landings in the jig fishery. Black rockfish catch in the jig fishery was inconsequential prior to 1980 and peaked in 1982 at 272 mt. Since 1996 nominal rockfish landings contain no black rockfish due to area restrictions that have forced jig fishers to target other rockfish species found farther offshore.

Black rockfish are the primary target of the coastal groundfish sport fishery with small catches first reported in the late 1970's that steadily increased to over 300 ton per year in the mid 1990's. Due to the implementation of a 10 fish bag limit in 1995 (Figure 7) and longer salmon seasons, annual catch of black rock declined to 188 mt in 2001. In recent years, sport catches increased to more than 300 mt. The coastal recreational rockfish fishery generally competed with sport salmon, halibut and tuna fisheries, and this is reflected in year-to-year variability in black rockfish catch.

Discard of black rockfish in Washington waters in either the commercial or recreational fisheries is likely very small. "Sebastes complex" trip limits in the line fishery were non-restrictive prior to 1999 since few landings ever achieved the trip limit, and there was no incentive to discard catch. Furthermore, Washington State waters (inside 3 miles) have been closed to directed non-trawl commercial fishing since 1996 and directed trawl fishing since 1999. Black rockfish represented only a small fraction of the nominal rockfish catch in the trawl fishery and it is unlikely they were discarded. Discard in the sport fishery is also insignificant since the vast majority of recreational fishers do not high-grade their rockfish catch. This is supported by recent sport fishery information that indicates discard is less than 16 mt on an annual basis.

Recent Black Rockfish Landings From Waters North of Cape Falcon, Oregon to the US-Canadian Border by Gear and Area

		Trawl Ge	ear		Non-	Trawl Gear		Re	ecreational	
	3A	3B	3CS	Total	3A	3B	Total	3A	3B	Total
1995	2.9	0.1	0.3	3.3	2.7	63.1	65.8	209.3	55.5	264.8
1996	0.0	0.0	0.0	0.0	4.8	3.8	8.6	199.7	64.6	264.2
1997	0.7	8.2	0.1	9.0	14.5	0.5	15.0	179.7	54.4	234.1
1998	72.5	0.3	0.3	73.1	0.4	4.5	4.8	195.2	64.2	259.4
1999	0.0	0.0	0.0	0.0	3.4	0.9	4.3	166.0	55.6	221.6
2000	0.0	0.0	0.0	0.0	0.5	0.7	1.2	157.6	67.2	224.8
2001	0.0	0.0	0.0	0.0	0.6	0.5	1.1	133.7	55.0	188.7
2002	0.1	0.1	0.0	0.2	0.4	1.0	1.5	173.0	66.0	238.9
2003	0.1	0.0	0.0	0.1	0.0	0.2	0.2	166.7	70.4	237.1
2004	0.6	0.0	0.0	0.6	0.4	0.3	0.7	173.4	94.6	268.0
2005	0.0	0.0	0.0	0.0	0.3	0.6	0.9	217.5	114.2	331.7
2006	1.2	0.2	0.0	1.4	0.8	0.4	1.2	246.7	74.9	321.5
Total	78	9	1	88	29	77	105	2.218	837	3.055



#### **Data and Assessment**


This portion of the U.S. black rockfish stock was last assessed in 1999 (Wallace et al 1999) with an population dynamics model constructed with AD model builder software (Fournier1997).

The current assessment employed Stock synthesis 2 (SS2V2.00c, compiled 3/27/2006) to model the dynamics of black rockfish population found between Cape Falcon, Oregon and North to the U.S./Canadian border in Coastal Waters. The model was specified to


begin in 1915 to ensure population equilibrium at the start of the modeling time period. Catch data were decayed from the last reliable catch estimates (1965) to 0 by 1940. Fisheries catch, size and age compositions were pooled into three fishery types including trawl, sport and non-trawl. The first size-age compositions were collected in the mid 1970's from the trawl fishery, but samples were not collected on a systematic basis until 1985. Growth (Lmin, Lmax and k) was estimated within the model to account for fishery selection of the larger individual fish at age. The population model was tuned to two fisheries-independent indices that include a tagging CPUE (1986-2007) and a tag abundance biomass index (2000-2007) both derived from WDFW black rockfish tagging information. Both STAT and STAR Panel members agreed that the available fishery dependent indices should not be incorporated due to potential bias resulting from bag limit changes and undocumented measures of fishing effort resulting from changes in search time across the time series.

#### **Unresolved Problems and Major Uncertainties**

Natural mortality is confounded with fishing mortality and is therefore one of the most challenging biological parameters to estimate. It is also one of the most important parameters due to affects on population dynamics including stock rebuild time and the estimation of virgin fishery biomass. In this assessment we explored direct methods to estimate natural mortality and compared to estimates derived from indirect methods (from other biological parameters, e.g., the growth constant and fecundity) in previous assessments. The estimated  $\hat{M}$  derived from direct methods was 0.223 (SE= 0.0071) and 0.272 (SE= 0.061) for males and females, respectively. Given the uncertainties, these estimates compared well with other existing indirect methods. The current base model assumes a female natural mortality rate to be age-specific for females using age at first and full maturity for inflections (10 and 15). A constant natural mortality rate of 0.16 was assumed for males and young females (< 10 years of age) and a rate of 0.2 was assumed for old females (>=15 years of age). This is higher than that used in the 2003 black rockfish assessment off Oregon and California (Ralston and E.J. Dick, 2003) which used a natural mortality of 0.1 and 0.2 for males and old females, respectively. It is apparent from our analysis using both direct and indirect methods that our current assumptions on natural mortality in the base model are within our limits to estimate this parameter and that the low natural mortality rate model is likely too low. Model sensitivity analysis show that other model configurations using higher natural mortality assumptions provide a better overall fit to the data.



Tagging is not incorporated in the model as a tagging experiment, which is not possible within the current SS2 modeling framework. The index for tagging abundance is not fit well and the model estimated effective q for the tagging index was 0.83. This is likely double what it should be based on STAT knowledge of available habitat off the Washington coast. Further, the north central Washington coast, where most of the nearshore rocky habitat exists, is inaccessible to most recreational fishers and is not part of the current tagging program. However, the estimation of q is complicated by the fact that the SS2 value of q is a function of selectivity that is strongly dome shaped for the fishery. Increasing the weighting on survey abundance shows that a better fit to the survey abundance index significantly improves our view of the current population status.



Without an objective evaluation of an informed prior on q it is difficult to compare a prior conception of q based on tagging and the one estimated by SS2. Other issues include the non-independence of the length/age compositions and non-independence of the tagging abundance and CPUE series.

# **Reference Points**

The Pacific Fisheries Management Council recommends that a default target fishing mortality rate of FSPR=0.5 be used for Council managed rockfish species. The current

assessment uses this default for harvest projections for black rockfish and based on the Councils control rule for groundfish would not be considered overfished.

# Model Reference Points

Unfished Stock	Value
Age 3+ Biomass (B <sub>0</sub> ) (mt)	11,390
Spawning Biomass SB( <sub>0</sub> ) (mt)	2,321
SPBio/Recruit (kg/fish)	0.687
Age1 Recruitment (R <sub>0</sub> ) (1,000's)	3,377
Steepness_R0_S0	0.6

Reference points based on


Exploited Stock	<b>Estimated MSY</b>	SB <sub>40%</sub>	SPR (SB <sub>0.5</sub> )
SPR (Spawning Biomass/Recruit)	0.418	0.400	0.40
F (Fishing Mortality Rate)	0.141	0.110	0.110
Exploitation Rate (Yield/Bsmry)	0.081	0.065	0.065
MSY (mt) or MSY proxy (mt)	423	408	408
Yield (mt)	700	928	928
SPBIO/SB(0)	30.1%	40.0%	40.0%
Age 3+ Biomass	5,218	6,264	6,264

#### **Stock Biomass**

The estimated current spawning biomass is 1,239 mt and unexploited spawning biomass is at 2,321 mt resulting in a current stock level that is 53.4% of the unfished. Model results indicate that spawning biomass and age 3+ biomass reached the lowest levels in 1995 following poor recruitment and intense fishing in the late 1980's.

#### Model Results

Year	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Spawning Biomass	707	762	826	891	959	1043	1114	1171	1211	1239
% of Virgin	0.304	0.328	0.356	0.384	0.413	0.449	0.480	0.505	0.522	0.534
Age 3+ Biomass	5977	6066	6147	6516	6739	7405	7485	7470	7564	7558




# Recruitment

Recent increases in biomass are the result of two prominent year classes in 1994 and in 1999. The 1999-year class is estimated to be the largest year class since the beginning of the estimation phase.

# **Model Results**

Year	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Age 1 Recruits (1,000's)	2,339	2,408	2,480	2,548	2,611	2,681	2,735	2,775	2,801	2,819

# ~95% Asymptotic confidence interval




# **Exploitation Status**


Exploitation of black rockfish reached a peak in 1988 of 13% of the Age 3+ biomass and remained near that level for 7 years dropping precipitously between 1995 and 2000. In recent years exploitation has been relatively low (4-6%).

#### Model Results


Recent Trends in black rocktish exploitaion										
Year	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total Exploitation Rate	0.042	0.035	0.027	0.027	0.022	0.028	0.027	0.030	0.037	0.036



Black rockfish stock abundance has been below the Councils' management target has dipped below the Councils' minimum stock size threshold in the last decade. The stock is currently above the management target of B40%.



Exploitation rate relative to spawning biomass indicate that harvest rates exceeded management targets between the mid 1980's through the mid 1990's.



#### **Management Performance**

Harvest has remained well below the harvest guideline of 517 mt (1997-1999) and 577 mt (+- 2CV's 523-632 mt's) that was established by the Council following the 1994 (Wallace et al., 1994) and the 1999 assessment (Wallace et al., 1999), respectively. The 1999 assessment estimated the 2001 spawning biomass of 646 mt (+- 2CV's 601-687 mt's) with an equilibrium spawning biomass of 451 mt (+- 2CV's 401-501 mt's) equating to a 2001 SB<sub>2001</sub>/SB<sub>Equil</sub> of 143%. The catch time series includes discard when existing, ABC is constant and changes in spawning biomass across the time series is not available.

# **Total Black Rockfish Catch By all Fisheries**

Year	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total Catch	312.0	263.7	222.8	225.9	190.4	239.2	238.4	268.9	334.3	323.3

#### **Forecasts**

Projections of future catches were based on a  $F_{SPR 50\%}$  rate of fishing mortality. We also assumed that the sport fishery would account for 100% of the catch and that selectivity would remain unchanged from that estimated within the model in the final year.

Year	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ABC (mt)	535	521	505	478	459	448	443	441	442	443
Spawning Biomass	1281	1317	1328	1270	1204	1140	1087	1048	1023	1010
% of Virgin	0.552	0.567	0.572	0.547	0.519	0.491	0.468	0.452	0.441	0.435

# **Decision Table**

Decision matrix based on a range of natural mortality rates where columns represent results from the assumed natural mortality model given catch rates that resulted from alternative states of nature (rows).

# 2007 Northern Black Rockfish Stock Assessment

2007 1	101 11101		M= 0.12 Males	SII SIUCK AS	<i>M</i> = 0.16 <i>Males</i>		M= 0.19 Males		
			M= 0.18 Femal	es	M= 0.24 Female	s	M= 0.285 Femal	les	
State of				al Mortality 1/	Base N		High Natural Mortality <sup>1/</sup>		
Nature	Year	ABC	l .	Depletion	SpawnBio	Depletion	SpawnBio	Depletion	
	2007 /2	140	•	18.2%	1281	55.2%	2571	86.5%	
	2008 /2	128		17.0%	1317	56.7%	2640	88.9%	
	2009	118		15.4%	1328	57.2%	2660	89.5%	
<del>5</del>	2010	132		16.7%	1368	58.9%	2678	90.2%	
Low Catch	2011	146		18.1%	1389	59.9%	2655	89.4%	
≥	2012	158	426	19.7%	1401	60.4%	2610	87.9%	
Ľ	2013	169	463	21.4%	1409	60.7%	2560	86.2%	
	2014	178	500	23.1%	1419	61.2%	2518	84.8%	
	2015	184	534	24.7%	1433	61.8%	2489	83.8%	
	2016	188	563	26.0%	1452	62.6%	2478	83.4%	
	2007 /2	535	393	18.2%	1281	55.2%	2571	86.5%	
_	2008 /2	521	368	17.0%	1317	56.7%	2640	88.9%	
Base Model Catch	2009	505	l .	15.4%	1328	57.2%	2660	89.5%	
రొ	2010	478	l .	11.7%	1270	54.7%	2582	86.9%	
<del> </del>	2011	459	185	8.6%	1204	51.9%	2475	83.3%	
§	2012	448	133	6.1%	1140	49.1%	2362	79.5%	
Se	2013	443	94	4.3%	1087	46.8%	2260	76.1%	
Ва	2014	441	l .	rash	1048	45.2%	2178	73.3%	
	2015	442	l .	rash	1023	44.1%	2121	71.4%	
	2016	443		rash	1010	43.5%	2087	70.3%	
	2007 /2	1190		18.2%	1281	55.2%	2571	86.5%	
	2008 /2	1153	l .	17.0%	1317	56.7%	2640	88.9%	
_	2009	1109	334	15.4%	1328	57.2%	2660	89.5%	
달	2010	1002		rash	1117	48.1%	2432	81.9%	
రొ	2011	923		ash	920	39.7%	2200	74.1%	
High Catch	2012	869		ash	754	32.5%	1989	67.0%	
Ī	2013	834		ash	624	26.9%	1815	61.1%	
	2014	814		ash	527	22.7%	1682	56.6%	
	2015	802	l .	ash	459	19.8%	1590	53.5%	
	2016	794	Cı	ash	411	17.7%	1531	51.5%	

<sup>1/</sup> The natural mortality rate of "young" females <= 10 years of age and males are equal. The natural mortality rate for "old" females between the ages of 11 and 15 is linearly increasing and then remains at the constant rate listed above.

#### **Research and Data Needs**

In order to objectively evaluate a prior on q for the tagging, information on habitat distribution within the stock boundary is necessary. A nearshore assessment should be completed using side-scan, backscatter and multi beam methods. This has already been completed for some portions of the coast and new information can be integrated.

# **Rebuilding Projections**

None required.

# **Regional Management Concerns**

Black rockfish is highly resident to specific reefs and are therefore susceptible to localized depletion especially during times of population decline. Because of this, relatively higher levels of abundance may be needed to meet recreational fishery objectives. For example, the recreational fishery industries need to maintain a sufficient success rate to be economically feasible.

#### 1.0 INTRODUCTION

The status of stocks for the "Northern" black rockfish stock found between Cape Falcon, Oregon and the U.S. Canadian border was last determined in 1999 (Wallace et al, 1999). The population was assessed using an AD model configuration where tag recovery was modeled explicitly. The population was regarded as healthy, stock abundance was estimated to be slightly increasing after passing through a low in the late 1980's and early 1990's. The recommended allowable annual yield was 577 mt based on an F45% exploitation strategy and a tag recovery rate of 50%. The estimated stock biomass ranged between 9,500-10,100 mt, depending on assumptions on tag reporting rates. The current analysis reprises estimates based on the 1999 model that uses an improved stock synthesis program (SS2) (Methot, 2006) and presents a completely new model specification. This assessment is distinguished from other more southerly black rockfish population assessment(s) by identifying it as the "northern" stock. However, we have no indication that there is any stock divide at the U.S.-Canadian border just that this assessment includes information only as far north as the U.S.-Canadian border.

Throughout the document we include model results that are based on both the "STAT best fit" model and the STAR base model. STAT best fit model natural mortality for "old" females is assumed to be 0.24 versus 0.20 in the STAR base model and all other parameter settings remain the same. Results in the STAT best fit model are based largely on new and expanded analyses following conclusion of the STAR Panel. The final base model recommended by the SSC and used for management is based upon the STAT best fit model.

# 1.1 Species Distribution, Stock Structure, and Management Units

Black rockfish (*Sebastes melanops*) are widely distributed along the Pacific coast from central California to the Gulf of Alaska inhabiting nearshore areas at bottom depths of less than 50 fathoms (Miller and Lea, 1972). Adults are schooling and associated with irregular, rocky bottom or underwater structures, though at times may be found actively feeding on the surface.

Washington tagging data suggest that Cape Flattery and Cape Falcon may represent area bounds for a coastal Washington-northern Oregon black rockfish stock. From over 54,000 tag releases in this area, no fish were recovered north of the Strait of Juan de Fuca and only 6 were recovered south of Cape Falcon in the 15-year study (Figure 1). To corroborate these results, a genetic stock identification study of coastal black rockfish populations was conducted from 1995-1997 {}(WDFW report in progress). Horizontal starch-gel electrophoresis was used to examine 10 black rockfish collections from northern California, Oregon, Washington and southern British Columbia. Significant heterogeneity occurred among Oregon collections, while less heterogeneity was found among Washington collections. Dendrogram and multidimensional scaling (MDS) analysis of genetic distances (Nei, 1978) revealed three major geographical groupings (Figure 2). The groups include samples from: 1) north of Cape Falcon, 2) south of Cape Falcon off the Oregon coast, and 3) a single collection from northern California (Port

Albion). The study concluded that there is an apparent large-scale geographical clustering of coastal black rockfish populations and there does not appear to be any geographical pattern to clustering of populations within each group. For this assessment, we assume that black rockfish distributed between Cape Falcon, Oregon and Cape Flattery, Washington represent a unit stock. All biological parameters, data analysis and yield projections presented in this assessment are intended to describe this portion of black rockfish coast-wide distribution.

It is interesting to note that although no black rockfish tags were recovered from southern British Columbia during the 15 year tagging study, fish collected just 20 km north of Cape Flattery in Barkley Sound, B.C. were genetically similar to the coastal Washington collections. The lack of recoveries from acoss the Strait of Juan de Fuca is likely due to a lack of any target fisheries in coastal B.C. waters or may indicate that the Strait provides an effective physical boundary, which few if any adult black rockfish will cross. Nearshore and oceanic drift likely influence gene flow during the three to four month planktonic stages. Survival during the early life stages is strongly influenced by oceanic processes and recruitment may be dependent upon the health of black rockfish populations both north and south of the Strait of Juan de Fuca.

# 1.2 Life History Overview

Like many rockfish species, black rockfish are slow growing, long lived and mature late in life. Black rockfish are recruited into the commercial and sport fishery at 4 years of age; age composition of the catch can span three decades. Early recruitment, delayed maturity and schooling behavior make black rockfish susceptible to over-exploitation. Furthermore, WDFW found evidence that, in at least one year, a number (approximately 10%) of mature females examined during parturition did not spawn during year of collection. Ovarian characteristics derived from histological preparations on these specimens indicated that although they had spawned in prior seasons, they had not advanced beyond the early yolk accumulation stage and were re-sorbing their oocytes. Thus, some fraction of the mature population may not spawn annually. If this behavior were common from year to year production would be accordingly reduced.

Another important aspect of black rockfish life history is differences in growth and apparent natural mortality rates between sexes. Composition sampling data show that the sex ratio before age 10 is nearly equal and then the percent female declines sharply thereafter (Figure 3). For the purposes of this assessment we interpret the loss of females due to increased natural mortality at age, which coincides with female transition into sexual maturity.

# 1.3 Review of Fishery

Recreational and commercial fishers have harvested black rockfish in nearshore areas off the Washington coast since the early 1960's. Commercial fisheries include salmon and bottomfish troll, jig and groundfish trawl. The recreational fishery is divided between charter and private boat operations. Due to restrictive regulations black rockfish landings

have steadily declined for commercial fisheries since the mid 1980's. Recreational landings peaked in the late 1980's and declined slightly in the 1990's and have increased slightly in the most recent years (Table 1 and Figure 4).

#### 1.3.1 Catch

Little information exists on the historical landings of black rockfish prior to the early 1960's. The first black rockfish catch of 151.5 mt was recorded in 1952 for trawl gear. Landings of rockfish peaked at nearly 25,000 mt in 1945 in support of the war effort, however, there is no known species composition estimates for these catches (Table 2). Due to the nearshore habitat of this species it is likely that very little of this catch was black rockfish. Catches prior to known estimates were decayed to zero back to 1940 within the model and these catches are presented in Table 3.

Predominate harvesters of black rockfish between 1963 and 1983 were commercial line and trawl fishers. Black rockfish trawl landings typically came from directed tows on nearshore rocky reefs and shipwrecks with few landings incidental to other targeted fisheries. Catch information has been updated since the 1999 assessment to reflect changes in species composition estimates derived from port sampling. These changes resulted in a slightly lower catch during the early part of the time series (Figure 5). Peak landings in the trawl fishery reached 350 mt in 1976 and declined to less than 10 mt in recent years due to area and catch restrictions (Figures 6-8).

The "non-trawl" fishery is composed of three distinct line fisheries and each differs in target species. Oregon and Washington fish receiving tickets show nominal rockfish catch as early as 1970 in the salmon troll fishery, during 1973 in the jig fishery and during 1979 in the bottomfish troll fishery. Black rockfish are generally caught as bycatch in the commercial salmon troll fishery; landings peaked in the late 70's (151 mt) and steadily decreased coincident with losses in fishing opportunities for coastal salmon. The bottomfish troll fishery generally targeted lingcod; rockfish landings are small and estimated black rockfish catch never exceeded 2 mt. The jig fishery is primarily composed of small vessels less than 26 feet in length that generally fish near their port of access. Black rockfish were targeted in nearshore areas and are a significant fraction of the nominal rockfish landings in the jig fishery. Black rockfish catch in the jig fishery was inconsequential prior to 1980 and peaked in 1982 at 272 mt. Since 1996 nominal rockfish landings contain no black rockfish due to area restrictions that have forced jig fishers to target other rockfish species found farther offshore.

Black rockfish have become the primary target of the coastal groundfish sport fishery since the mid 1980's (Table 1 and Figure 4). Small black rockfish catches were reported in the late 1970's and steadily increased to over 300 ton per year in the mid 1990's. Due to the implementation of a 10 fish bag limit in 1995 (Figure 7) and longer salmon seasons, annual catch of black rock declined to 188 mt in 2001. In recent years, sport catches increased to more than 300 mt. The coastal recreational rockfish fishery generally competed with sport salmon, halibut and tuna fisheries, and this is reflected in year-to-year variability in black rockfish catch.

#### 1.3.2 Discard

Discard of black rockfish in Washington waters in either the commercial or recreational fisheries is likely very small. "Sebastes complex" trip limits in the line fishery were non-restrictive prior to 1999 since few landings ever achieved the trip limit, and there was no incentive to discard catch. Furthermore, Washington State waters (inside 3 miles) have been closed to directed commercial fishing since 1995. Black rockfish represented only a small fraction of the nominal rockfish catch in the trawl fishery and it is unlikely they were discarded. Discard in the sport fishery is also insignificant since the vast majority of recreational fishers do not high-grade their rockfish catch. This is supported by recent sport fishery information that indicates discard is less than 16 mt on an annual basis (Table 4).

#### 1.3.3 Effort

Coastal Washington recreational effort has steadily increased since the early 1980's with some declines in the late 1990's (Table 5). Increase in the popularity of bottomfish fishing has been coincident with loss of salmon fishing opportunities and a genuine increase in public interest in recreational groundfish fishing. Though a multiple target strategy may be used by sport fishermen, the bottomfish-only trips consisted about 15%-20% of the total activities in the Washington recreational fisheries.

# 1.4 Fishery Management

#### 1.4.1 ABC/HG and Management Performance

The black rockfish resource was first assessed in 1994 (Wallace and Tagart, 1994). Estimated biomass declined to 60% and female egg production decreased to 43% of the unfished level. The 1995 forecasted yield ( $F_{45\%}$ ) and harvest guideline (HG) for combined fisheries was 517 mt. Black rockfish harvest has remained below the HG at 298, 244, 242 and 309 mt for 1995, 1996 1997 and 1998, respectively. Harvest has also remained well below the harvest guideline of 577 mt that was established by the Council following the 1999 assessment (Wallace et al., 1999).

#### 1.4.2 Review of Regulatory Changes

In recognition of the recreational fishery dependence on black rockfish and to address concerns over localized declines in availability, state and federal regulations have significantly restricted commercial and recreational harvest over the last decade (Figures 5-7). In 1992, the recreational bag limit was reduced from 15 to 12 rockfish off most of Washington, and commercial line fisheries were limited to 100 lbs of black rockfish or 30% of total catch on board except when fishing in the area between Destruction Island and Cape Alava or south of Leadbetter Point. The area restrictions were intended to reduce commercial harvest of black rockfish in areas heavily utilized by recreational fishers. WDFW imposed further restrictions in 1995 that prohibit commercial line harvest (except for bycatch in the salmon troll fishery) inside state waters, imposed trawl gear restrictions and reduced the recreational bag limit to 10 fish. These regulations are still in effect today.

# 1.5 Sampling Regime

Oregon and Washington routinely collect commercial and sport groundfish biological samples at various ports of landing (Tables 6 and 7). ODFW sampling is stratified by port complex, gear, market category, and quarter and generally follow methodology describe by Sen (1984). Oregon samples of interest for this assessment include only those samples collected from the sport fishery fishing north of Cape Falcon and landing into Garibaldi. WDFW black rockfish age composition sampling is stratified by time (year) and area (3B and 3A). Washington samples are collected from the trawl fishery throughout the year, and between March and October for the sport and commercial line fisheries concurring with the spring to fall fishing season.

Both Oregon and Washington regularly collect species composition samples for mixed rockfish market categories in the trawl fishery. Samples are used to derive catch estimates for various species including black rockfish and are available from PacFIN. WDFW periodically collected species composition samples from nominal rockfish landings in the commercial line fishery and these were used to estimate black rockfish catch in mixed rockfish categories.

#### 2.0 Data

#### 2.1 Catch

Black rockfish catch data are compiled from a variety of sources including PacFIN, agency reports, fish ticket information and communication with agency personnel. Rockfish landings from the domestic trawl fishery are routinely sampled for species composition by coastal port samplers. Revised estimates of catch for Washington and Oregon were obtained from PacFIN, fish tickets, and species composition sampling in coastal ports in Oregon and Washington (Tables 1-2). Revised catch estimates were slightly smaller in most years prior to 1983 (Figure 8).

Estimates of Washington coastal sport catch and effort is produced from creel and exit count data collected by WDFW's Ocean Sampling Program (OSP). WDFW instituted the OSP in the 70's to estimate catch. The program was later refined to provide necessary information to meet the goals of the Magnuson Fishery Conservation and Management Act of 1976. Estimation procedures for sport groundfish landings are not well documented in earlier years, but species-specific catches were reported in a series of WDFW technical publications since the 1970's. Lai, et al. (1991) describes estimation methodologies beginning in the late 1980's. Variance estimates for catch are available since 1990. Black rockfish discard data in sport fisheries are available since 2002 (Table 3). Proportion-at-size and proportion-at-age by sex and fisheries where derived from biological samples collected from coastal Washington and Oregon landings north of Cape Falcon (Figures 9 and 10).

# 2.2 Biology

# 2.2.1 Sampling

Biological sampling of fisheries for black rockfish age and length compositions goes back as far as 1976 in the trawl fishery. Coverage of the commercial fisheries in the last 10 years is nil due to restrictive management. The sport fishery has been relatively well sampled over the last two decades (Tables 6 and 7).

# 2.2.2 Length weight relationship

Random samples were collected in 1984 (1,157 fish) and from 1988-2001 (1,397 fish), with fork length (cm) and weight (kg) measurements. We modeled the length weight relationship as  $W = aL^b + \varepsilon$ , where W and L were the weight and fork length,  $\varepsilon \sim N(0,\sigma^2)$  and the parameters a and b were to be determined. For male black rockfish,  $\hat{a}$  and  $\hat{b}$  were 3.796x10<sup>-5</sup> kg cm<sup>-2.782</sup> (3.303x10<sup>-6</sup> kg cm<sup>-2.782</sup>) and 2.782 (SE=0.02309), respectively (Figure 11). For female black rockfish,  $\hat{a}$  and  $\hat{b}$  were 4.030x10<sup>-5</sup> kg cm<sup>-2.768</sup> (3.334x10<sup>-6</sup>) and 2.768 (SE=0.02188), respectively (Figure 9). There was no statistical difference (P>0.05) between the male and female length weight relationships.

#### 2.2.2 Growth

Random samples of black rockfish (14,919 male, 12,304 female, and 213 of unknown sex) were collected in 1984 and from 1988-2006 with age and fork length measurements. Most of the fish with unknown sex were juveniles with the smallest age equal to one. , The Schnute three parameter von Bertalanffy growth model was used to model growth, with the assumption of no variation in growth among years. The growth equation is  $l_t = L_{\infty} + (L_1 - L_{\infty})(1 - e^{-K(t-1)}) + \varepsilon$ ,

where  $l_t$  is the fork length at age t,  $L_1$ ,  $L_\infty$  and K are unknown to be determined,  $\varepsilon \sim N(0, \sigma^2)$ .  $L_1$  is the length at age one;  $L_\infty$  and K are von Bertalanffy growth parameters, limited size and growth constant.

Due to the restriction of  $L_1$ , which was assumed to be the same in male and female fish, we proposed the use of a dummy variable in the Schnute three parameters growth model. The use of dummy variable was to test the growth difference between male and female fish. The proposed model was

$$l_{t} = L_{\infty} + D_{L}z + (L_{1} - L_{\infty} - D_{L}z)(1 - e^{-(K + D_{k}z)(t - 1)}) + \varepsilon,$$

where z was a dummy variable (male =0, female =1),  $D_L$  and  $D_k$  were two additional unknown variables to be determined. In this model, males and females have the same growth curves before age 1.

The parameters  $\hat{L}_{\infty}$ ,  $\hat{K}$ ,  $\hat{L}_1$ ,  $\hat{D}_L$  and  $\hat{D}_K$  were 46.370 cm (SE=0.215 cm), 0.194/yr (SE=0.00628/yr), 20.123 cm (SE=0.583 cm), 3.903 cm (SE=0.347 cm) and -0.0299/yr (SE=0.00472/yr), respectively. In Figure 12, there are plots of the expected age fork length relationships of male and female black rockfish. Both  $\hat{D}_L$  and  $\hat{D}_K$  were highly significant (P<0.001).; this implied the expected  $\hat{L}_{\infty}$  and  $\hat{K}$  were different between the sexes. The expected limited size of male rockfish and the growth constant were 46.370 cm and 0.194/yr. The expected limited size of female rockfish and the growth constant were 50.273 cm and 0.164/year. The estimated expected limited sizes of male and female black rockfish were similar to the expected limited sizes of male (46.611 cm) and female (51.225 cm) estimated by the capture-recapture data with time at large and size measurements but not the growth constants. The difference in the growth constants estimation might be due to the assumption of age at zero and the aging of fish with an integer scale.

#### 2.2.3 Aging error

Since 1992, 3,147 black rockfish were sequentially selected and aged with two age readers independently. We modeled the aging error with a simple regression with no intercept. The estimated slope was 0.9977 (s.e.=0.001858). The CV of the aging error was small (0.18%). Figure 13, shows a scatter plot of the age data from the two readers. Figure 14 shows the between reader age specific variation that was used for data input in the SS2 stock assessment.

# 2.2.4 Age weight conversion errors

There were aging errors, age to length conversion errors, and length weight conversion errors in age to weight conversion:

$$W = a[(L_{\infty} + D_L z + (L_1 - L_{\infty} - D_L z)(1 - e^{-(K + D_k z)(t + 1)})]^b.$$

We assumed all these errors were independently normally distributed. The Delta method was employed to estimate the overall standard errors. The estimated male and female black rockfish age to weight and standard errors are presented in Table 8.

# 2.2.5 Age and fork length sex maturity relationship

A random sample of 352 female black rockfish captured in 1998 were selected for the estimation of black rockfish maturity (Table 9). A generalized linear model with a binomial (logit link) was used to model the age of 50% maturity. Bootstrapping was used to estimate the 95% confidence intervals of the age of 50% maturity. The estimated age of 50% maturity was 10.31 year and the 95% confidence intervals by bootstrapping was (9.72 year, 11.24 year). The estimated probability of maturity with age was

$$\hat{\pi} = \frac{e^{-7.13 + 0.69t}}{1 + e^{-7.13 + 0.69t}}.$$

The estimated probability of sex maturity curve with age for females is plotted in Figure. 15. Female black with fork length (l) 25-49 cm captured in 1998 (391 fish) were randomly selected for the estimation of maturity of rockfish (Table 10). The estimated

length of 50% was 42.15 cm and the 95% confidence intervals by bootstrapping was (41.49 cm, 42.87 cm). The estimated probability of maturity with fork length was

$$\hat{\pi} = \frac{e^{-17.05 + 0.40l}}{1 + e^{-17.05 + 0.40l}}.$$

The estimated probability of sex maturity curve with fork length is plotted in Figure. 16.

Fecundity estimates are based on 47 mature black rockfish ovaries collected during parturition between 1989 and 1991 off the central Washington coast. Estimated fecundity ranged from 117,550 eggs for a 37 mm fish to 1.2 million eggs for a 490 mm fish. Fecundity at a mean size of 41 cm is 544,528. There is a significant relationship between fecundity and length (M, E+6larvae/cm) = 0.0634L-2.0586 and fecundity and weight (M, E+6larvae/kg) = 0.7674W-0.3657 (Figure 17). Fecundity at weight parameters are provided as data input to synthesis and since larval output are in 1.0E+6 units, spawning biomass from the model should be multiplied by 106 to obtain the absolute spawning output. An increasing larval output by older, larger fish has a significant impact on the population dynamics such that a lightly exploited population with and age structure shifted towards older fish, would have greater spawning potential than a population shifted towards younger fish even if the biomass of spawning females were the same. This effect is significantly amplified in the black rockfish populations because it appears that larvae from larger, older black rockfish appear to be more viable than those from younger fish (Berkeley, 2004). This further implies that maintaining a black rockfish population that preserves the older segment of the population may be very important for reproductive success of this species.

# 2.2.6 Total mortality

The mortality model we used assumes direct density dependence. If the population at time t is N(t), then

$$\frac{dN(t)}{dt} = -ZN(t) ,$$

where M is the termed the instantaneous coefficient of total mortality. This model is popular for fish stock assessment because it is simple, because data are usually not available to support more complex representations, and because it often makes reasonable assumptions for the exploited age classes. The population size at time t is

$$N(t) = N(0)e^{-Zt}$$
.

We assume that fishing mortality (F) and natural mortality (M) sum to total mortality (Z), where Z=F+M.

Taking the log of both sides of the equation, we get

$$\log(N(t)) = \log(N(0)) - Zt.$$

For the rockfish length frequency composition data, we need to convert the fork length into age. The inverse von Bertalanffy growth equation is

$$t(L) = t_0 - \frac{\log(1 - \frac{L}{L_{\infty}})}{K}$$
. We set  $t_0 = 0$  for simplicity. Now,

$$\log(\frac{N(t)}{t(L+\Delta L)-t(L-\Delta L)}) = \log(\frac{N(0)}{2(t(L+\Delta L)-t(L))}) - Zt(L).$$

The length interval for the frequency data is 3 cm. Assuming there is error in the data, we can fit a regression line with

$$y = \log(\frac{N(t)}{t(L + \Delta L) - t(L - \Delta L)})$$
 and  $x = t(L)$  with  $\Delta L = 1.5$  cm.

The above method is equivalent to the method of Pauly (1983).. who derived it by using the Baranov catch equation,

$$C(t_1, t_2) = N(t_1) \frac{F}{Z} \{1 - \exp[-Z(t_2 - t_1)]\},$$

where  $C(t_1, t_2)$  is the total between age class  $t_1$  and  $t_2$ . He approximated part of the catch equation

$$\log C(t_1, t_2) = d - Zt_1 + \log\{1 - \exp[-Z(t_2 - t_1)]\}\$$

with  $\log(1 - \exp(-\Delta t)) \approx \log(\Delta t) - \frac{\Delta t}{2}$ . Both results are similar.

Black rockfish length frequency data have been collected from port sampling and recreational surveys since 1984. Both male and female black rockfish length frequency data show peaks near 36 cm, presumably due to fishery selectivity. Thus, for the purposes of this analysis, black rockfish with size greater than 36 cm were used to estimate the total mortality. We estimated the total mortalities of black rockfish by sex. The estimated male and female total mortality coefficients from 1984 to 2006 and number of samples are listed in Table 11. Plot of expected male and female estimated total mortality coefficients against total fishing effort are shown in Figure 18. The estimated intercept (~0.2 for males and ~0.26 for females) in each sub graph is the estimated natural mortality (where effort=0) using the mortality model described above and assuming direct density dependence. The estimated female total mortality coefficients were greater than the estimated male total mortality coefficients from 1984 to 2006 and beginning in 2000 there was a decreasing trend observed in both male and female black rockfish total mortality (Figure 19).

# 2.2.7 Natural mortality

Fish natural mortality is confounded with fishing mortality, so it is one of the most challenging fish biological parameters to be estimated. It significantly affects the stock rebuild time and the estimation of virgin fishery biomass. There are both direct and indirect methods to estimate the natural mortality of fish species. Indirect methods are derived from other biological parameters, eg., the growth constant and fecundity (Wallace et.al., 1994 and Wallace et.al., 1999). It is difficult to estimate the uncertainties from indirect methods.

In this assessment, we attempted to estimate the natural mortality of black rockfish with a direct method. We assumed F = qE, where q was catchability coefficient and E was fishing effort. Natural mortality could be estimated with the relationship

$$Z = F + M$$
.

After 1995, the bag limit for recreational catch dropped to 10; thus, we only included the recreational rockfish trip effort (fish/angler) and the total catch in this analysis. We assumed constant M with annual variation and the total fishing effort at year t would result in the total mortality in year t+1. The proposed model was

$$Z_{\scriptscriptstyle t+1} = qE_{\scriptscriptstyle t} + M + \varepsilon_{\scriptscriptstyle t}\,,$$

where  $\varepsilon_r \sim \text{NID}(0, \sigma^2)$ , q and M were the unknown to be determined.

Plot of expected male and female estimated total mortality coefficients against total fishing effort where the intercept was the estimated natural mortality is shown in Figure 18. The estimated linear relationship between male and female black rockfish is shown in Figure 19 and a time series plot of the estimated male and female black rockfish total mortalities is shown in Figure 20. The estimated  $\hat{M}$  of male and female black rockfish were 0.223 (SE= 0.0071) and 0.272 (SE= 0.061). The relationship of  $\hat{M} \approx \hat{K}$  was observed in male black rockfish, while  $M=1.6\hat{K}$  was observed in female black rockfish. All these values agreed with other existing indirect methods.

#### 2.3 Abundance Indices

#### 2.3.1 Bottom trawl surveys

The NMFS has conducted the West Coast triennial bottom trawl survey of groundfish resources since 1977. Survey depth range in most years has been from 30-200 fm (Wilkins et.al, 1995). This is outside the normal depth range of black rockfish and only 233 fish in 27 tows have been captured to date. Therefore, we incorporated no triennial trawl survey data into this assessment.

#### 2.3.2 Recreational CPUE

Abundance indices are assumed to be proportional to population abundance. The catchability coefficient (Q) is the factor that relates the units of the index to the abundance of the population. Random variability in the coefficient may occur, but if there is a trend over time or if the coefficient varies with population abundance, then the assessment may be biased. Sport fishery catch rates will be influenced by undocumented

search time at sea, and the observed decline in CPUE indices would be underestimated. There is no information to evaluate annual differences in effort for specific individual target species such as black rockfish. April-September estimates of catch and effort (by trip type) for the sport fishery from coastal Washington ports are available from the WDFW Ocean Sampling Program since 1990. Black rockfish abundance trends were explored using methods described below, but not used in the current assessment due to 1) changes in bag limits, 2) a switch to bait in the early –mid 1990's, and 3) a bag limit that may have capped the trend since the mid-late 1990's that may have biased the population trend.

#### **Delta lognormal model**

A delta lognormal model (Lo et al. 1992) has been commonly used for modeling the abundance of marine species from trawling data. It uses generalized linear models GLMs in both stages, where  $P_{ij}$  is the probability of abundance of observation j in year i and  $C_{ij}$  is the catch per unit effort (CPUE). CPUE can be catch per angler hr, catch per trip, or catch per angler. The distribution of  $C_{ij} > 0$  usually follows a lognormal distribution.

The distribution of  $P_i$  follows a binomial distribution. The modeling of  $P_{ij}$  and  $C_{ij}$  through a two stage process with other predictor variables is commonly called delta lognormal model (Lo et al., 1992). This approach affords the opportunity to investigate the probability of abundance on a spatial scale with other predictor variables, which include both geographical information and environmental variables. Problems associated with zero values in catch data can be avoided by using the delta lognormal model, which only fits the positive catch data. There is, however, a possible bias induced by using a two stage model process. Lo et. al. (1992) and Syrjala (2000) attempted to estimate the bias of the estimated variance in this model using both simulation and approximation techniques. Both  $P_{ij}$  and  $C_{ij}$  do not assume normally distributed (binomial, lognormal) in the two stages model process and there is possible correlation between them. Also, the use of the delta lognormal method to estimate the variance of the final estimate is questionable. This can be overcome by non-parametric bootstrapping.

#### First stage model

The response variable  $P_{ij}$  is a Bernoulli component (presence-absence) of CPUE j in year i. The choice of the logit link function is standard (McCullagh and Nelder, 1989; Cheng and Gallinat, 2004). The link function is

$$g(P_{ij}) = \log(\frac{P_{ij}}{1 - P_{ij}}) = x_i,$$

where  $x_i$  is a factor variable (annual effect).

# Second stage model

We model  $C_{ij} > 0$  in terms of the covariates  $x_{ij}$ . It is a truncated Poisson distribution.

# Bootstrapping method and non-parametric coefficient of variation

The nonparametric bootstrap method (Efron 1982; Hall 1992; Jackson and Cheng 2001) was used to estimate the 95% confidence intervals for the mean CPUE in both the mean estimates and the estimates which resulted from the delta lognormal model. Due to the computational intensity required when applying GLMs and a large data set, K = 200 to 1000 samples have been used. We have rerun the bootstrapping three times and compared the precision of the estimates of the 2.5%, 15.87%, 84.13%, and 97.5% quantiles. The estimates of the quantiles are correct to the first 3 significant places due to the very large dataset. The coefficient of variation of a data X,

$$CV_X = \frac{\sigma_X}{\mu_X} \approx \frac{\hat{\sigma}_X}{\overline{X}},$$

is commonly used to describe the variation (one standard deviation) of the data compared with the mean of the data. The parameters  $\sigma_X$  and  $\hat{\sigma}_X$  are population X standard deviation and estimate population X standard deviation. Letting  $q_{X,0.025}$  be the 2.5% quantile of data X, we define the ad hoc CV for the non-normal distribution as

$$CV_X = \frac{q_{X,0.8413} - q_{X,0.1587}}{2\mu_X} \approx \frac{\hat{q}_{X,0.8413} - \hat{q}_{X,0.1587}}{2\overline{X}} \ .$$

For the sample mean, we use

$$CV_{\overline{X}} = \frac{q_{X,0.8413-}q_{X,0.1587}}{2\sqrt{n}\mu_{Y}} \approx \frac{\hat{q}_{X,0.8413-}\hat{q}_{X,0.1587}}{2\sqrt{n}\overline{X}},$$

where n is the sample mean.

The sample mean of the CPUE in each year was compared with delta lognormal model results. Black rockfish length frequency data have been collected since 1990 in both recreational and commercial fisheries. Plots of the estimated recreational fishery CPUEs from mean estimators and the delta lognormal model for all areas combined is shown in Figure 21 and Figure 22 shows results from Area 2 only, A summary of the number of recreational data recorded in all areas (Areas 1,2 3, 74 and 84) and the proportion of these from 1990 to 2006 is given in Table 12. Area 2 was the major fishing area and the fishing effort was roughly proportional to the catch. Area 2 was the major rockfish area. Tables 13 and 14 provide summaries of the estimated CPUEs from the mean estimator and the delta lognormal model for all areas combined and area 2, respectively. Undoubtedly, Area 2 had a higher CPUEs compared with the other areas. Although the bag limit changed from 15, to 12 to 10 during 1990 to 1995, the estimated CPUEs reflected the changes from 1990 to 1993. From 1995 onwards, there was an increasing trend in CPUEs with a constant bag limit (Figures 19-20).

# 2.3.3 Tagging CPUE

Since the start of the coastal Washington black rockfish tagging program in 1981 information on catch and rod hours have been recorded. These data represent the total number of fish caught and angler hours at each specific fishing location during a trip. The number of fish tagged (and released) was typically less because of mortality from hooking or barotraumas. The tag CPUE in the current assessment represents the mean annual CPUE across all trips (by year) for the Central Washington Coast between Grays Harbor and Sea Lion Rock since 1985 (Table 15 and Figure 24).

# 2.3.4 Mark-recapture tagging study

From 1981 to 1990 and resuming again in 1998, black rockfish has been the subject of multiple tagging experiments along the coast of Northern Oregon and Washington Since 1998, internally implanted coded wire tags (CWT) were employed to ensure that tag recovery was not dependent upon tag reporting by fishers. Information from the first two years of recovery was suspect and was dropped from the tag abundance index.

#### 2.3.5 CWT tag loss rate

Double CWT tagging experiments were conducted between 1998 and 2006 to estimate the tag loss rates. The estimated tag loss rates were used to adjust the number of tag returns. In 1998, 2262 black rockfish were released with double CWT tags on both the left and right sides of the fish in order to estimate the instant CWT tag loss rate per year. In total, there were 2209 fish returned with double CWT tags, 58 fish returned with left CWT tag loss and 66 fish returned with right CWT tag loss (Table 16 and Figure 23). The estimate the instant rate of tag loss per year was – 0.0017 (st. err=0.0003, P=0.0035).

# 2.3.6 Population estimate

Petersen's method was used to estimate the population size from capture and recapture data. The method requires only two survey periods; the first survey involves the initial marking of  $n_1$  fish, of which m tagged fish are recovered from  $n_2$  fish sampled in the second survey. The estimated population size is

$$\hat{N} = \frac{(n_1 + )(n_2 + 1)}{m + 1} - 1$$

and

$$\operatorname{Var}(\hat{N}) = \frac{(n_1 + 1)(n_2 + 1)(n_1 - m)(n_2 - m)}{(m+1)^2(m+2)}.$$

The assumptions are

- i) The tags are not lost and always identified on recapture.
- ii) The population is closed.
- iii) Every individual has the same probability of recapture.

When the tag loss rate is known, the new estimate is  $\hat{m} = \frac{m}{1 + \hat{\beta}}$ .

The estimated population from 2000 to 2006 are given in Table 17.

# 3.0 Modeling History

In the 1994 stock synthesis model configuration, two auxiliary data sets were used as black rockfish abundance indicators: tagging CPUE, and coastal recreational bottomfish directed effort (Wallace and Tagart, 1994).

In 1999 we constructed an assessment model by using the AD model builder software (Fournier1997) to assess black rockfish abundance (Wallace et al 1999). The three key

features of the 1999 model were (1) the parameterization of the expected catches at age, (2) the definitions of the sampling unit for the different types of data input, and (3) the integration of tagging data explicitly. The parameterization chosen mostly affected parameter bias whereas the sampling unit designation mostly affected estimator variance. Both bias and variance were components of overall parameter uncertainty. The parameterization and the sampling unit definitions were both designed to conform to the actual sampling protocol used, thereby propagating sampling uncertainty through to the final biomass estimates.

# 4.0 Current Model Description

The current assessment employed Stock synthesis 2 (SS2V2.00c, compiled 3/27/2006) to model the dynamics of black rockfish population found between Cape Falcon, Oregon and North to the U.S./Canadian border in Coastal Waters. This model is a forward projecting, separable, age-structured model developed by Methot (2006). The convergence criterion for maximum gradient was set to 1.0e-5. The model was specified to begin in 1915 to ensure population equilibrium at the start of the modeling time period. Catch data were decayed from the last reliable catch estimates (1965) to 0 by 1940. Fisheries catch, size and age compositions were pooled into three fishery types including trawl, sport and non-trawl. The first size-age compositions were collected in the mid 1970's from the trawl fishery, but samples were not collected on a systematic basis until 1985. Growth (Lmin, Lmax and k) was estimated within the model to account for fishery selection of the larger individual fish at age. The population model was tuned to two fisheries-independent indices that include a tagging CPUE (1986-2007) and a tag abundance biomass index (2000-2007) both derived from WDFW black rockfish tagging information. Both STAT and STAR Panel members agreed that the available fishery dependent indices should not be incorporated due to potential bias resulting from bag limit changes and undocumented measures of fishing effort resulting from changes in search time across the time series. The black rockfish STAR base and STAT best fit models down-weights size composition for all fisheries (emphasis=0.1) to improve model fit to the age composition and indices rather than length. Given that length compositions are from all samples including aged samples, down weighting mitigates effects that may be contributed to the model by "double-counting" composition sampling.

There are 10 likelihood components for data including: 1) tag abundance, 2) tag CPUE, 3) trawl size compositions, 4) sport size compositions, 5) non-trawl size compositions, 6) tag survey size compositions, 7) trawl age compositions, 8) sport age compositions, 9) non-trawl age compositions and 10) mean size at age (Table 18).

There are a total of 76 parameters estimated within the base model and assumptions on priors are listed in Table 19. We modeled the black rockfish spawner-recruit relationship using the Beverton-Holt curve. The key steepness parameter (h), which determines overall productivity of a stock, is fixed at 0.6 and the prior on h is set to 0.3 in the STAR base model and in the STAT best fit model. Based on Dorn's (personal communication) Bayesian meta-analysis of productivity for west coast rockfish stocks, steepness parameter (h) for black rockfish should be in the 0.6-0.7 range and variation about the

stock-recruit curve (Sigma R) would be near 0.57. The natural mortality for females is assumed to be constant (0.16) for ages <=10 and then increasing to 0.2 by age 15, and males were assumed to have a constant natural mortality of 0.16 for all ages in the STAR base model. The natural mortality for females is assumed to be constant (0.16) for ages <=10 and then increasing to 0.24 by age 15, and males were assumed to have a constant natural mortality of 0.16 for all ages in the STAT best fit model.

# Sample size and effective sample size

Initial sample size inputs were based upon methods presented at the NMFS 2006 Stock Assessment Data and Modeling workshop that incorporates objective weighting for length- and age-frequency data for West coast groundfish fisheries where: Fishery data:

```
Effective \ N = ((0.138*FPS)+1)*NS \quad where: FPS < 44 \\ Effective \ N = 7.06*NS \quad where: FPS > 44 \\ Survey \ data: \\ Effective \ N = ((0.070*FPS)+1)*NS \quad where: FPS < 55 \\ Effective \ N = 4.89*NS \quad where: FPS > 55 \\ NS = Number \ of \ samples
```

Comparison of input sample size and the effective sample sizes estimated by the STAR base model are provided in Tables 20 and 21.

# 5.0 Model Selection and Evaluation

FPS = Average number of fish per sample

A large number of model structures were initially explored prior to establishing a base black rockfish model. Our primary goal in model selection was to ensure fit to the tag abundance index and age composition data while minimizing the overall likelihood. This is because we have confidence in the study design and methodology of our current tagging program and the resulting abundance estimates. In addition, we have collected numerous age samples from the fisheries during the last two decades that likely represents the underlying age structure of the population.

Natural mortality for mature females (>10 years of age) was assumed to be 0.20 (STAR base) and 0.24 (STAT best fit) and constant 0.16 for males and females < age 11. These rates are within the range of natural mortality rates estimated external to synthesis. Both male and female natural mortality rates are lower than that estimated in the 1999 assessment (Figure 25) and somewhat lower than the 1982 catch curve estimate of 0.265 (Wallace et al., 1994). The natural mortality in the current assessment is higher than that used in the 2003 assessment for black rockfish populations off Oregon and California (Ralston and E.J. Dick, 2003), which used a natural mortality of 0.1 and 0.2 for males and females, respectively.

Results of the model sensitivity analysis on natural mortality (Table 24) indicate that the STAT best fit model provided a better overall fit to the data compared to the STAR base

model and estimates of fishing mortality is closer to tagging study results (Figure 26). We conclude that the STAR base model should be used to base management decisions and set allowable harvest. A list of supporting information include: 1) the assumed rate of natural mortality in the "Low Natural Mortality" state of nature is lower than any previous assessment for the "Northern" population, is lower that any external estimation by direct and indirect methods, 2) biomass results from the "Low Natural Mortality" indicate that the population declined to less than 13% of the unfished population in the mid-1990's yet we have no indication from the fishery or from our tagging study that there was localized depletion during this time period, 3) sensitivity analyses indicate "Low Natural Mortality" model fit to the data is very poor relative to other model results that assume a higher rates of natural mortality, 4) the estimated q for the survey is likely double what it should be based on STAT knowledge of available habitat off the Washington coast, 5) tagging data are not fit well and tagging estimates external to the model indicate that the population is larger and fishing mortality is lower compared to STAR base model run results, 6) other model runs with higher steepness and Sigma R fit the data better and improved our view of the current population status above both the STAR base and "Best Fit" model runs and finally, 7) compared to the STAT best fit model a model with high natural mortality for females (where M=0.16 for males and females <=10 years of age and M for females linearly increasing from age 11 to age 15 to 0.26) fit the data equally well. This model resulted in an improved view of current population status above both the STAR base and STAT best-fit model runs. However, results from this model were not incorporated in the decision table because the higher natural mortality on females (0.26) fell outside the range considered at the STAR Panel.

Convergence properties using a parameter jitter of 0.001 was also explored in the base model and results indicate no other local minima (Figure 27). Growth was assumed linear to age 5 where variation in fork-length at age was stabilized across ages (Figure 28). Growth was fully (Lmin, Lmax and k) estimated within the model to account for fishery selection that favors the largest individuals at age. Model estimates of growth compared reasonably to external estimates and there were no apparent differences in estimates of growth between STAR and STAT models (Figure 29).

Both the STAR nor STAT models underestimated the increasing trend in tag abundance and tag CPUE indices in most recent years (Figure 30). We believe this is due to several factors including that tagging is not incorporated in the model as a tagging experiment, which is not possible within the current SS2 modeling framework. Further, the model estimated effective q for the tagging index was 0.83 and this is likely double what it should be based on STAT knowledge of available habitat off the Washington coast. The north central Washington coast, where most of the nearshore rocky habitat exists, is inaccessible to most recreational fishers and is not part of the current tagging program. However, the estimation of q is complicated by the fact that the SS2 value of q is a function of selectivity that is strongly dome shaped for the fishery. Increasing the weighting on survey abundance demonstrates that a better fit to the survey abundance index significantly improves our view of the current population status (Figure 31). Additionally, a retrospective analysis of the STAR base and STAT best-fit models shows that the indices strongly influence population trends and that the population trajectory in

most recent years was highly influenced by the large (estimated) 1999 year class (Figure 32).

Without an objective evaluation of an informed prior on q it is difficult to compare a prior conception of q based on tagging and the one estimated by SS2. Other issues include the non-independence of the length/age compositions and non-independence of the tagging abundance and CPUE series. However, both STAR and STAT conclude the current model configuration(s) represented the "best available" scientific information and should be used for management.

There appears to be some pattern in the size composition residuals such that model estimates for small fish were much higher than that observed in the trawl fishery fit. However, forcing the model to fit the size compositions degraded the fit to the age composition. Fit to size compositions in the sport, non-trawl and survey showed little trend. Overall, fit (or lack of) to the size composition data did not draw significant debate at the STAR panel and model fit to size compositions is likely within the uncertainty (Figures 33-36). Model fit to the age composition data showed relatively inconsistent patterns and was considered to be within model uncertainty (Figures 37-39). There was an obvious trade off where forcing the model to fit the age data degraded the fit to the size composition data. This was not fully resolved and is discussed below in the uncertainty section.

During the September, 2007 Council meeting the SSC recommended that management of the "Northern" black rockfish stock be based on the "STAT best fit" model.

#### 6.0 Base-run Results

Comparison of STAR base model recruitment estimates to the previous assessments and the STAT best fit model indicates similar estimated recruitment patterns (Figure 40). It is apparent that the large estimated recruitment in 1994 and 1999 is highly influential in determining current stock status. Due to lack of good recruitments and intense fishing by multiple fisheries, highest fishing mortality rates occurred in the late 1980's (Figure 41). Selectivity was domed-shaped (STAR and STAT models) in both the tagging survey and sport fishery and asymptotic in the trawl and non-trawl commercial fisheries (Figure 42). Comparison of STAR base model spawning biomass estimates to the previous assessments indicate a similar declining trend through the mid 1990's and then sharply increasing to 43% of the unfished biomass by 2006, though the trend is lower in the current model (Figure 43). The STAT best fit model resulted in a slightly smaller unfished biomass and a larger ending biomass compared to the STAR base model, biomass estimates show little difference in population trend (Figure 44).

Black rockfish stock abundance was below the Councils' management target a number of years and also dipped below the Councils' minimum stock size threshold in the STAR base model. The STAT best fit model population trajectory remained just above minimum stock size threshold. Both models indicate that the stock is currently well above the management target of B40% (Figure 45). The corresponding exploitation rate

relative to spawning biomass shows similar trend and harvest rates have exceeded management targets between the mid 1980's through the mid 1990's (Figure 46). Biomass, numbers-at-age, exploitation and selectivity resulting from the final base model is shown in Tables 25-27.

# 7.0 Uncertainty and Sensitivity

A retrospective analysis of the STAT best-fit models indicates that 1) indices strongly influence population trends and 2) that the population trajectory in most recent years was highly influenced by the large (estimated) 1999 year class (Figure 32).

Natural mortality is confounded with fishing mortality making it one of the most difficult biological parameters to estimate. In this assessment we explored direct methods to estimate natural mortality and compared to estimates derived from indirect methods (from other biological parameters, e.g., the growth constant and fecundity) in previous assessments. It is apparent from our analysis using both direct and indirect methods that our current assumptions on natural mortality in the STAR base model are within our limits to estimate this parameter and that the "low natural mortality rate" model used to bracket uncertainty is likely too low. Model sensitivity analysis show that other model configurations using higher natural mortality assumptions such as the STAT best fit model provides a better overall fit to the data (Figure 47).

Tagging is not incorporated in the model as a tagging experiment, which is not possible within the current SS2 modeling framework. The index for tagging abundance is not fit well and the model estimated effective q for the tagging index was 0.83. This is likely double what it should be based on STAT knowledge of available habitat off the Washington coast. Further, the north central Washington coast, where most of the nearshore rocky habitat exists, is inaccessible to most recreational fishers and is not part of the current tagging program. However, the estimation of q is complicated by the fact that the SS2 value of q is a function of selectivity that is strongly dome shaped for the fishery. Increasing the weighting on survey abundance shows that a better fit to the survey abundance index significantly improves our view of the current population status (Figure 48). Without an objective evaluation of an informed prior on q it is difficult to compare a prior conception of q based on tagging and the one estimated by SS2. Other issues include the non-independence of the length/age compositions and non-independence of the tagging abundance and CPUE series.

Likelihood profile of the STAR base assessment model for different fixed values of the Beverton-Holt steepness parameter (h) and Sigma R show that higher values (STAR base and STAT best-fit model had the steepness fixed at 0.6 and Sigma R tuned to 0.35) of both parameters improved the overall fit to the data (Figure 49). Our assumption on h is well within the uncertainties based on the Dorn meta-analysis, but assumptions on Sigma R may be too low (Dorn personal communication).

Changes in likelihood profile for various components of the base assessment model following changes in the emphasis (weight) of the recruitment Dev and Dev time series indicate very modest changes in fit for weighting between 0.1 to 100 with fit improving to age compositions and declining fit to size compositions with increasing emphasis (Figure 50).

Likelihood profile for various components of the base assessment model following changes in the emphasis (weight) on the abundance and tag CPUE indices indicate a slight improvement in fit by increasing emphasis to 10 on most components with the exception to the fit to sport size and age that declined (Figure 51). Increasing emphasis on the age composition for all fisheries above 1 improves fit to the abundance indices but increased likelihood for the fishery size components (Figure 52). The model was very sensitive to increasing emphasis on the size compositions and declined fit to all age and index components substantially (Figure 53).

# 8.0 Reference Points and Forecast

The Pacific Fisheries Management Council recommends that a default target fishing mortality rate of FSPR=0.5 be used for Council managed rockfish species. The current assessment uses this default for harvest projections and based on the Councils control rule for groundfish would not be considered overfished. Reference points and benchmark fishing mortality rates from this model are shown in Table 23. Forecast ABC's, Spawning biomass and depletion is shown for both the "STAR base" and STAR base model and the STAT best fit model in Table 24. During the September, 2007 Council meeting the SSC recommended that management of the "Northern" black rockfish stock be based on the "STAT best fit" model.

# 9.0 ACKNOWLEDGEMENTS

We would like to acknowledge the dedication of numerous volunteers who donated their time to the tagging survey and technicians who spent countless hours collecting fishery and biological information making this assessment possible. We would also like to thank Rick Methot (NWFSC) and Ian Taylor (NWFFSC) for their advice in model structuring and for providing parsing programs that aided immensely in analyzing results. Meetings with Industry and Managers also provided greatly to our understanding of the fishery and perceived stock condition. In great admiration, we would especially like to thank WDFW manager Brian Culver for his unyielding conservation ethic and his contributions to wise management of a valuable public resource for future generations.

#### 10.0 LITERATURE CITED

Berkeley, S.A., Chapman, C., and Sogard, S.M. 2004. Maternal age as a determinant of larval growth and survival in a marine fish, Sebastes Melanops. Ecology 85(5):1258-1264.

Bobko, S.J. and Berkeley, S.A. 2004. Maturity, ovarian, cycle, fecundity, and age-specific parturition of black rockfish (Sebastes melanops). Fish. Bull. 102:418-429.

Cheng and Gallinat, 2004) – Need cite.

Dorn 2002. Need cite.

Efron, B. 1982. *The Jacknife, the Bootstrap and Other Resampling Plans*. SIAM (Regional Conference Series in Applied Mathematics, 38). Philadelphia.

Fournier, D. 1997. An Introduction to AD Model Builder for use in non-linear Modeling and statistics. Otter Research Ltd.

Hall P. 1992. The Bootstrap and Edgeworth Expansion, Springer, New York. Lo, N.C.H, L.D. Jacobson and J.L. Squire 1992. Indices of relative abundance from fish spotter data based on delta-lognormal models. *Can. J. Fish. Aquat. Sci.* 49: 2515-2526.

Jackson, J. and Y.W. Cheng. 2001. Improving parameter estimation for daily egg production method of stock assessment of pink snapper in Shark Bay, Western Australia. *Journal of Agricultural, Biological and Environmental Statistics* 6: 243-257.

Lai H, R. Moore, and J.V. Tagart. 1991. Methodologies for estimating catch and effort statistics of ocean sport fishery off the Washington coast with users. Guide for the program "OSP.FOR'. Wash. Dept. Fish. Prog. Rept. No. 289: 35p.

Lo et al 1992.(need cite for delta lognormal model).

Miller, D.J. and R.N. Lea. 1972. Guide to the coastal marine fishes of California. Calif. Dept. Fish. Game Fish Bull. 157:1-249.

McCullagh and Nelder, 1989 – Need Cite.

Methot (2006)

Nei(1978)

Pauly, D., 1983. Length-converted catch curves. A powerful tool for fisheries research in the tropics. (Part I). *ICLARM Fishbyte*, 1(2):9-13

Peterson – need cite fo Peterson estimate.

Ralston and Dick 2003 - need cite

Sen, A.R. 1984. Sampling commercial rockfish landings in California. NOAA Tech. Memo. NMFS-SWFC-45, 95p.

Schnute growth model – need cite.

Syrjala, S.E. 2000. Critique on the use of the delta distribution for the analysis of trawl survey data. *ICES Journal of Marine Science* 57: 831-842.

Wallace, F.R. and J.V. Tagart, 1994. Status of the coastal black rockfish stocks in Washing and northern Oregon in 1994. *In* Pacific Fisheries Management Council. 1994. Appendix F: Status of the Pacific Coast Groundfish Fishery Through 1994 and Recommended Acceptable Biological Catches for 1995: Stock Assessment and Fishery Evaluation. Pacific Fisheries Management Council, Portland OR.

Wallace, F.R., Hoffmann, A., and Tagart, J.V. Status of the Black Rockfish Resource in 1999. In Pacific Fishery Management Council. 1999. Appendix: Status of the Pacific Coast Groundfish Fishery Through 1999 and Recommended Biological Catches for 2000: Stock Assessment and Fishery Evaluation. Pacific Fishery Management Council, 2130 SW Fifth Avenue, Suite 224, Portland, Oregon 97201.

(WDFW report in progress)-need genetics work cite

Wilkins, M.E., M. Zimmermann, and K.L. Weinberg. 1995. The 1995 Pacific coast bottom trawl survey of groundfish resources: Estimates of distribution, abundance, and length and age composition. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-89, 138 p. plus Appendices.

Table 1. Black rockfish catch North of Cape Falcon, Oregon by gear and year 1963-2006 (blanks indicate no data).

Coastal black rockfish catch North of Cape Falcon, Oregon

	Cod	ch by G	oar.	Catch by PMFC Area					
Year	Trawl		<i>ear</i> Ion-Trawl	3A	3B		TOTAL		
1963	19.0	Эроп	ion-irawi	19.0	0.0	0.0	19.0		
1963	8.0			8.0	0.0	0.0			
	108.0			108.0	0.0		8.0 108.0		
1965						0.0			
1966	186.0			186.0	0.0	0.0	186.0		
1967	234.0			234.0	0.0	0.0	234.0		
1968	122.0			122.0	0.0	0.0	122.0		
1969	261.0		20.5	261.0	0.0	0.0	261.0		
1970	303.0		20.5	320.4	3.1	0.0	323.5		
1971	134.1		17.5	147.6	4.0	0.0	151.6		
1972	116.0		29.3	137.7	7.5	0.0	145.3		
1973	48.0		26.8	63.7	11.1	0.0	74.8		
1974	75.0		51.2	106.8	19.4	0.0	126.2		
1975	156.0	62.3	36.9	216.7	38.5	0.0	255.2		
1976	347.2	36.8	32.7	384.5	32.3	0.0	416.7		
1977	15.0	76.0	52.2	96.1	47.2	0.0	143.2		
1978	96.0	94.2	89.8	185.2	94.8	0.0	280.1		
1979	321.3	150.7	104.0	500.5	75.5	0.0	576.0		
1980	64.6	144.8	70.5	228.9	51.0	0.0	279.9		
1981	213.0	213.8	81.8	436.6	72.0	0.0	508.6		
1982	185.1	135.7	128.9	364.8	84.9	0.0	449.7		
1983	327.5	244.3	134.1	458.2	247.7	0.0	705.9		
1984	218.9	302.2	145.8	513.2	153.8	0.0	666.9		
1985	127.3	305.3	272.0	407.8	296.8	0.0	704.6		
1986	158.6	391.1	103.0	534.0	118.8	0.0	652.8		
1987	82.0	389.3	220.1	494.3	197.0	0.0	691.3		
1988	129.0	414.2	129.3	521.1	151.5	0.0	672.5		
1989	124.4	369.6	165.3	469.3	188.0	2.0	659.3		
1990	43.3	387.2	119.4	386.9	163.0	0.0	549.9		
1991	46.2	332.3	83.4	320.3	139.6	1.9	461.9		
1992	71.4	342.9	132.3	327.2	219.3	0.0	546.5		
1993	46.8	316.9	88.4	298.3	152.9	1.0	452.2		
1994	1.0	358.6	106.3	323.9	141.6	0.4	465.9		
1995	3.3	264.8	65.8	214.9	118.7	0.3	333.9		
1996	0.0	264.2	8.6	204.4	68.4	0.0	272.8		
1997	9.0	234.1	15.0	194.8	63.2	0.1	258.1		
1998	73.1	259.4	4.8	268.1	69.0	0.3	337.4		
1999	0.0	221.6	4.3	169.4	56.5	0.0	225.9		
2000	0.0	224.8	1.2	158.2	67.9	0.0	226.1		
2001	0.0	188.7	1.1	134.3	55.5	0.0	189.8		
2002	0.2	238.9	1.5	173.5	67.1	0.0	240.6		
2003	0.1	237.1	0.2	166.8	70.6	0.0	237.4		
2004	0.6	268.0	0.7	174.4	94.9	0.0	269.3		
2005	0.0	331.7	0.9	217.8	114.7	0.0	332.5		
2006	1.4	321.5	1.2	248.7	75.4	0.0	324.1		
Mean	101.7	253.8	68.8	261.5	82.6	0.1	344.2		
Last 10 y	8.4	252.6		190.6	73.5	0.0	264.1		
Last 5 y	0.5	279.4		196.2	84.5	0.0	280.8		
•									

Table 2. Historical catch of rockfish by known rockfish catch categories between 1936 and 1969.

# Historical catch of rockfish by rockfish catch categories for coastal Washington Waters

wasiiiigioii waters		
Year	Black Rockfish	Rockfish
1936	-	0.1
1937	-	219.0
1938	-	273.7
1939	-	290.8
1940	-	330.2
1941	-	554.7
1942	-	1,925.0
1943	-	5,811.7
1944	-	9,084.7
1945	-	25,969.7
1946	-	11,322.2
1947	-	2,970.8
1948	-	5,192.1
1949	-	5,943.5
1950	-	151.1
1951	-	6,777.8
1952	151.5	-
1953	8.0	153.1
1954	16.1	2.8
1955	5.0	76.5
1956	7.8	-
1957	19.1	76.5
1958	71.8	33.1
1959	26.6	36.2
1960	96.2	32.7
1961	40.7	40.5
1962	12.5	22.5
1963	-	279.9
1964	3.4	38.7
1965	-	347.8
1966	1.0	36.6
1967	-	167.7
1968	-	130.9
1969	-	151.4

Note: Data from WDFW annual catch reports.

Table 3. Assumed catch and data input of black rockfish between 1940 and 1975. Bold italics represents catch assumptions and normal italics indicate the actual catch estimate based on fish ticket and species composition data.

	Ca	tch by Ge	ar
Year	Trawl	Sport	Non-Trawl
Initial	2.0	2.0	1.0
1940	3.2	2.8	0.0
1941	9.2	4.6	0.0
1942	15.2	6.3	0.0
1943	21.2	8.1	0.0
1944	27.2	9.8	0.0
1945	33.2	11.6	0.0
1946	39.2	13.3	0.0
1947	<i>52.0</i>	15.1	0.0
1948	51.2	16.8	0.0
1949	<i>57.2</i>	18.6	0.0
1950	63.2	20.3	1.5
1951	69.2	22.1	2.5
1952	<i>75.2</i>	23.8	3.5
1953	81.2	25.6	4.5
1954	87.2	27.3	5.5
1955	93.2	29.1	6.5
1956	99.2	30.8	7.5
1957	105.2	32.6	8.5
1958	111.2	34.3	9.5
1959	117.2	36.1	10.5
1960	123.2	37.8	11.5
1961	129.2	39.6	12.5
1962	135.2	41.3	13.5
1963	141.2	43.1	14.5
1964	108.0	44.8	15.5
1965	186.0	46.6	16.5
1966	234.0	48.3	17.5
1967	122.0	50.1	18.5
1968	261.0	51.8	19.5
1969	303.0	53.6	20.5
1970	134.1	55.3	17.5
1971	116.0	57.1	29.3
1972	48.0	58.8	26.8
1973	75.0	60.6	51.2
1974	156.0	62.3	36.9
1975	347.2	62.3	32.7

Presumed Catch SS2 input in bold italics.

Table 4. Estimated black rockfish discard in the Washington recreational sport fishery.

Black Rockfish Discard in the Washington Sport Fishery

Year	# of Fish	Mean Weight (kg)	Assumed Mortality	Catch Weight (mt)
2002	5,719	1.17	90%	6.0
2003	4,554	1.21	90%	5.0
2004	9,764	1.18	90%	10.4
2005	15,085	1.19	90%	16.2
2006	8,733	1.22	90%	9.6

Note: Discard not available prior to 2002

Table 5. Total effort (expanded) in Washington sport fisheries.

	All Trip Types		Bottom-Fish-Only Trips	
	ANGLERS	BOAT TRIPS	ANGLERS	BOAT TRIPS
1985	177,305	36,486	31,200	5,984
1986	213,459	47,941	36,223	6,536
1987	245,293	60,622	45,115	9,268
1988	254,412	67,793	47,793	9,299
1989	301,922	80,913	32,506	6,217
1990	198,095	50,245	36,572	7,109
1991	216,554	60,133	37,416	7,437
1992	174,219	48,476	40,248	8,960
1993	230,890	68,690	42,022	9,446
1994	55,288	12,039	40,005	8,009
1995	115,954	28,775	36,120	8,425
1996	144,324	39,575	32,950	6,822
1997	111,714	32,792	29,937	6,593
1998	81,429	22,740	29,818	6,012
1999	81,182	21,764	24,269	4,737
2000	113,869	31,976	22,563	4,169
2001	208,076	59,325	20,385	4,068
2002	153,200	40,120	20,394	3,817
2003	180,360	48,437	18,453	3,548
2004	184,615	51,119	22,188	4,733
2005	150,017	39,433	28,645	6,451
2006	122,067	31,743	30,138	6,321

Table 6. Summary of size composition data collected from commercial and recreational fisheries during 1976-2006.

Year	Number	of field sa	mples	Number of le	ngth meas	urements	Mea	n size (cn	1)
	Sport	Trawl	Non-Trawl	Sport	Trawl	Non-Trawl	Sport	Trawl	Non-Trawl
1976		4			782			47.5	
1977									
1978									
1979	7			508			46.4		
1980	8	2	2	703	206	96	45.4	46.1	45.9
1981	23	4		1468	400		43.3	46.8	
1982	9	4	1	263	400	29	40.7	44.5	40.1
1983	1	8	2	10	800	124	36.9	45.3	41.2
1984	21	3	1	835	300	100	40.4	44.7	40.9
1985	2			160			43.1		
1986	21	13	27	512	322	527	41.8	45.5	44.4
1987	23	16	25	645	401	722	43.3	47.3	43.2
1988	18	4	17	451	100	424	41.9	47.3	43.8
1989	16	9	12	397	225	299	42.2	49.1	44.7
1990	11	10	4	290	249	125	41.6	47.1	36.7
1991	22	12	19	720	302	500	40.8	47.1	40.2
1992	34	8	11	890	200	275	41.3	46.3	40.0
1993	35	5	13	866	125	325	40.6	46.9	40.4
1994	35	2	9	868	49	250	40.9	46.2	38.1
1995	32	2	9	814	50	225	40.5	45.7	39.6
1996	33			834			39.5		
1997	36	2		900	31		40.5	46.6	
1998	37	2		1327	85		39.8	43.6	
1999	34			1673			39.5		
2000	33	1		1650	3		40.0	47.3	
2001	36	1		1777	1		40.2	53.0	
2002	56	1		2629	50		40.9	47.8	
2003	58	1		2323	3		41.4	45.7	
2004	44	2		2002	15		41.0	51.7	
2005	61		1	2228		1	41.2		43.0
2006	152		2	2854		20	41.1		48.1

Table 7. Summary of age composition data collected from commercial and recreational fisheries during 1976-2006.

Year	Number	of field sar	nples	Nun	nber of ag	е	M	ean age	
	Sport	Trawl	Non-Trawl	Sport	Trawl	Non-Trawl	Sport		Non-Trawl
1976		2			238			11.3	
1977									
1978									
1979									
1980	4	2		364	195		14.3	13.0	
1981	2	4		71	394		10.6	11.8	
1982		3			295			10.2	
1983		8	1		794	100		10.2	11.8
1984	20	3	1	828	298	99	9.7	12.6	11.3
1985	2			160			10.8		
1986	21	13	27	506	321	525	9.3	10.1	11.9
1987	23	16	25	642	401	720	11.5	11.3	10.9
1988	18	4	17	448	99	416	10.0	12.1	10.8
1989	16	9	12	395	224	297	9.3	10.5	10.8
1990	11	10	4	289	249	125	9.4	11.2	7.3
1991	22	12	19	717	301	500	9.2	12.2	8.7
1992	34	8	11	889	200	275	9.7	10.1	9.0
1993	35	5	13	863	125	324	9.0	10.9	8.5
1994	35	2	9	866	48	250	9.6	13.4	7.7
1995	32	2	9	813	49	225	8.6	12.0	7.7
1996	33			829			8.5		
1997	36			893			9.6		
1998	37			1323			9.4		
1999	34			1655			9.1		
2000	33			1644			9.6		
2001	36			1773			9.7		
2002	38			1894			9.8		
2003	37			1841			9.6		
2004	33			1645			9.4		
2005	33			1603			9.6		
2006	30		1_	1484		19	9.5		14.3

Table 8. Summary of male and female black rockfish age to weight data with estimated errors in each conversion.

Age	N	Male	Fe	Female		
$\hat{t}$ (st. err.)	$\hat{L}$ (st.err.)	$\hat{W}$ (st.err.)	$\hat{L}$ (st.err.)	$\hat{W}$ (st.err.)		
1(0.002)	20.123(0.583)	0.161(0.019)	20.123(0.583)	0.163(0.019)		
2(0.004)	24.754(0.406)	0.286(0.029)	24.689(0.396)	0.288(0.028)		
3(0.006)	28.568(0.281)	0.426(0.040)	28.564(0.262)	0.431(0.039)		
4(0.007)	31.709(0.195)	0.569(0.052)	31.851(0.170)	0.583(0.051)		
5(0.009)	34.296(0.138)	0.708(0.064)	34.641(0.113)	0.735(0.063)		
6(0.011)	36.427(0.105)	0.838(0.076)	37.008(0.083)	0.883(0.076)		
7(0.013)	38.181(0.092)	0.955(0.086)	39.017(0.070)	1.022(0.088)		
8(0.015)	39.626(0.094)	1.059(0.096)	40.722(0.064)	1.150(0.099)		
9(0.017)	40.816(0.106)	1.149(0.104)	42.168(0.062)	1.267(0.109)		
10(0.019)	41.796(0.122)	1.228(0.111)	43.396(0.063)	1.371(0.118)		
11(0.020)	42.603(0.140)	1.295(0.117)	44.437(0.068)	1.465(0.126)		
12(0.022)	43.268(0.159)	1.352(0.123)	45.321(0.077)	1.547(0.133)		
13(0.024)	43.815(0.177)	1.400(0.127)	46.071(0.090)	1.618(0.139)		
14(0.026)	44.266(0.195)	1.441(0.131)	46.707(0.106)	1.681(0.144)		
15(0.028)	44.637(0.211)	1.474(0.134)	47.247(0.123)	1.735(0.149)		
16(0.030)	44.943(0.227)	1.503(0.137)	47.706(0.141)	1.782(0.153)		
17(0.032)	45.195(0.241)	1.526(0.139)	48.095(0.158)	1.823(0.157)		
18(0.033)	45.402(0.253)	1.546(0.141)	48.425(0.175)	1.858(0.160)		
19(0.035)	45.573(0.265)	1.562(0.143)	48.705(0.192)	1.888(0.163)		
20(0.037)	45.714(0.275)	1.576(0.144)	48.942(0.207)	1.913(0.165)		
21(0.039)	45.829(0.284)	1.587(0.146)	49.144(0.221)	1.935(0.167)		
22(0.041)	45.925(0.292)	1.596(0.147)	49.315(0.234)	1.954(0.169)		
23(0.043)	46.003(0.299)	1.603(0.147)	49.460(0.246)	1.970(0.171)		
24(0.045)	46.068(0.306)	1.610(0.148)	49.583(0.257)	1.983(0.172)		
25(0.046)	46.121(0.311)	1.615(0.149)	49.688(0.267)	1.995(0.173)		
26(0.048)	46.165(0.316)	1.619(0.149)	49.776(0.275)	2.005(0.174)		
27(0.050)	46.201(0.320)	1.623(0.150)	49.852(0.283)	2.013(0.175)		
28(0.052)	46.231(0.323)	1.626(0.150)	49.916(0.291)	2.020(0.176)		
29(0.054)	46.256(0.326)	1.628(0.150)	49.970(0.297)	2.026(0.177)		
30(0.056)	46.276(0.329)	1.630(0.150)	50.016(0.303)	2.032(0.177)		
31(0.058)	46.293(0.331)	1.632(0.151)	50.055(0.308)	2.036(0.178)		
32(0.059)	46.306(0.333)	1.633(0.151)	50.088(0.312)	2.040(0.178)		
33(0.061)	46.318(0.335)	1.634(0.151)	50.116(0.316)	2.043(0.179)		
34(0.063)	46.327(0.336)	1.635(0.151)	50.140(0.320)	2.046(0.179)		
35(0.065)	46.334(0.338)	1.636(0.151)	50.160(0.323)	2.048(0.179)		
36(0.067)	46.341(0.339)	1.636(0.151)	50.177(0.325)	2.050(0.179)		
37(0.069)	46.346(0.339)	1.637(0.151)	50.192(0.328)	2.051(0.180)		
38(0.071)	46.350(0.340)	1.637(0.151)	50.204(0.330)	2.053(0.180)		
39(0.072)	46.354(0.341)	1.638(0.151)	50.215(0.332)	2.054(0.180)		
40(0.074)	46.357(0.341)	1.638(0.151)	50.224(0.333)	2.055(0.180)		
41(0.076)	46.359(0.342)	1.638(0.151)	50.231(0.335)	2.056(0.180)		
42(0.078)	46.361(0.342)	1.638(0.152)	50.238(0.336)	2.057(0.180)		
43(0.080)	46.363(0.343)	1.639(0.152)	50.243(0.337)	2.057(0.180)		
44(0.082)	46.364(0.343)	1.639(0.152)	50.248(0.338)	2.058(0.180)		
45(0.084)	46.365(0.343)	1.639(0.152)	50.252(0.339)	2.058(0.180)		
46(0.085)	46.366(0.343)	1.639(0.152)	50.255(0.340)	2.059(0.180)		
47(0.087)	46.367(0.343)	1.639(0.152)	50.258(0.340)	2.059(0.181)		

48(0.089)	46.367(0.344)	1.639(0.152)	50.260(0.341)	2.059(0.181)
49(0.091)	46.368(0.344)	1.639(0.152)	50.262(0.341)	2.059(0.181)
50(0.093)	46.368(0.344)	1.639(0.152)	50.264(0.342)	2.060(0.181)
51(0.095)	46.369(0.344)	1.639(0.152)	50.265(0.342)	2.060(0.181)
52(0.097)	46.369(0.344)	1.639(0.152)	50.267(0.342)	2.060(0.181)
53(0.098)	46.369(0.344)	1.639(0.152)	50.268(0.343)	2.060(0.181)
54(0.100)	46.369(0.344)	1.639(0.152)	50.269(0.343)	2.060(0.181)
55(0.102)	46.369(0.344)	1.639(0.152)	50.269(0.343)	2.060(0.181)
56(0.104)	46.370(0.344)	1.639(0.152)	50.270(0.343)	2.060(0.181)
57(0.106)	46.370(0.344)	1.639(0.152)	50.270(0.343)	2.060(0.181)
58(0.108)	46.370(0.344)	1.639(0.152)	50.271(0.343)	2.060(0.181)
59(0.110)	46.370(0.344)	1.639(0.152)	50.271(0.344)	2.060(0.181)
60(0.112)	46.370(0.344)	1.639(0.152)	50.272(0.344)	2.061(0.181)

Table 9. Summary of the number of black rockfish fish sampled with age in maturity study and the expected probability of maturity with age.

Age	No. of immature fish	No. of mature fish	Expected probability of maturity
4	1	0	0.01
5	12	0	0.02
6	50	1	0.05
7	73	7	0.09
8	65	13	0.17
9	38	22	0.29
10	22	12	0.45
11	6	15	0.62
12	2	5	0.76
13	2	2	0.87
14	0	2	0.93
15	0	0	0.96
16	0	0	0.98
17	0	0	0.99
18	0	0	1.00
19	0	0	1.00
20	0	2	1.00

Table 10. Summary of the number of black rockfish fish sampled with fork length in maturity study and the expected probability of maturity with fork length.

Fork length (cm)	No. of immature	No. of mature fish	Expected
	fish		probability of
			maturity
25	1	0	0
26	1	0	0
27	1	0	0
28	2	0	0
29	3	0	0
30	7	0	0.01
31	3	1	0.01
32	5	0	0.02
33	13	0	0.02
34	18	0	0.03
35	30	0	0.05
36	32	3	0.07
37	37	4	0.11
38	30	8	0.15
39	35	10	0.22
40	27	12	0.29
41	20	13	0.38
42	14	9	0.48
43	8	10	0.59
44	4	11	0.68
45	2	2	0.76
46	0	7	0.83
47	1	3	0.88
48	0	2	0.92
49	0	2	0.94

Table 11. Summary of the estimated total mortality coefficients of male and female black rockfish from 1984 to 2006.

Year	Ma	ale	Fen	nale	
	N	$\hat{Z}$ (st. err.)	n	$\hat{Z}$ (st. err.)	
1984	267	0.162(0.068)	429	0.267(0.005)	
1988	128	0.169(0.098)	148	0.341(0.207)	
1989	180	0.256(0.112)	217	0.205(0.071)	
1990	132	0.200(0.044)	158	0.407(0.129)	
1991	326	0.213(0.050)	394	0.259(0.031)	
1992	424	0.187(0.080)	457	0.325(0.011)	
1993	364	0.270(0.048)	495	0.277(0.028)	
1994	399	0.244(0.013)	465	0.348(0.016)	
1995	372	0.304(0.009)	440	0.370(0.039)	
1996	399	0.394(0.080)	432	0.387(0.014)	
1997	437	0.298(0.079)	438	0.361(0.031)	
1998	947	0.315(0.043)	874	0.400(0.013)	
1999	851	0.320(0.034)	822	0.353(0.013)	
2000	741	0.316(0.071)	909	0.406(0.056)	
2001	800	0.353(0.026)	974	0.427(0.053)	
2002	783	0.324(0.064)	1066	0.298(0.057)	
2003	793	0.290(0.055)	1009	0.327(0.069)	
2004	731	0.254(0.066)	922	0.297(0.032)	
2005	681	0.238(0.092)	982	0.339(0.069)	
2006	806	0.220(0.074)	802	0.323(0.035)	

Table 12. Summary of the proportion by area and the number of recreational observations taken from 1990 to 2006.

Year			Area		
	1	2	3	74	84
1990	5102(2.87%)	159462(89.83%)	2202(1.24%)	5601(3.16%)	5144(2.90%)
1991	2156(1.43%)	138150(91.69%)	2602(1.73%)	3122(2.07%)	4643(3.08%)
1992	3422(2.82%)	97598(80.29%)	4159(3.42%)	10128(8.33%)	6252(5.14%)
1993	5636(5.13%)	88923(81.01%)	3153(2.87%)	6115(5.57%)	5942(5.41%)
1994	7754(4.37%)	148419(83.69%)	7552(4.26%)	7275(4.10%)	6340(3.58%)
1995	3442(2.42%)	112959(79.57%)	5118(3.61%)	10172(7.17%)	10271(7.24%)
1996	5018(3.02%)	133094(80.22%)	4179(2.52%)	8263(4.98%)	15349(9.25%)
1997	5771(4.67%)	100816(81.61%)	1729(1.40%)	5814(4.71%)	9400(7.61%)
1998	8048(5.79%)	110960(79.78%)	2711(1.95%)	4645(3.34%)	12720(9.15%)
1999	1951(1.77%)	93642(84.92%)	2801(2.54%)	4412(4.00%)	7470(6.77%)
2000	3524(3.09%)	93927(82.31%)	3125(2.74%)	6625(5.81%)	6918(6.06%)
2001	3814(4.01%)	77415(81.37%)	2232(2.35%)	5322(5.59%)	6355(6.68%)
2002	4610(4.54%)	79168(77.89%)	2823(2.78%)	8967(8.82%)	6079(5.98%)
2003	6589(7.25%)	68067(74.87%)	2735(3.01%)	6757(7.43%)	6766(7.44%)
2004	4599(4.66%)	74905(75.93%)	3706(3.76%)	6047(6.13%)	9399(9.53%)
2005	4136(3.43%)	84719(70.28%)	7052(5.85%)	9351(7.76%)	15280(12.68%)
2006	5769(4.31%)	106803(79.75%)	4558(3.40%)	6307(4.71%)	10492(7.83%)

Table 13. Summary of the recreational fishery CPUEs estimated from mean estimator and delta lognormal model for all areas.

Year	Total cate	ch/total an	glers		Delta lognormal model			
	Estimates	$q_{ar{X},2.5\%}$	$q_{ar{X},97.5\%}$	$CV_{\overline{X}}$	Estimates	$q_{ar{X},2.5\%}$	$q_{ar{X},97.5\%}$	$CV_{\overline{X}}$
1990	8.58	8.33	8.85	0.02	5.73	5.52	5.92	0.02
1991	7.37	7.18	7.60	0.02	5.43	5.24	5.61	0.02
1992	6.14	5.92	6.37	0.02	4.77	4.63	4.92	0.02
1993	5.83	5.61	6.11	0.02	4.24	4.13	4.42	0.02
1994	6.87	6.70	7.04	0.01	4.43	4.29	4.56	0.01
1995	5.94	5.75	6.10	0.01	4.07	3.94	4.18	0.02
1996	6.37	6.22	6.53	0.01	4.57	4.45	4.69	0.01
1997	5.78	5.64	5.94	0.02	3.93	3.81	4.05	0.02
1998	6.35	6.17	6.50	0.01	4.80	4.66	4.91	0.01
1999	6.93	6.73	7.07	0.01	4.86	4.70	4.99	0.02
2000	6.83	6.63	6.98	0.01	5.03	4.87	5.18	0.02
2001	6.46	6.25	6.66	0.01	4.29	4.13	4.44	0.02
2002	7.03	6.86	7.20	0.01	5.01	4.86	5.17	0.02
2003	6.93	6.75	7.12	0.01	4.95	4.75	5.14	0.02
2004	7.14	6.94	7.33	0.01	5.57	5.41	5.73	0.02
2005	6.98	6.80	7.13	0.01	5.36	5.21	5.48	0.01
2006	7.29	7.15	7.42	0.01	5.20	5.06	5.33	0.01

Table 14. Summary of the recreational sport CPUEs estimated from mean estimator and delta lognormal model for Area 2.

Year	Total catch/total anglers					Delta lognormal model			
	Estimates	$q_{\overline{X},2.5\%}$	$q_{ar{X},97.5\%}$	$CV_{\overline{X}}$	Estimates	$q_{ar{X},2.5\%}$	$q_{ar{X},97.5\%}$	$CV_{\overline{X}}$	
1990	10.98	10.66	11.29	0.02	10.84	10.51	11.18	0.01	
1991	8.75	8.54	8.96	0.01	8.35	8.11	8.56	0.01	
1992	7.35	7.01	7.63	0.02	6.85	6.53	7.11	0.02	
1993	7.52	7.24	7.85	0.02	7.16	6.92	7.48	0.02	
1994	9.64	9.43	9.86	0.01	9.33	9.13	9.55	0.01	
1995	8.31	8.16	8.46	0.01	7.81	7.64	8.01	0.01	
1996	8.03	7.83	8.23	0.01	7.63	7.45	7.81	0.01	
1997	7.23	7.01	7.44	0.01	6.37	6.12	6.59	0.02	
1998	7.44	7.20	7.63	0.01	6.76	6.57	6.95	0.01	
1999	8.54	8.35	8.72	0.01	7.70	7.46	7.91	0.02	
2000	8.36	8.18	8.58	0.01	7.80	7.60	7.99	0.01	
2001	8.25	8.03	8.47	0.01	7.08	6.81	7.35	0.02	
2002	8.85	8.63	9.05	0.01	7.95	7.68	8.24	0.02	
2003	8.46	8.24	8.68	0.01	6.83	6.51	7.11	0.02	
2004	8.10	7.86	8.31	0.01	6.86	6.58	7.12	0.02	
2005	8.77	8.60	8.93	0.01	7.80	7.60	8.03	0.02	
2006	8.92	8.78	9.05	0.01	8.16	7.96	8.33	0.01	

Table 15. Central Washington coastal tagging mean catch per trip (catch/hours fished).

## Central Washington Coast Tagging CPUE Mean Catch Per Hour

Year	<b>Across All Trips</b>	In(1+cv)
1981	4.8	0.666
1986	2.3	0.5993
1987	1.2	0.6344
1988	8.0	0.5539
1989	1.2	0.9771
1990	1.0	0.8439
1998	2.5	0.813
1999	3.1	0.7407
2000	2.2	0.5684
2001	4.7	0.6076
2002	5.5	0.5034
2003	6.2	0.5913
2004	9.4	0.5149
2005	10.2	0.7579
2006	10.5	0.4205

Table 16. Summary of the return of tagged fish from the CWT double tags experiment.

Year			i	No. of one tag return $(r_{si})$		No. of two tags return $(r_{di})$
				left	right	
199	8	0		8	17	691
199	9	1		14	11	542
200	00	2		14	18	433
200	)1	3		14	8	276
200	)2	4		6	8	160
200	)3	5		2	2	73
200	)4	6		0	2	34

Table 17. Summary of the year, the no. of fish tagged, no. of fish sampled, the numbers of fish return with tags, tag on the right, tag on the left, double tag, the estimated population size and variance, the adjusted no. of tag return with tag loss, the estimated population size with tag loss adjustment and variance.

Year	$n_1$	$n_2$	m	$m_r$	$m_l$	$m_d$	Ñ	$\operatorname{Var}(\hat{N})$	m	Ñ	$\operatorname{Var}(\hat{\ddot{N}})$
1998	2456	46951	14	1	1	12	7.69E+06	3.67E+12	14.08	7.65E+06	4.53E+12
1999	3479	66253	43	1	0	42	5.24E+06	6.02E+11	43.01	5.24E+06	6.46E+11
2000	2789	65276	130	3	5	122	1.39E+06	1.39E+10	130.13	1.39E+06	1.53E+10
2001	3210	64440	68	2	1	65	3.00E+06	1.26E+11	68.03	3.00E+06	1.35E+11
2002	4089	68475	143	1	1	141	1.94E+06	2.51E+10	143.01	1.94E+06	2.66E+10
2003	6747	77622	246	1	8	237	2.12E+06	1.74E+10	246.09	2.12E+06	1.86E+10
2004	4209	53385	74	1	1	72	3.00E+06	1.16E+11	74.01	3.00E+06	1.23E+11
2005	3913	70482	54	0	0	54	5.02E+06	4.43E+11	54.00	5.02E+06	4.66E+11

Table 18. Likelihood components from the STAR base (top) and STAT best-fit (bottom) northern black rockfish models.

#### STAR Base Model

Likelihood Components	<b>Emphasis</b>	Likelihood
indices		_
Tag Abundance	1.0	43.4
Tag CPUE	1.0	11.7
discard	0.0	0.0
length_comps		
Trawl	0.1	67.6
Sport	0.1	32.3
Non-Trawl	0.1	38.1
Tag	0.1	18.3
age_comps		
Trawl	1.0	187.2
Sport	1.0	395.3
Non-Trawl	1.0	187.0
size-at-age	0.0	105.9
mean_body_wt	0.0	0.0
Equil_catch	1.0	0.0
Recruitment	0.1	14.5
Parm_priors	1.0	0.0
Parm_devs	0.1	0.0
penalties	0.0	0.0
Forecast_Recruitment	0.0	0.2
	·	1101 6

1101.6

STAT Best Fit Model

<b>Likelihood Components</b>	<b>Emphasis</b>	Likelihood
indices		
Tag Abundance	1.0	41.5
Tag CPUE	1.0	10.4
discard	0.0	0.0
length_comps		
Trawl	0.1	69.2
Sport	0.1	32.5
Non-Trawl	0.1	39.4
Tag	0.1	19.0
age_comps		
Trawl	1.0	180.6
Sport	1.0	386.8
Non-Trawl	1.0	185.7
size-at-age	0.0	106.5
mean_body_wt	0.0	0.0
Equil_catch	1.0	0.0
Recruitment	0.1	15.4
Parm_priors	1.0	
Parm_devs	0.1	0.0
penalties	0.0	0.0
Forecast_Recruitment	0.0	0.2
-	•	1007 15

1087.15

Table 19. Assumptions and Priors used in the Northern black rockfish STAR base model. The only change in the STAT Best Fit model is an increase in the "old" female natural mortality rate from 0.20 to 0.24.

# Assumptions and Priors used in the 2007 Northern Black Rockfish Model Negative Phase indicates that it was not estimated.

Gender:_1Pat	tern:_1	Phase	#_size_sel:_1_Trawl
1	0.16	-2	1 47.2587 2
2	0.24	-2	2 3.9375 3
3	27.4969	2	3 4 -3
4	47.6046	2	4 2.2 -3
5	0.257678	3	5 -8 2
6	0.08	-2	6 8.99978 2
7	0.08	-3	#_male
Gender:_2Pat			#_size_sel:_2-Sport
8	0.16	-2	7 40.1805 2
9	0.16	-2	8 -6 3
10	28.4314	2	9 3.5 -3
11	45.3143	2	10 3 -3
12	0.239616	3	11 -8 2
13	0.07	-3	12 0.55712 2
14	0.07	-3	#_male
biology_parms			#_size_sel:_3_Non-Trawl
15	4.03E-05	-3	13 39.8172 2
16	2.768	-3	14 3.38361 3
17	42.6	-3	15 3.60126 3
18	-0.4	-3	16 4.57262 3
19	-0.3657	-3	17 -8 2
20	0.7674	-3	18 8.99922 2
21	3.80E-05	-3	#_male
22	2.782	-3	#_size_sel:_4
SR_parms	0.40400		19 1 -2
1	8.12489	1	20 19 -3
2	0.6	-5	#_male
3	0.3	-4	#_size_sel:_5
4	0	-3	21 1 -2 22 19 -3
5	0	-1	
6	0	-99	#_male
			#_size_sel:_6
			23 40.4008 2
			24 -0.682246 3
			25 3.70382 3
			26 1.10507 3 27 -7.10564 2
			28 -0.475443 2

Table 20. Average Pearson residual by fishery (Trawl=1, Sport=2, Non-Trawl=3) by likelihood component.

Used	1
year	(All)

				Data						For Age & Len:
kind	fleet	season	mkt	Average of A	verage of effN	ا Average	Min of Pe I	Max of P∈S	tdDev of Pearson	effN/inputN
AGE	1	1	0	66	103	-0.026	-2.39	4.86	0.823880087	1.57
	2	1	0	312	343	0.027	-2.67	3.84	0.885559936	1.10
	3	1	0	116	157	0.004	-2.44	8.05	0.988462445	1.35
LEN	1	1	0	110	44	0.785	-3.78	14.38	2.878351254	0.40
	2	1	0	135	340	0.161	-3.08	7.96	1.137126017	2.52
	3	1	0	199	139	0.430	-3.72	9.64	2.065073952	0.70
	6	1	0	217	204	0.001	-2.71	4.79	1.166505276	0.94
L@A	2	1	0	42	1.2326	-0.454	-4.37	3.95	1.501360074	0.03

Table 21. Average Pearson residual by fishery (Trawl=Top 2 rows, Sport=Middle 2 rows, Non-Trawl Bottom 2 rows) , age and sex.

Mean Pearson residule by Age and Fleet (Trawl=1, Sport =2 and Non-Trawl=3)																											
Average of Pearson			bin																								
fleet	gender		3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	30	Grand Total
1	1	1	0.33	-0.35	-0.22	-0.38	-0.06	-0.02	0.29	-0.10	0.11	0.20	0.08	-0.30	-0.35	-0.40	-0.24	-0.45	-0.56	-0.44	-0.49	-0.62	-0.31	-0.29	-0.63	-0.16	-0.20
		2	1.95	-0.19	0.00	0.21	0.64	1.01	0.82	0.43	-0.20	0.24	0.24	-0.13	-0.04	-0.08	-0.20	-0.26	-0.27	-0.19	0.23	0.00	-0.13	-0.03	-0.04	0.03	0.12
2	2	1	-0.22	0.00	0.17	0.05	0.16	0.59	0.46	0.56	0.53	0.46	0.32	0.10	0.26	-0.01	-0.08	-0.05	-0.18	-0.27	-0.25	-0.43	-0.29	-0.42	-0.33	0.60	0.10
		2	-0.63	-0.25	0.04	-0.02	-0.09	-0.09	-0.68	-0.90	-0.64	-0.81	-0.43	-0.28	0.02	-0.05	0.26	0.23	0.25	0.58	0.45	0.25	0.36	0.40	0.47	0.47	-0.04
3	3	1	-0.15	0.73	0.80	0.24	0.25	0.09	0.22	-0.36	-0.09	0.21	0.48	0.03	0.30	0.28	-0.10	0.00	-0.29	-0.32	-0.56	0.09	-0.57	-0.40	-0.47	1.25	0.10
		2	0.00	-0.24	0.58	-0.05	-0.31	-0.29	-0.58	-0.25	-0.68	-0.24	-0.05	-0.10	-0.10	-0.30	0.05	-0.16	-0.02	0.19	0.11	0.02	0.15	-0.01	0.05	0.82	-0.07
Grand Total			-0.20	-0.02	0.24	0.00	0.09	0.25	0.08	-0.10	-0.13	-0.02	0.07	-0.12	0.02	-0.09	-0.03	-0.08	-0.12	0.01	0.02	-0.08	-0.02	0.00	0.06	0.47	0.01
Grand Total	3	1 2 1 2	-0.22 -0.63 -0.15 0.00	0.00 -0.25 0.73 -0.24	0.17 0.04 0.80 0.58	0.05 -0.02 0.24 -0.05	0.16 -0.09 0.25 -0.31	0.59 -0.09 0.09 -0.29	0.46 -0.68 0.22 -0.58	0.56 -0.90 -0.36 -0.25	0.53 -0.64 -0.09 -0.68	0.46 -0.81 0.21 -0.24	0.32 -0.43 0.48 -0.05	0.10 -0.28 0.03 -0.10	0.26 0.02 0.30 -0.10	-0.01 -0.05 0.28 -0.30	-0.08 0.26 -0.10 0.05	-0.05 0.23 0.00 -0.16	-0.18 0.25 -0.29 -0.02	-0.27 0.58 -0.32 0.19	-0.25 0.45 -0.56 0.11	-0.43 0.25 0.09 0.02	-0.29 0.36 -0.57 0.15	-0.42 0.40 -0.40 -0.01	-0.33 0.47 -0.47 0.05	0.60 0.47 1.25 0.82	

Table 22. Model sensitivity by likelihood component to rates of natural mortality relative to that assumed in the STAR base model. Values represent the change in likelihood relative to the base models such that negative values indicate a better fit. Female natural mortality rates for ages less than 11 are assumed to be equal to that assumed for males.

Total Like	otal Likelihood Female Natural Mortality Rate											
		0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3
	0.1	270.0	186.0	128.4	90.6	67.2	54.2	48.8	49.2	54.1	62.4	73.6
te a	0.12	226.1	140.6	84.5	48.5	26.2	13.6	16.4	8.1	12.4	20.0	30.5
Ra	0.14	205.1	124.0	70.6	35.7	13.0	-0.9	-8.1	-10.0	-7.5	-1.6	7.1
ž ž	0.16	203.4	127.5	76.5	41.1	16.5	0.0	-10.0	-14.7	-15.0	-11.6	-5.1
Male Natural Mortality Rate	0.18	216.3	144.4	95.1	58.6	31.3	11.4	-2.2	-10.3	-12	-13.5	-9.9
ŽΣ	0.2	240.0	171.6	122.8	89	54.6	30.9	13.4	1.3	-6.0	-9.2	-8.7
	0.22	272.4	206.4	112.8	102.7	90.0	57.4	33	19.2	7.8	0.7	-2.4
Fit to Tag	Abund	dance										
		0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3
	0.1	127.1	95.1	71.4	54.9	43.7	35.9	30.5	26.6	23.7	21.5	19.8
te a	0.12	97.1	62.1	40.2	26.9	18.9	13.8	7.4	8.3	6.8	5.7	4.9
atu 7 Ra	0.14	72.1	39.9	21.8	12.2	7.0	3.9	2.1	0.9	0.1	-0.3	-0.7
Male Natural Mortality Rate	0.16	54.1	26.3	11.9	5.2	1.8	0.0	-1.0	-1.5	-1.8	-1.9	-2.0
Mai	0.18	41.7	18.1	6.8	2.0	0.0	-0.9	-1.3	-1.4	-1	-1.2	-1.1
_ ≥	0.2	33.1	13.5	4.3	2.1	-0.1	-0.3	-0.1	0.2	0.5	0.8	1.1
	0.22	27.5	10.9	2.5	2.8	3.2	1.5	2	2.7	3.3	3.8	4.2
Fit to Tag	CPUE											
		0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3
	0.1	26.4	22.6	19.3	16.7	14.6	13.0	11.8	10.8	10.0	9.4	9.0
ral ate	0.12	22.9	17.9	13.8	10.8	8.6	7.0	5.8	4.9	4.3	3.8	3.4
Male Natural Mortality Rate	0.14	19.4	13.7	9.3	6.2	4.2	2.7	1.7	1.0	0.5	0.1	-0.1
a e	0.16	16.3	10.5	6.0	3.0	1.2	0.0	-0.8	-1.3	-1.7	-2.0	-2.2
Mal	0.18	13.6	7.9	3.6	1.0	-0.7	-1.6	-2.3	-2.7	-3	-3.2	-3.3
2	0.2	11.5	5.9	2.2	0	-1.8	-2.5	-3.0	-3.3	-3.5	-3.6	-3.7
	0.22	9.8	4.6	-3.7	-3.6	-3.5	-2.9	-3	-3.4	-3.5	-3.6	-3.6
Fit to all I	ndices											
		0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3
	0.1	153.4	117.7	90.7	71.6	58.3	48.9	42.2	37.4	33.7	30.9	28.8
te a	0.12	120.0	80.0	54.0	37.7	27.5	20.8	13.1	13.3	11.1	9.5	8.3
Male Natural Mortality Rate	0.14	91.6	53.7	31.0	18.5	11.1	6.6	3.8	1.9	0.7	-0.2	-0.8
a E	0.16	70.4	36.8	17.9	8.2	3.0	0.0	-1.8	-2.8	-3.5	-3.9	-4.2
Mal	0.18	55.4	26.0	10.5	3.0	-0.7	-2.6	-3.6	-4.1	-4	-4.4	-4.4
_ 2	0.2	44.6	19.5	6.5	2.3	-1.8	-2.8	-3.1	-3.1	-3.0	-2.8	-2.6
	0.22	37.3	15.5	-1.2	-0.8	-0.3	-1.5	-1	-0.7	-0.2	0.2	0.6

Note: Square indcates the Base Model and bold font indicates best fit.

Table 22. Continued.

Fit to Length Composition					Female Natural Mortality Rate										
			0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3		
		0.1	2.0	-1.6	-3.1	-3.2	-2.4	-1.1	0.5	2.4	4.3	6.3	8.3		
	e a	0.12	5.6	8.0	-1.5	-2.3	-2.0	-0.9	17.1	2.3	4.2	6.2	8.1		
	atur Ra	0.14	8.0	4.6	1.0	-0.7	-1.1_	-0.5	0.7	2.2	4.0	5.9	7.9		
	Male Natural Mortality Rate	0.16	8.2	9.4	4.1	1.2	0.0	0.0	0.8	2.0	3.6	5.4	7.3		
	Male orta	0.18	9.1	10.7	7.5	3.3	1.1	0.4	0.7	1.7	-8.4	4.7	6.6		
	ŽΣ	0.2	10.7	11.5	11.6	5.5	2.2	0.7	0.5	1.1	2.2	3.7	5.5		
		0.22	12.7	13.0	10.3	8.3	4.8	0.8	-18.4	0.2	1.1	2.4	4.1		
Fit to	o Age C	omposi	ition												
			0.1	0.12	0.14	0.16	0.18	0.2	0.22	0.24	0.26	0.28	0.3		
		0.1	89.8	58.9	35.4	19.4	9.6	5.0	4.7	8.0	14.3	23.1	34.1		
	g a	0.12	91.0	57.5	32.4	14.4	2.2	-5.0	-18.6	-7.0	-2.9	3.9	13.1		
	Male Natural Mortality Rate	0.14	102.8	66.8	40.8	20.2	4.7	-6.0	-12.2	-14.4	-13.1	-8.9	-2.0		
	<u> </u>	0.16	126.3	83.4	57.1	33.7	14.5	0.0	-10.0	-15.8	-17.8	-16.5	-12.3		
	Malo	0.18	155.7	111.2	79.6	53.8	30.9	12.4	-1.7	-11.3	46.3	-19.1	-17.9		
	_ ≥	0.2	190.0	145.1	106.8	57.4	53.7	31.0	12.7	-1.2	-10.9	-16.7	-19.0		
		0.22	228.5	183.0	59.1	55.7	54.0	55.8	114.8	14.6	0.7	-9.2	-15.3		
Dep	letion L	.evel													
			0.10	0.12	0.14	0.16	0.18	0.20	0.22	0.24	0.26	0.28	0.3		
		0.1	0.05	0.06	0.08	0.11	0.13	0.15	0.17	0.19	0.21	0.23	0.300		
	ate a	0.12	0.05	0.08	0.12	0.15	0.19	0.23	0.25	0.29	0.32	0.34	0.24		
	Male Natural Mortality Rate	0.14	0.07	0.11	0.16	0.22	0.27	0.32	0.37	0.41	0.45	0.48	0.37		
	ality N	0.16	0.08	0.13	0.21	0.29	0.36	0.43	0.49	0.54	0.58	0.62	0.51		
	Mal	0.18	0.10	0.17	0.26	0.35	0.45	0.53	0.60	0.67	0.72	0.76	0.65		
	_ ≥	0.2	0.12	0.20	0.29	0.46	0.53	0.63	0.72	0.79	0.85	0.90	0.80		
	1	<b>0.22</b> Note: Sau	0.14 lare indca	0.23	0.84 Jase Mod	0.86 el and bol	0.87 d font indi	0.73	1.27 st fit.	0.90	0.97	1.02	0.94		
	·	1010. 040													
B0			0.10	0.12	0.14	0.16	0.18	0.20	0.22	0.24	0.26	0.28	0.3		
		0.1	2620	2458	2301	2159	2034	1923	1825	1738	1661	1592	1531		
		0.12	2700	2548	2404	2281	2171	2071	1986	1901	1827	1761	1701		
	ural Rate	0.12	2741	2579	2486	2396	2315	2238	2164	2096	2031	1971	1915		
	Male Natural Mortality Rate	0.14	2780	2577	2540	2513	2474	2433	2389	2344	2298	2253	2207		
	ale rtali	0.18	2809	2615	2583	2620	2656	2674	2683	2685	2676	2670	2652		
	₽ĕ	0.16	2836	2652	2574	2725	2875	2992	3102	3205	3297	3378	3443		
		0.22	2856	2684	13213	10728	8357	3460	3792	4124	4496	4887	5283		

Note: Square indcates the Base Model and bold font indicates best fit.

Table 23. Comparison of Councils' default target fishing mortality rates and reference points between the STAR base model and STAT best fit model. The default target fishing mortality rate of FSPR=0.5 is used in this assessment for both models and that used for other Council managed rockfish species.

#### STAR Base Model Results

Unfished Stock	Value
Age 3+ Biomass (B <sub>0</sub> ) (mt)	10,813
Spawning Biomass SB(0) (mt)	2,429
SPBio/Recruit (kg/fish)	0.780
Age1 Recruitment (R <sub>0</sub> ) (1,000's)	3,113
Steepness_R0_S0	0.6

#### Reference points based on

Exploited Stock	Estimated MSY	SB <sub>40%</sub>	SPR (SB <sub>0.5</sub> )
SPR (Spawning Biomass/Recruit)	0.413	0.400	0.400
F (Fishing Mortality Rate)	0.132	0.101	0.101
Exploitation Rate (Yield/Bsmry)	0.076	0.060	0.060
MSY (mt) or MSY proxy (mt)	377	361	361
Yield (mt)	718	972	972
SPBIO/SB(0)	29.6%	40.0%	40.0%
Age 3+ Biomass	4,947	6,012	6,012

#### STAT Best Fit Model Results

Unfished Stock	Value
Age 3+ Biomass (B <sub>0</sub> ) (mt)	11,390
Spawning Biomass SB( <sub>0</sub> ) (mt)	2,321
SPBio/Recruit (kg/fish)	0.687
Age1 Recruitment (R <sub>0</sub> ) (1,000's)	3,377
Steepness_R0_S0	0.6

#### Reference points based on

Exploited Stock	Estimated MSY	SB <sub>40%</sub>	SPR (SB <sub>0.5</sub> )
SPR (Spawning Biomass/Recruit)	0.418	0.400	0.40
F (Fishing Mortality Rate)	0.141	0.110	0.110
Exploitation Rate (Yield/Bsmry)	0.081	0.065	0.065
MSY (mt) or MSY proxy (mt)	423	408	408
Yield (mt)	700	928	928
SPBIO/SB(0)	30.1%	40.0%	40.0%
Age 3+ Biomass	5,218	6,264	6,264

Table 24. Comparison of ABC's, Spawning biomass and depletion between the STAR base (top) and STAT best fit model (bottom).

#### STAR Base Model

Year	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ABC (mt)	394	377	361	350	345	344	346	350	354	357
Spawning Biomass (mt)	1064	1071	1060	1036	1005	977	956	944	940	943
% of Virgin	0.438	0.441	0.436	0.426	0.414	0.402	0.394	0.389	0.387	0.388

#### STAT "Best Fit" Model and SSC recommendation

Year	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
ABC (mt)	535	521	505	478	459	448	443	441	442	443
Spawning Biomass	1281	1317	1328	1270	1204	1140	1087	1048	1023	1010
% of Virgin	0.552	0.567	0.572	0.547	0.519	0.491	0.468	0.452	0.441	0.435

Table 25. Biomass time series for final base model (STAT best fit).

						Exploitation (Popes approx)					
year	bio-all	bio-smry	SpBio	SpBio/B0	recruit-0	Trawl	Sport	Non-Trawl			
VIRG	13,226	11,390	2,321		3,377			_			
INIT	13,226	11,390	2,321		3,377						
1915-1939	,	11,390	2,321	1.00	3,377	0.000	0.000	0.000			
1940		11,389	2,320	1.00	3,377	0.001	0.000	0.000			
1941	13,220	11,384	2,319	1.00	3,377	0.002	0.001	0.000			
1942		11,372	2,315	1.00	3,376	0.003	0.001	0.000			
1943	•	11,355	2,309	0.99	3,375	0.004	0.001	0.000			
1944	•	11,334	2,301	0.99	3,373	0.005	0.001	0.000			
1945	•	11,308	2,292	0.99	3,370	0.006	0.002				
1946	•	11,278	2,281	0.98	3,368	0.007	0.002				
1947		11,246	2,270	0.98	3,365	0.009	0.002				
1948		11,204	2,255	0.97	3,361	0.009	0.002				
1949	12,994	11,167	2,242	0.97	3,358	0.010	0.003				
1950		11,128	2,228	0.96	3,354	0.011	0.003				
1951	12,908	11,084	2,214	0.95	3,350	0.013	0.003				
1952		11,039	2,198	0.95	3,346	0.014	0.003				
1953		10,990	2,182	0.94	3,342	0.015	0.004				
1954		10,939	2,165	0.93	3,338	0.016	0.004				
1955		10,886	2,148	0.93	3,333	0.017	0.004				
1956		10,832	2,130	0.92	3,328	0.019	0.004				
1957	•	10,775	2,112	0.91	3,323	0.020	0.005				
1958	•	10,717	2,093	0.90	3,317	0.021	0.005				
1959	•	10,657	2,074	0.89	3,312	0.023	0.005				
1960		10,596	2,055	0.89	3,306	0.024	0.006				
1961	12,331	10,534	2,035	0.88	3,300	0.025	0.006				
1962		10,471	2,015	0.87	3,294	0.027	0.006				
1963		10,406	1,994	0.86	3,288	0.028	0.006				
1964		10,340	1,974	0.85	3,281	0.022	0.007				
1965		10,308	1,965	0.85	3,279	0.038	0.007				
1966		10,207	1,933	0.83	3,268	0.048	0.007				
1967		10,073	1,889	0.81	3,253	0.026	0.008				
1968		10,050	1,882	0.81	3,869	0.055	0.008				
1969	•	9,904	1,834	0.79	3,252	0.065	0.009				
1970	•	9,735	1,777	0.77	2,832	0.030	0.009				
1971	11,461	9,847	1,777	0.77	2,829	0.026	0.009	0.004			

Table 25. (Continued)

						Exploitation (Popes approx)						
year	bio-all	bio-smry	SpBio	SpBio/B0	recruit-0	Trawl	Sport	Non-Trawl				
1,972	11,402	9,852	1,781	0.77	2,891	0.011	0.010	0.004				
1,973	11,527	9,839	1,806	0.78	3,569	0.016	0.010	0.007				
1,974	11,741	9,796	1,816	0.78	4,242	0.034	0.010					
1,975	11,603	9,689	1,806	0.78	2,795	0.075	0.010	0.005				
1,976	11,267	9,517	1,744	0.75	2,646	0.003	0.006					
1,977	11,269	9,772	1,787	0.77	2,817	0.021	0.012					
1,978	11,210	9,605	1,777	0.77	3,374	0.072	0.016					
1,979	10,806	9,217	1,680	0.72	2,596	0.015	0.026					
1,980	10,603	9,148	1,655	0.71	2,093	0.051	0.025					
1,981	10,211	9,013	1,593	0.69	1,937	0.045	0.037					
1,982	9,718	8,640	1,528	0.66	1,920	0.083	0.025					
1,983	9,313	8,134	1,443	0.62	2,625	0.058						
1,984	8,876	7,639	1,360	0.59	2,286	0.036	0.061	0.048				
1,985	8,398	7,060	1,257	0.54	2,470	0.048						
1,986	7,902	6,739	1,186	0.51	1,689	0.026	0.090					
1,987	7,543	6,235	1,085	0.47	3,027	0.046						
1,988	7,037	5,836	981	0.42	1,931	0.049						
1,989	6,950	5,314	853	0.37	4,011	0.020						
1,990	6,753	5,240	774	0.33	2,433	0.023	0.129					
1,991	6,614	5,023	710	0.31	2,361	0.037						
1,992	6,437	5,158	653	0.28	2,267	0.026	0.124					
1,993	6,445	5,128	619	0.27	2,623	0.001	0.116					
1,994	6,860	5,122	608	0.26	4,625	0.002	0.130					
1,995	6,910	5,180	606	0.26	2,349	0.000	0.092					
1,996	7,070	5,423	649	0.28	2,157	0.005	0.084					
1,997	7,361	5,977	707	0.30	3,104	0.036	0.071	0.001				
1,998	7,509	6,066	762	0.33	2,705	0.000	0.076					
1,999	8,186	6,147	826	0.36	5,355	0.000	0.063					
2,000	8,405	6,516	891	0.38	2,429	0.000	0.059	0.000				
2,001	8,515	6,739	959	0.41	2,075	0.000	0.046					
2,002	8,635	7,405	1,043	0.45	2,283	0.000	0.056	0.000				
2,003	8,737	7,485	1,114	0.48	2,536	0.000	0.054					
2,004	8,835	7,470	1,171	0.50	2,702	0.000	0.060	0.000				
2,005	9,018	7,564	1,211	0.52	2,780	0.000	0.070					
2,006	9,063	7,558	1,239	0.53	2,819	0.000	0.066	0.000				

Table 26. Numbers-at-age (thousands of fish) by year for the final base model (STAT best fit).

Females Ages 0-20

<b>Female</b>	s Ag	es Q-	<b>20</b> ,	2	4	_		7	٥	0	10	4.4	40	10	4.4	45	10	47	10	10	20
year VIRG	1689	1439	1226	3 1045	4 890	5 759	6 647	7 551	8 470	9 400	10 341	11 291	12 244	13 201	14 163	15 131	16 103	17 81	18 64	19 50	20 39
INIT	1689	1439	1226	1045	890	759	647	551	470	400	341	291	244	201	163	131	103	81	64	50	39
1915-1939	1689	1439	1226	1045	890	759	647	551	470	400	341	291	244	201	163	131	103	81	64	50	39
1940 1941	1689 1689	1439 1439	1226 1226	1045 1045	890 890	759 759	647 647	551 551	469 469	400 400	341 341	291 290	244 243	201 201	163 163	131 130	103 103	81 81	64 63	50 50	39 39
1942	1688	1439	1226	1045	890	759	647	551	469	399	340	290	243	201	163	130	103	81	63	50	39
1943	1687	1438	1226	1045	890	759	647	551	469	399	340	289	242	200	162	130	102	80	63	50	39
1944	1686	1438	1226	1045	890	759	647	551	469	399	339	288	242	199	162	129	102	80	63	49	39
1945 1946	1685 1684	1437 1436	1225 1225	1045 1044	890	759 750	646 646	550 550	468 468	398 398	338 338	288 287	241 240	198 197	161 160	129 128	101 101	80 79	63 62	49 49	39 38
1947	1682	1435	1223	1044	890 890	759 758	646	550	468	397	337	286	239	196	159	127	100	78	62	49	38
1948	1681	1434	1223	1043	889	758	646	550	467	396	336	285	238	195	158	126	99	78	61	48	38
1949	1679	1432	1222	1042	889	758	646	550	467	396	335	284	237	194	157	125	98	77	61	48	37
1950 1951	1677 1675	1431 1429	1220 1219	1041 1040	888 887	757 757	645 645	549 549	467 466	395 395	335 334	283 282	236 235	193 192	156 155	124 123	97 96	76 76	60 59	47 47	37 37
1952	1673	1428	1218	1039	886	756	644	548	465	394	333	281	234	191	154	122	96	75	59	46	36
1953	1671	1426	1216	1038	885	755	644	548	465	393	332	280	233	190	153	121	95	74	58	45	36
1954	1669	1424	1215	1037	884	754	643	547	464	392	331	279	231	189	152	120	94	73	57	45	35
1955 1956	1666 1664	1422 1420	1213 1212	1035 1034	883 882	753 753	642 642	546 546	463 462	391 390	330 329	278 276	230 229	188 186	151 150	119 118	93 92	72 72	57 56	44 44	35 34
1957	1661	1418	1212	1034	881	752	641	545	461	389	327	275	227	185	148	117	91	71	55	43	34
1958	1659	1416	1208	1031	880	751	640	544	460	388	326	274	226	184	147	116	90	70	54	42	33
1959	1656	1413	1206	1030	879	750	639	543	459	387	325	272	225	182	146	115	89	69	54	42	33
1960 1961	1653 1650	1411 1409	1204 1202	1028 1026	877 876	749 747	638 637	542 541	458 457	386 385	324 322	271 270	223 222	181 179	144 143	114 112	88 87	68 67	53 52	41 41	32 31
1962	1647	1406	1202	1025	875	746	636	540	456	383	321	268	220	178	142	111	86	66	51	40	31
1963	1644	1404	1198	1023	873	745	635	539	455	382	319	267	219	177	140	110	85	66	51	39	30
1964	1641	1401	1196	1021	872	744	634	538	454	381	318	265	217	175	139	109	84	65	50	39	30
1965 1966	1639 1634	1398 1397	1194 1191	1019 1017	870 868	743 741	633 632	538 536	454 451	381 378	318 315	265 262	217 214	175 172	138 136	108 106	83 82	64 63	49 48	38 37	30 29
1967	1627	1392	1190	1017	867	740	631	534	449	374	310	257	210	168	133	103	79	61	47	36	28
1968	1935	1386	1187	1014	865	738	630	534	450	375	311	258	210	168	132	103	79	60	46	36	27
1969	1626	1649	1181	1011	864	737	628	532	446	371	307	253	205	164	129	100	76	58	45	34	26
1970 1971	1416 1415	1386 1207	1405 1181	1007 1197	861 858	736 734	627 627	530 531	443 445	367 370	302 304	247 249	199 201	159 159	124 124	96 96	73 73	56 56	43 42	33 32	25 25
1972	1445	1205	1028	1006	1020	731	624	530	445	371	307	251	202	160	125	96	73	55	42	32	25
1973	1784	1232	1027	876	857	869	622	529	446	373	311	256	207	164	127	98	74	56	43	33	25
1974	2121	1520	1049	875	746	730	739	525	444	372	310	258	209	166	129	99	75	56	43	33	25
1975 1976	1398 1323	1808 1191	1296 1540	894 1104	746 762	636 635	621 541	624 523	440 518	368 359	307 296	255 245	208 199	165 159	129 124	99 95	74 72	56 54	43 41	32 31	25 23
1977	1408	1127	1015	1312	941	649	541	458	441	435	301	249	202	162	127	98	74	55	42	31	24
1978	1687	1200	960	865	1118	801	552	455	381	364	358	247	200	160	126	97	74	56	42	31	24
1979	1298	1438	1023	818	737	952	680	461	373	307	289	281	190	151	119	92	70 69	53	40	30	22
1980 1981	1047 969	1106 892	1225 942	871 1044	697 742	627 594	808 532	570 676	382 468	306 308	251 244	236 199	225 183	150 172	117 112	90 87	66	52 50	40 38	30 29	22 22
1982	960	825	760	803	889	632	503	442	547	372	242	191	152	138	128	82	62	47	36	27	21
1983	1312	818	703	647	684	757	535	418	357	432	289	186	143	112	100	90	57	43	33	25	19
1984 1985	1143 1235	1118 974	697 953	599 594	551 510	582 469	640 490	441 519	333 343	278 253	332 208	220 247	139 161	105 100	81 74	71 56	63 48	40 43	30 27	23 21	17 16
1986	844	1052	830	812	506	434	396	402	411	265	193	158	184	118	72	53	39	34	30	19	14
1987	1513	720	897	707	691	430	365	319	308		195	141	114	130	82	49	36	27	23		13
1988	965	1290	613	764	602	588		294		228		141	101	80	90	56	33		18		14
1989 1990	2005 1216	823 1709	1099 701	522 936	650 445	512 553	493 430	287 394	220 218		161 127	156 117	97 112	68 68	53 47	59 36	36 40	21 24	15 14	12 10	10 8
1991	1180	1036	1456	597	797	378	464	344	298	160	117	92	83	78	47	32	24	27	16	10	7
1992	1134	1006	883	1240	508	677	317		258	216	114	83	64	57	53	32	21		18	11	6
1993	1312	966	857	752	1056	432	569	254	280	189	156	82	59	45	39	36	21	14	11	12	7
1994 1995	2312 1174	1118 1970	823 952	730 701	641 622	898 545	363 755	456 291	193 347	207 143	139 152	115 102	60 83	42 43	32 30	27 22	25 19	15 17	10 10	7 7	8 5
1996	1078	1001	1679	811	597	529	460	620	232		112	119	79	63	32	22	16	14	12	7	5
1997	1552	919	853	1431	691	508	447	379	495	182	213	87	92	60	47	24	16	12	10	9	5
1998	1353	1322	783	727	1219	588	430	370	304	390	142	165	67	69	44	35	17	11	8	7	6
1999 2000	2677 1214	1153 2282	1127 982	667 960	619 568	1037 527	498 879	356 414	298 290	242 240	309 195	113 249	129 89	51 101	52 40	33 40	25 25	12 19	8 9	6 6	5 5
2001	1037	1035	1944	837	818	484	447	732	338	235	194	157	198	70	78	30	30	19	14	7	5
2002	1141	884	882	1657	713	697	411	374	604	277	192	158	126	157	55	60	23	23	14	11	5
2003	1268	973	753	751	1411	607	591	342	306	490	224	155	126	99	122	42	45	17	17	11	8
2004 2005	1351 1390	1080 1151	829 921	642 706	640 547	1202 545	515 1019	493 429	281 402	249 227	397 201	182 320	124 144	100 97	77 77	93 59	31 70	34 24	13 25	13 10	8 10
2006	1409	1185	981	784	602	466		844		322			252		74	58	44		18		7

Table 26. (Continued)

#### Females Ages 21-40

Table 26. (Continued) *Males Ages* 0-20

Year	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
VIRG	1689	1439	1226	1045	890	759	647	551	470	400	341	291	248	211	180	153	131	111	95	81	69
INIT	1689	1439	1226	1045	890	759	647	551	470	400	341	291	248	211	180	153	131	111	95	81	69
1915-1939	1689	1439	1226	1045	890	759	647	551	470	400	341	291	248	211	180	153	131	111	95	81	69
1940 1941	1689 1689	1439 1439	1226 1226	1045 1045	890 890	759 759	647 647	551 551	469 469	400 400	341 341	291 290	248 247	211 211	180 180	153 153	131 130	111 111	95 95	81 81	69 69
1942	1688	1439	1226	1045	890	759	647	551	469	400	340	290	247	210	179	153	130	111	95	81	69
1943	1687	1438	1226	1045	890	759	647	551	469	399	340	289	246	210	179	152	130	111	94	80	68
1944	1686	1438	1226	1045	890	759	646	551	469	399	339	289	246	209	178	152	129	110	94	80	68
1945	1685	1437	1225	1045	890	759	646	551	468	398	339	288	245	209	178	151	129	110	93	80	68
1946	1684	1436	1225	1044	890	759	646	550	468	398	338	287	244	208	177	150	128	109	93	79	67
1947 1948	1682 1681	1435 1434	1224 1223	1043 1043	890	758 750	646	550 550	468 468	398 397	338 337	287 286	243 242	207 206	176	150 148	127 126	108	92	79 78	67 66
1948	1679	1434	1223	1043	889 889	758 758	646 646	550	468	397	336	285	242	205	175 174	148	125	107 107	91 91	77	66
1950	1677	1431	1220	1042	888	757	645	550	467	396	336	284	241	204	173	147	124	106	90	76	65
1951	1675	1429	1219	1040	887	757	645	549	467	396	335	284	240	203	172	146	124	105	89	76	64
1952	1673	1428	1218	1039	886	756	644	549	466	395	335	283	239	202	171	145	123	104	88	75	64
1953	1671	1426	1216	1038	885	755	644	548	465	394	334	282	238	201	170	144	122	103	87	74	63
1954	1669	1424	1215	1037	884	754	643	547	465	394	333	281	237	200	169	143	121	102	86	73	62
1955 1956	1666 1664	1422 1420	1213 1212	1035 1034	883 882	753 753	642 642	547 546	464 463	393 392	332 331	280 279	236 235	199 198	168 167	142 141	120 119	101 100	85 84	72 71	61 60
1956	1661	1420	1212	1034	881	753 752	641	545	462	392	330	279	235	198	166	139	119	99	84	71	60
1958	1659	1416	1208	1031	880	751	640	544	461	390	329	277	233	195	164	138	116	98	83	70	59
1959	1656	1413	1206	1030	879	750	639	544	461	389	328	275	231	194	163	137	115	97	82	69	58
1960	1653	1411	1204	1028	877	749	638	543	460	388	326	274	230	193	162	136	114	96	81	68	57
1961	1650	1409	1202	1026	876	747	637	542	459	387	325	273	229	192	161	135	113	95	80	67	56
1962	1647	1406	1200	1025	875	746	636	541	458	386	324	272	227	190	159	133	112	94	78	66	55
1963	1644	1404	1198	1023	873	745 744	635	540	457 456	385	323 322	270 269	226 225	189	158	132	110	92	77 76	65	55 54
1964 1965	1641 1639	1401 1398	1196 1194	1021 1019	872 870	744	634 633	539 538	455	384 383	321	269	224	187 187	156 156	131 130	109 109	91 91	76 76	64 63	53
1966	1634	1397	1191	1013	868	741	632	537	453	381	319	266	222	185	154	128	103	89	74	62	52
1967	1627	1392	1190	1015	867	740	631	535	451	378	315	262	218	181	151	125	104	87	72	60	50
1968	1935	1386	1187	1014	865	738	629	535	451	378	316	263	218	181	150	125	103	86	72	60	50
1969	1626	1649	1181	1011	864	737	628	533	449	375	312	259	214	177	147	121	101	83	69	58	48
1970	1416	1386	1405	1007	861	736	627	531	446	372	308	254	209	173	142	117	97	80	67	55	46
1971	1415 1445	1207	1181	1197	858	734	626	531	447	374	310	256	210	173	142 143	117	97	80	66	55	45 45
1972 1973	1784	1205 1232	1028 1027	1006 876	1020 857	731 869	624 622	530 529	447 447	374 376	311 314	257 261	212 215	174 177	145	117 119	97 98	80 81	66 67	55 55	45 45
1974	2121	1520	1049	875	746	730	739	526	445	374	313	261	216	178	147	120	99	81	67	55	45
1975	1398	1808	1296	894	746	636	621	625	442	371	310	258	215	178	146	120	98	81	66	55	45
1976	1323	1191	1540	1104	762	635	541	524	522	364	302	251	207	171	141	116	95	78	64	52	43
1977	1408	1127	1015	1312	941	649	540	458	443	439	306	254	210	174	144	118	97	80	65	53	44
1978	1687	1200	960	865	1118	801	551	456	383	367	362	252	208	172	142	118	97	79	65	53	44
1979	1298	1438	1023	818	737	952	680	463	376	312	295	288	199	164	135	111	92	75 75	62	51	41
1980 1981	1047 969	1106 892	1225 942	871 1044	697 742	627 593	808 532	571 677	384 472	310 313	255 250	241 204	235 192	162 187	133 128	110 105	90 87	75 71	61 59	50 48	41 40
1982	960	825	760	803	889	632	502	443	552	378	248	197	160	150	145	100	82	67	55	46	38
1983	1312	818	703	647	684	757	535	419	362	442	298	193	152	123	114	111	76	62	51	42	35
1984	1143	1118	697	599	551	582	639	443	338	285	344	229	148	116	93	87	84	57	47	39	32
1985	1235	974	953	594	510	469	490	522	349	259	216	257	171	110	86	69	64	62	43	35	29
1986	844	1052	830	812	506	434	396	403	418	273	200	165	195	129	83	65	52	49	47	32	26
1987	1513	720	897 613	707 764	691	429 597	364 361	320 296	313	315	202 232	146 147	120 106	143 87	94 103	61 68	47 44	38 34	35 28	34 26	23 25
1988 1989	965 2005	1290 823	613 1099	764 522	602 650	587 511	492	289	249 225	236 182	168	164	103	74	61	72	48	30	24	19	18
1990	1216	1709	701	936	445	552	429	397	223	168	133	122	118	75	54	44	52	34	22	17	14
1991	1180	1036	1456	597	797	378	463	345	305	165	122	96	88	85	54	39	32	38	25	16	12
1992	1134	1006	883	1240	508	676	316	372	264	225	119	87	68	62	60	38	27	22	27	18	11
1993	1312	966	857	752	1056	432	567	255	286	196	163	86	63	49	45	43	27	20	16	19	13
1994	2312	1118	823	730	640	897	362	458	197	214	145	120	63	46	36	33	32	20	15	12	14
1995 1996	1174 1078	1970 1001	952 1679	701 811	621 597	544 529	753 459	292 621	354 235	147 279	158 115	106 123	88 82	46 68	34	26 26	24	23 19	15 18	11 12	9 8
1996	1552	919	853	1431	691	508	447	380	500	186	219	90	96	68 64	36 53	26 28	21 21	16	15	14	9
1998	1353	1322	783	727	1218	588	430	371	308	398	146	171	70	74	50	41	22	16	13	11	11
1999	2677	1153	1127	667	619	1037	497	357	301	246	316	116	135	55	59	40	33	17	13	10	9
2000	1214	2282	982	960	568	527	878	415	292	244	198	254	93	108	45	47	32	26	14	10	8
2001	1037	1035	1944	837	818	484	446	734	341	237	197	160	204	75	88	36	38	26	21	11	8
2002	1141	884	882	1657	713	696	410	374	607	280	194	161	130	167	61	72	29	31	21	18	9
2003	1268	973	753	751 642	1411	607	590 514	343	308	495	227	157	130	105	135	49	58 40	24 47	25	17	14
2004 2005	1351 1390	1080 1151	829 921	642 706	640 547	1201 545	514 1018	494 430	283 405	251 229	401 203	183 323	127 148	105 102	85 85	109 69	40 88	47 32	19 38	21 16	14 17
2006	1409	1185	981	784	601	465	461			326		162		118	82	68	55		26		


Table 26. (Continued)

#### Males Ages 21-40

Table 27. Final base model length selectivity (cm) by fishery.

Length(bin)	Trawl	Sport	Non-Trawl
21	0.000	0.000	0.000
23	0.000	0.000	0.001
25	0.000	0.001	0.003
27	0.001	0.006	0.012
29	0.003	0.023	0.041
31	0.008	0.079	0.120
33	0.024	0.211	0.282
35	0.064	0.445	0.531
37	0.146	0.737	0.805
39	0.287	0.959	0.982
41	0.488	1.000	1.000
43	0.717	0.989	1.000
45	0.911	0.883	1.000
47	0.999	0.752	1.000
49	1.000	0.673	1.000
51	1.000	0.644	1.000
53	1.000	0.637	1.000
55	1.000	0.636	1.000
57	1.000	0.636	1.000

### **Figures**






Figure 1. Location of black rockfish tag release area (top) and tag recovery locations (bottom).




Figure 2. Dendogram showing results of cluster analysis of ten black rockfish collections using Nei's (1978) unbiased genetic distance at 20 polymorphic loci.




Figure 3. Relative abundance of females with age in pooled age data for Washington fisheries.




Figure 4. Total black rockfish catch by gear and year for areas North of Cape Falcon to the U.S. Canadian border.




Figure 5. Regulation changes in commercial fisheries




Figure 6. Maximum retainable rockfish catch per trip for the sport fisheries.




Figure 7. Maximum retainable black for the limited entry commercial fisheries.




Figure 8. Comparison of catch estimates between the 1999 and the current assessment.




Figure 9. Proportion at size by sex and fisheries from 1984 to 2006.





Figure 10. Proportion at age by sex and fisheries from 1984 to 2006.






Figure 11. Scatter plot of fork length and weight of male (top panel) and female (bottom panel) black rockfish and the expected length weight relationship.




Figure 12. Scatter plots of male (top panel) and female (bottom panel) age and fork length data and the estimated growth curves.




Figure 13. Scatter plot of age reading from two independent age readers and the expected relationship of age reading between the two age readers.




Figure 14. Standard deviation of ageing error.




Figure 15. Plot of the estimated probability of maturity against the estimated age of female black rockfish. The intervals are the 95% confidence intervals estimated by bootstrapping.





Figure 16. Plot of the estimated probability of maturity against the fork length of female black rockfish. The intervals are the 95% confidence intervals estimated by bootstrapping.







Figure 17. Relationship between fecundity and size (top panel) and fecundity and body weight (bottom panel).



0.5 - (reg of the control of the con

Figure 18. Plot of expected male (top panel) and female (bottom panel) estimated total mortality coefficients against total fishing effort. The estimated intercept in each sub graph was the estimated natural mortality.




Figure 19. Scatter plot of estimated female black rockfish mortality coefficients versus estimated male black rockfish mortality coefficients, and the estimated linear relationship.




Figure 20. Time series plot of the estimated male and female black rockfish total mortalities.




Figure 21. Time series plot of the estimated CPUEs of recreational survey data in all areas from 1990 to 2006. The estimated CPUEs were done by mean estimator and delta lognormal model.




Figure 22. Time series plot of the estimated CPUEs of recreational survey data in Area 2 from 1990 to 2006. The estimated CPUEs were done by mean estimator and delta lognormal model.




Figure 23. Plot of accumulated CWT tag lost rate with time.




Figure 24. Time series of the tagging CPUE of the central Washington coast.




Figure 25. Comparison of natural mortality rates between males and females as defined in the STAT Best Fit Model. In the STAR base model Female natural mortality asymptotes at 0.20 at age 15 instead of 0.24 in the STAT Best Fit model.




Figure 26. Comparison of fishing mortality rates estimated from STAR Base, STAT Best Fit model and the tagging model (assuming M=0.2 for both sexes).




Figure 27. Convergence properties of the STAR base model.




Figure 28. Variation in fork-length at age by sex.




Figure 29. Comparison of growth curves estimated from STAR base, STAT best-fit model and external estimates to the mean size at observed age.




Figure 30. STAR base and STAT "best fit" model fit to tagging abundance (top panel) and tagging CPUE (bottom panel) data by fishery.






Figure 31. Comparison of model results using increasing weighting on the abundance trend time series.



#### Spawning Biomass by Retrospective Year (BestFit1624) Spawning Biomass Year ♦ 1996 △ 2000 • 2002 **x** 2004

Figure 32. Retrospective analysis of the northern black rockfish STAR base (top panel) and STAT best-fit (bottom panel) models. Observation data are ignored and there is no recruitment deviations estimated beyond retrospective year

#### Female whole catch length fits for fleet 1



#### Male whole catch length fits for fleet 1





Figure 33. STAR base model fit to female (top) and male (bottom) length composition samples collected from the trawl fishery.

#### Female whole catch length fits for fleet 2



#### Male whole catch length fits for fleet 2





Figure 34. STAR base model fit to female (top) and male (bottom) length composition samples collected from the sport fishery.

#### Female whole catch length fits for fleet 3



## Male whole catch length fits for fleet 3




Figure 35. STAR base model fit to female (top panel) and male (bottom panel) length composition samples collected from the non-trawl fishery.

# Combined sex whole catch length fits for fleet 6





Figure 36. STAR base model fit to male length composition samples (combined sex) collected from the trawl fishery.

#### Female whole catch age fits for fleet 1



#### Male whole catch age fits for fleet 1





Figure 37. STAR base model fit to female (top panel) and male (bottom panel) age composition samples collected from the trawl fishery.

#### Female whole catch age fits for fleet 2



#### Male whole catch age fits for fleet 2





Figure 38. STAR base model fit to female (top panel) and male (bottom panel) age composition samples collected from the sport fishery.

#### Female whole catch age fits for fleet 3



#### Male whole catch age fits for fleet 3




Figure 39. STAR base model fit to female (top panel) and male (bottom panel) age composition samples collected from the non-trawl fishery.

1000 Recruits (Age-6) Recruits (Age-1) **YEAR** - Current Assessment -- 1999 Assessment -- 1994 Assessment





Figure 40. Comparison of STAR base model recruitment estimates to the previous northern black rockfish assessments (top panel) and to the STAT best-fit model recruitment estimates (bottom panel).






Figure 41. Northern black rockfish STAR base model estimated fishing mortality rates by year and fishery (top panel) and cumulative fishing mortality by year and fishery (bottom panel).




Figure 42. Trawl, sport, non-trawl, and tagging survey selectivity estimated by the STAR base model.




Figure 43. Comparison of STAR base model spawning biomass estimates to the previous two assessments for the northern black rockfish assessment.




Figure 44. Comparison of STAR base model spawning biomass estimates to the STAT best-fit spawning biomass estimates for the Northern black rockfish assessment.




Figure 45. Comparison of stock abundance resulting from the STAT base model (left panel) and the STAT best fit model (right panel) to the Councils' minimum stock size threshold and management target.




Figure 46. Plot of population status in relation to fishery management benchmarks for the STAR base (left panels) and STAT best fit models (right panels). Fspr versus (Bunfished/B40) in top panel and spawning depletion in relation to management target of B40% and B 25% in bottom panel.




Figure 47. Change in total likelihood relative to the STAR base model. Negative values indicate a better fit.




Figure 48. STAR base model sensitivity to increased weight on the tagging CPUE and tagging population estimates of abundance.






Figure 49. Likelihood profile of the STAR base assessment model for different fixed values of the Beverton-Holt steepness parameter (h) and Sigma R. The STAR base and STAT best-fit model had the steepness fixed at 0.6 and Sigma R tuned to 0.35.








Lambda on RecDev and Dev series

Figure 50. Likelihood profile for various components following simultaneous changes in the emphasis (weight) of the Recruitment Dev and Recruitment Dev time series for the STAR base (top panel) and STAT best-fit models (bottom panel).







Figure 51. Likelihood profile for various components of the STAR base model (top panel) and STAT best-fit model (bottom panel) following changes in the emphasis (weight) on the abundance and tag CPUE indices.







Figure 52. Likelihood profile for various components of the STAR base model and the STAT best-fit model following changes in the emphasis (weight) on the age composition for all fisheries.



Lambda on size composition




Figure 53. Likelihood profile for various components of the STAR base model and STAT best-fit model following changes in the emphasis (weight) on the length composition for all fisheries.

### Appendix A: SS2 2.00c Control and Data Files

```
#_data_and_control_files:
                            ageonly.DAT
                                         //
                                                  ageonly.CTL
       #_N_Growth_Patterns
1
       #_N_submorphs
1
       #_N_areas
           1
                  1 1
       #_area_assignments_for_each_fishery_and_survey
#_recruit_design_(G_Pattern_x_birthseas_x_area)_X_(0/1_flag)
1
       #4_single
                     "season,area,and"
                                          growth "pattern, then=1"""
0
       #Allow_recr_distr_interaction
0
       #Allow_migration
0
                            movement
                                           from
                                                  area
                                                          1
                                                                 to
                                                                        area
              season 1
                            (0=no; start age=1; End
                                                          age
                                                                 =1)
2
       #_Nblock_Designs
              #_N_Blocks_per_Pattern
1996 2006 #_begin_and_end_year_for_each_Block_in_Pattern_1
1989 2006 #
0.5
       #_fracfemale
1
       #_submorph_between/within
-1
       #vector_submorphdist_(-1_first_val_for_normal_approx)
                          &c
       Natural Mortality
                                  Maturity
10
       #_natM_amin
15
       #_natM_amax
5
       #_Growth_Age-at-L1
20
       #_Growth_Age-at-L2
0
       #_SD_add_to_LAA
       #_CV_Growth_Pattern
0
       #_maturity_option
1
4
       #_First_Mature_Age
1
       #_parameter_offset_approach
       #MG_Adjustment_method
1
       #_MGparm_Dev_Phase
#_growth_parms
            INIT PRIOR PR_type SD_Prior
                                                  PHASE env-var use_dev dev_minyr
                     STD_4elements_in_Dev_Vector
       dev_maxyr
                                                  Block Block_Fxn
#Females
                                         -2
       0.2
              0.16
                     0.3
                                  0.9
                                                 0
                                                          0
                                                                       0
                                                                                0.5
0.1
                            -1
       0
              0
                     #_Gpattern:_1_Gender:_Female_M1_natM_young
```

```
0.2 0.3 -1 0.9 -2 0 0
                                                        0
-3
      3
                                                               Ο
                                                                       0.5
            0
      #M1_natM_old_4_intermediateages_do_a_linear_interpolation_of_NM_on_age
            34.4 13.5 -1 10 2 0 0
                                                                       0.5
10
      40
                   #M1_Lmin_Body_length_at_Amin_(units_in_cm)
      0
             0
             50.37 49.3 -1 10 2 0 0
30
      70
                                                                       0.5
                   #M1_Lmax_Body_length_at_Amax_(units_in_cm)
      0
            Ω
0.01
      0.4
             0.181 0.1745 -1 0.9 3 0 0
                                                                0
                                                                       0.5
             0
                   #M1_VBK
      0
-3
      3
             0.08
                   0.0622 -1
                              0.9 -2
                                            0
                                                   Ω
                                                                Ω
                                                                       0.5
      0
            0
                  #M1_CV-young_Variability_for_size-at-age_at-
age<=AFIX_(units_are_fraction)Units_CV_or_stddev_depending_on_assigned_value_of_CV_patter
n
                 0.0721 -1
-3
      3
             0.08
                               0.9
                                     - 3
                                            0
                                                   0
                                                                       0.5
                  #M1_CV-old_Variability_for_size-at-age_at-
age>=AFIX2_do_a_linear_interpolation_of_CV_on_mean_size-at-age
#Males
            0.16 0.1 -1
                               0.9 -2 0 0
                                                                0
0.1
      0.2
                                                                       0.5
                   #_Gpattern:_1_Gender:_Male__M1_natM_young
      0
            0
0.1
      0.2
            0.16
                  0.1 -1
                               0.9
                                    -2
                                                                Λ
                                                                       0.5
             0
      0
      #M1_natM_old_4_intermediateages_do_a_linear_interpolation_of_NM_on_age
10
      40
             34.2 15.0 -1
                                0.9
                                      2
                                             Ω
                                                   Ω
                                                          Ω
                                                                       0.5
                                                                Ω
      0
             0
                   #M1_Lmin
                                       2
                                                   Λ
30
      70
             47.3
                  46.6 -1
                                0.9
                                             Λ
                                                          Λ
                                                                Ω
                                                                       0.5
      0
            0
                   #M1 Lmax
            0.191 0.1982 -1
0.1
      0.3
                                0.9
                                      3
                                             0
                                                   0
                                                          0
                                                                0
                                                                       0.5
                   #M1_VBK
      0
             0
             0.07
                   0.06 -1
0 05
      0 25
                                0 9
                                      - 3
                                             Λ
                                                   Λ
                                                          Λ
                                                                Λ
                                                                       0.5
      0
             0
                   #M1_CV-young
             0.07
                  0.0567 -1
                                0.9
                                      -3
                                             0
                                                    0
                                                                 0
-3
      3
                                                          0
                                                                       0.5
      0
            0
                 #M1_CV-old
#Females_wtln_Maturity_fec
            -3
                                                                 0
                                                                       0
      0
1_coefficient_to_convert_L_in_cm_to_Wt_in_kg
            2.768 2.768 -1 0.9 -3
                                             0
                                                   0
                                                          0
-3
      3
                                                                 0
                                                                       0.5
             0
                   #Female_wt-len-2_Exponent_in_female_L-W_conversion
-3
            42.6
                  42.6 -1 0.9 -3 0
                                                   0
                                                          0
                                                                       0.5
      3
                  #Female_Maturity_logistic_inflection
      0
            0
-3
      3
             -0.4 \quad -0.4 \quad -1 \quad 0.9 \quad -3 \quad 0
                                                                       0.5
                   #Female_Logistic_slope
            Ω
      0
-3
      3
            -0.3657 -0.3657 -1 0.9 -3
                                                   0
                                                          0
                                                                 0
                                                                       0.5
            0 #-0.3657Female_eggs/gm_intercept
      Ω
            0.7674 0.7674 -1 0.9 -3
- 3
      3
                                                    Ω
                                                                 Ω
                                                                       0.5
      0
            0 #0.7674Female_eggs/gm_slope
#Male_wtln
            3.80E-05 3.80E-05 -1 99 0.5 0 #Male wt-len-
      3
                                                   - 3
                                                          Ω
                                                                Ω
                                                                       Ω
-3
      0
1_coefficient_to_convert_L_in_cm_to_Wt_in_kg
             2.782 2.782 -1 0.9 -3
                                            0
                                                   0
                                                          0
                                                                       0.5
      0
            0
                   #Male_wt-len-2_Exponent_in_female_L-W_conversion
-4
      4
            0
                   1 -1 0.9 -3 0
                                                   0
                                                          0
                                                                 0
                                                                       0.5
      0
            0
                   #_recrdistribution_by_growth_pattern
                   1 -1 0.9 -3 0
-4
            Λ
                                                   Λ
                                                          Λ
                                                                Λ
                                                                       0.5
      4
                   #_recrdistribution_by_area_1
      0
            0
                   1 -1 0.9 -3 0
                                                   Ω
                                                          Ω
                                                                Ω
-4
      4
            Ω
                                                                       0.5
      n
             Ω
                   #_recrdistribution_by_season_1
                                                   0
-1
      1
            1
                   1 -1 0.9 -3
                                                                       0.5
      Ω
             0
                   #_cohort_growth_deviation
0
      #_custom_MG-env_setup
      #_custom_MG-block_setup
```

<sup>#</sup>\_Spawner-Recruitment

<sup>3 #</sup>\_SR\_function

```
#_LO
       ΗI
               INIT
                      PRIOR PR_type SD
                                              PHASE
1
       15
               12
                       6.7
                               0
                                      10
                                              1
                                                      #log(R0)
0.2
               0.6
                       0.566
                                       0.181
                                              -5
       1
                               2
                                                      #steepness
               0.3
                       0.65
                               0
                                       0.4
                                               -4
                                                      #sigma-r
0
       2
                                                      #env-linkrecruitment-
-5
       5
               Ω
                       0
                                               -3
                               Λ
                                       1
environmental_linkage_coefficient
-5
       5
               0
                       Ω
                               0
                                       1
                                              - 1
       #log(R1)offsetfor_initial_equil_recruitment_relative_to_virgin_recruitment_(usuall
y0)
               0
                       0
                               -1
                                       0
                                               -99
0
       Ω
                                                      #autocorrelation_parameter_for_S-R
0
       #_SR_env_link
       #_SR_env_target_1=devs;_2=R0;_3=steepness
1
       #do_recr_dev: 0=none; 1=devvector;_2=simple_deviations
1
1968
       #Begin RecDevs
2001
       #End_recr_Dev
       #Min_Value4Rec_Dev
-15
       #Max_Value4RecDev
15
3
       #Phaseto
                       begin_Estimation
       #_first_yr_fullbias_adj_in_MPD
1492
#_initial_F_parms
                      PRIOR PR_type SD 0.0001 -1 99
               INIT
#_LO
       ΗI
                                              PHASE
0
       0.6
               0.000
                                               -1
               0.000
                      0.0001 -1
                                       99
0
       0.6
                                               -1
               0.000
                       0.0001 -1
Ω
       0.6
                                       99
                                               -1
#_Q_setup
       A=do
               "power,"
                           "B=env-var," C=extra "SD," "D=devtype(<0=mirror,"
                       "2=cons,"
                                      "3=rand,"
       "0/1=none,"
                                                      4=randwalk); "E=0=num/1=bio,"
       F=err_type
#_A
                       D
                                       F
       В
               C
                               Е
0
       0
               0
                       0
                               1
                                       0
0
       0
               0
                       0
                               1
                                       0
                                       Ω
Ω
       0
               Ω
                       Ω
                               1
       0
               0
                               0
       0
               0
                               0
                                       0
Ω
                       Ω
0
       0
               0
                       0
                               0
                                       Ω
#_Q_parms(if_any)
#_size_selex_types
#_Pattern_DiscardMale_Special
       0
               0
                       0
24
       Ω
                                       2
               Ω
                       Ω
24
       0
               0
                       0
                                       3
5
       0
                       2
                                       4
5
       0
               0
                       6
                                       5
24
       0
               0
                       0
#_age_selex_types
               Discard Male
#_Pattern
                               Special
10
       0
               O
                       Ω
                                       1
10
       0
               0
                       0
                                       2
10
       0
               0
                       0
                                       3
10
       0
               0
                       0
                                       4
10
       0
               0
                       0
                                       5
10
       0
               0
                       0
                               #
                                       6
```

#\_selex\_parms

#_LO	HI	INIT	PRIOR	PR_type	SD	PHASE	env-var	use_dev	dev_min	yr	
	dev_max	xyr	dev_std		Block	Block_F	`xn				
#_size_	_sel:1										
#19	70	45.57	50	1	0.05	2	0	0	0	0	0.5
π⊥Э	0	0		or_logis		2	0	O	O	O	0.5
#0.01	60	6.6	15	1	0.05	2	0	0	0	0	0.5
	0	0		th_for_l							
30	60	46	51.2	-1	0.05	2	0	0	0	0	0.5
6	0	0	2 6	1	0 05	2	0	0	0	0	0 E
-6	4 0	1.6 0	-2.6	-1	0.05	3	U	U	U	U	0.5
-1	9	4	5.2	-1	0.05	-3	0	0	0	0	0.5
	0	0									
-1	9	2.2	6	-1	0.05	-3	0	0	0	0	0.5
0	0	0	2 5		0 05	0	0	0	0	0	۰ -
-8	9 0	-4 0	-3.7	-1	0.05	2	0	0	0	0	0.5
-5	9	-1	0.1	-1	0.05	2	0	0	0	0	0.5
	0	0	0.1	_	0.00	_	ŭ	Ü	Ü	Ü	0.5
#_size_	_sel:2										
		4-									
#19	70 0	45 0	50	1 or_logis	0.05	2	0	0	0	0	0.5
#0.01	60	20	15	1	0.05	2	0	0	0	0	0.5
110.01	0	0		th_for_l			o .	Ü	Ü	Ü	0.5
#											
20	60	41.5	41.2	-1	0.05	2	0	0	0	0	0.5
-6	0 4	0 -4	-2.6	-1	0.05	3	0	0	0	0	0.5
-0	0	0	-2.0		0.03	3	U	U	U	U	0.5
-1	9	3.5	5.2	-1	0.05	-3	0	0	0	0	0.5
	0	0									
-1	9	3	6	-1	0.05	-3	0	0	0	0	0.5
0	0	0	2 5		0 05	0	0	0	0	0	۰ -
-8	9 0	-3.7 0	-3.7	-1	0.05	2	0	0	0	0	0.5
-5	9	-1	0.1	-1	0.05	2	0	0	0	0	0.5
	0	0									
#_size_	_sel:3										
U.1.O	п.	4.1 .	F 0	1	0.05	0	0	0	0	0	۰ -
#19	70 0	41.6	50	1 or_logis	0.05	-2	0	0	0	0	0.5
#0.01	60	0 9.3	15	1	0.05	-2	0	0	0	0	0.5
110.01	0	0		th_for_l				Ü	Ü		0.5
30	60	46	41.2	-1	0.05	2	0	0	0	0	0.5
	0	0									
-6	4	-0.747	-2.6	-1	0.05	3	0	0	0	0	0.5
-1	0 9	0 4.83454	5 2	-1	0.05	3	0	0	0	0	0.5
_	0	0	3.2	_	0.03	5	0	O	O	O	0.5
-1	9	4	6	-1	0.05	3	0	0	0	0	0.5
	0	0									
-8	9	-4	-3.7	-1	0.05	2	0	0	0	0	0.5
-5	0 9	0 2	0.1	-1	0.05	2	0	0	0	0	0.5
-5	0	0	0.1	-1	0.05	2	U	U	U	U	0.5
#_size_		· ·									
1	19	1	5		0.05	-2	0	0	0	0	0.5
-	0	0	#Mirror	_	0 0=	2	0	0	0	0	o =
1	19 0	19	5 #Mirror	-1 Tag	0.05	-3	0	0	0	0	0.5
#_size_		0	#Mirror	ıay							
1	19	1	5	-1	0.05	-2	0	0	0	0	0.5
	0	0	#Mirror	TagCPUE							

1	19	19	5	-1	0.05	-3	0	0	0	0	0.5	
0 0 #MirrorTagCPUE #_size_sel:6												
#1	19	30	5	-1	0.05	2	0	0	0	0	0.5	
#1	0 19	0 19	# 5	-1	0.05	3	0	0	0	0	0.5	
20	0 60	0 39.513	# 41.2	-1	0.05	2	0	0	0	0	0.5	
-6	0 4	0 -3.41	-2.6	-1	0.05	3	0	0	0	0	0.5	
-1	0	0 3.7	5.2	-1	0.05	3	0	0	0	0	0.5	
-1	0 9	0 3.5	6	-1	0.05	3	0	0	0	0	0.5	
-8	0 9	0 -4.69	-3.7	-1	0.05	2	0	0	0	0	0.5	
-5	0	0 -3.95	0.1	-1	0.05	2	0	0	0	0	0.5	
#_age_	0 0 #_age_sel:1											
#0	40	1	5	0	99	2	0	0	0	0	0.5	
#0.01	0	0 2	# 2	3	99	3	0	0	0	0	0.5	
#1	0 20	0 7	# 5	4 0	99	2	0	0	0	0	0.5	
#1	0 25	0 17	# 2	5	99	3	0	0	0	0	0.5	
0 0 # 6 #_age_sel:2												
#1	20	7	5	0	99	2	0	0	0	0	0.5	
#1	0 25	0 17	# 2	5 0	99	3	0	0	0	0	0.5	
0 0 # 6 #_age_sel:3												
#0	40	1	5	0	99	2	0	0	0	0	0.5	
#0.01	0 10	0	# 2	7	99	3	0	0	0	0	0.5	
#1	0 20	0 7	# 5	8 0	99	2	0	0	0	0	0.5	
#1	0 25	0 17	# 2	5 0	99	3	0	0	0	0	0.5	
#_age_	0 sel:4	0	#	6								
#0	40	1	5	0	99	2	0	0	0	0	0.5	
#0.01	0 10	0 2	# 2	9 0	99	3	0	0	0	0	0.5	
1	0 #_Selpa	0 arm_Adju	# .st_Metho	10 od								
0	#_cust	om_sel-e	nv_setu	ō								
0		om_sel-b		tup								
#-6	60	44	-2.6	1	0.05	4						
#-10 #-10	10 10	.0	.1 .1	1 1	0.05 0.05	4 4						
-1	#_selpa	armdev-p	hase									
#_Vari	iance_adjustments_to_input_values											
#_1	2											
0	0 0	0 0	0	0	0		to_surve to_disca					
0	0	0	0	0	0	#_add_t	to_bodyw _by_lenc	rt_CV				

```
2
                             1
                                     1
                                               #_mult_by_agecomp_N
                                               #_mult_by_size-at-age_N
1
               1
                       1
                               1
                                       1
30
        #_DF_for_discard_like
        #_DF_for_meanbodywt_like
30
1
        #_maxlambdaphase
0
       #_sd_offset
#_lambdas_(columns_for_phases)
1
       #_Fishery:_1
       #_Fishery:_2
1
1
       #_Fishery:_3
       #_OSP_CPUE:_4
0
        #_TagAbundance:_5
1
       #_TagCPUE:_6
1
       #_discard:_1
0
0
        #_discard:_2
        #_discard:_3
0
0
        #_discard:_4
        #_discard:_5
0
        #_discard:_6
0
       #_meanbodyweight
0
.1
       #_lencomp:_1
.1
        #_lencomp:_2
        #_lencomp:_3
.1
0
        #_lencomp:_4
        #_lencomp:_5
0
.1
        #_lencomp:_6
       #_agecomp:_1
1
        #_agecomp:_2
1
        #_agecomp:_3
1
0
        #_agecomp:_4
0
        #_agecomp:_5
        #_agecomp:_6
0
1
        #_size-age:_1
1
       #_size-age:_2
0
       #_size-age:_3
        #_size-age:_4
O
        #_size-age:_5
0
        #_size-age:_6
        #_init_equ_catch
1
        #_recruitment Deveations
       #_parameter-priors
1
1
        #_parameter-dev-vectors
1000
        #_crashPenLambda
0.99
       #_maximum_allowed_harvest_rate
999
```

```
#
       SS2
                      File
               Data
1915
        #
               start
                      year
2006
        #
               end
                       year
1
        #
               number seasons
12
        #
               months per
                              season
1
        #
               spawning
                               season
               number of
                               fleets
3
        #
3
        #
               number of
                               surveys
Trawl_1%Sport_2%Line_3%SPTCPUE_4%TagAbun_5%TagCPUE_6 #
                                                             Fleets &
                                                                             Surveys
0.5
       0.5
               0.5
                      0.5
                               0.5
                                      0.5
                                                      Timing of
                                                                     Catch
                                                                            and
                                                                                     Survey
2
        #
               number of
                               genders
40
       #
                                      Plus
               Maximum Age
                               in
                                              Group
#
#
       Landings
#
                               Initial Landings
0
               0
                                                      MT
                                                              Opposite
                                                                             Time
                                                                                     Series
       Estimated
                       Time
                               Series
```

0	0	0	#	1915						
0	0	0	#	1916						
0	0	0	#	1917						
0	0	0	#	1918 1919						
0	0	0	#	1920						
0	0	0	#	1921						
0	0	0	#	1922 1923						
0	0	0	#	1923						
0	0	0	#	1925						
0	0	0	#	1926						
0	0	0	#	1927 1928						
0	0	0	#	1929						
0	0	0	#	1930						
0	0	0	#	1931 1932						
0	0	0	#	1932						
0	0	0	#	1934						
0	0	0	#	1935						
0.00	0	0	#	1936 1937						
0.00	0	0	#	1938						
0.00	1	0	#	1939						
3.2	2.8	0 1	#	1940	Landings	MT	1.4	312	0.4	2
9.2	4.6	0	#	1941	Landings	MT	1.4	312	1.2	3.2
15.2	6.3 4.6	0	#	1942	Landings	MT	0	315.5	0.9	9.2
21.2	8.1 6.3	0	#	1943	Landings	MT	0.6	257.6	0.7	15.2
27.2	9.8 8.1	0	#	1944	Landings	MT	0.1	232.2	0.2	21.2
33.2	11.6 9.8	0	#	1945	Landings	MT	0.2	232.9	1.5	27.2
39.2	13.3 11.6	0	#	1946	Landings	MT	0	188.7	1.1	33.2
52	15.1 13.3	0	#	1947	Landings	MT	0	224.8	1.2	39.2
51.2	16.8 15.1	0	#	1948	Landings	MT	0	221.6	4.3	5.2
57.2	18.6 16.8	0	#	1949	Landings	MT	73.1	259.4	4.8	51.2
63.2	20.3 18.6	1.5 0	#	1950	Landings	MT	9	234.1	15	57.2
69.2	22.1 20.3	2.5 1.5	#	1951	Landings	MT	0	264.2	8.6	63.2
75.2	23.8 22.1	3.5 2.5	#	1952	Landings	MT	3.3	264.8	65.8	69.2
81.2	25.6	4.5	#	1953	Landings	MT	1	358.6	106.3	75.2
87.2	27.3	5.5 4.5	#	1954	Landings	MT	46.8	316.9	88.4	81.2
93.2	29.1	6.5 5.5	#	1955	Landings Landings	MT	71.4	342.9	132.3	87.2
99.2	30.8	7.5 6.5	#	1956	3	MT	46.2	332.3	83.4	93.2
105.2	32.6	8.5 7.5	#	1957	Landings	MT	43.3	387.2	119.4	99.2
111.2	34.3 32.6	9.5 8.5	#	1958	Landings	MT	124.4	369.6	165.3	105.2
117.2	36.1 34.3	10.5 9.5	#	1959	Landings	MT	129	414.2	129.3	111.2
123.2	37.8 36.1	11.5	#	1960	Landings	MT	82 159 6	389.3	220.1	117.2
129.2	39.6 37.8	12.5	#	1961	Landings Landings	MT	158.6	391.1	103	123.2
135.2	41.3 39.6	13.5 12.5	#	1962	rangings	MT	127.3	305.3	272	129.2

141.2	43.1 41.3	14.5 13.5	#	1963	Landings	MT	218.9	302.2	145.8	135.2
108	44.8	15.5	#	1964	Landings	MT	327.5	244.3	134.1	141.2
186	43.1 46.6	14.5 16.5	#	1965	Landings	MT	185.1	135.7	128.9	108
234	44.8	15.5	#	1966	Landings	MT	213	213.8	81.8	186
122	46.6 50.1	16.5 18.5	#	1967	Landings	MT	64.6	144.8	70.5	234
261	48.3 51.8	17.5 19.5	#	1968	Landings	MT	321.3	150.7	104	122
303	50.1 53.6	18.5 20.5	#	1969	Landings	MT	96	94.2	89.8	261
134.1	51.8 55.3	19.5 17.5	#	1970	Landings	MT	15	76	52.2	303
116	53.6 57.1	20.5 29.3	#	1971	Landings	MT	347.2	36.8	32.7	134.1
48	55.3 58.8	17.5 26.8	#	1972	Landings	MT	156	62.3	36.9	116
75	57.1 60.6	29.3 51.2	#	1973	Landings	МТ	75	60.6	51.2	48
	58.8	26.8			_					75
156	62.3	51.2	#	1974	Landings	MT	48	58.8	26.8	
347.2	62.3 62.3	32.7 36.9	#	1975	Landings	MT	116	57.1	29.3	156
15	36.8 36.8	52.2 32.7	#	1976	Landings	MT	134.1	55.3	17.5	347.2
96	76 76	89.8 52.2	#	1977	Landings	MT	303	53.6	20.5	15
321.3	94.2 94.2	104 89.8	#	1978	Landings	MT	261	51.8	19.5	96
64.6	150.7 150.7	70.5 104	#	1979	Landings	MT	122	50.1	18.5	321.3
213	144.8 144.8	81.8	#	1980	Landings	MT	234	48.3	17.5	64.6
185.1	213.8 213.8	128.9 81.8	#	1981	Landings	MT	186	46.6	16.5	213
327.5	135.7 135.7	134.1 128.9	#	1982	Landings	MT	108	44.8	15.5	185.1
218.9	244.3 244.3	145.8 134.1	#	1983	Landings	MT	141.2	43.1	14.5	327.5
127.3	302.2 302.2	272 145.8	#	1984	Landings	MT	135.2	41.3	13.5	218.9
158.6	305.3 305.3	103 272	#	1985	Landings	MT	129.2	39.6	12.5	127.3
82	391.1 391.1	220.1 103	#	1986	Landings	MT	123.2	37.8	11.5	158.6
129	389.3 389.3	129.3 220.1	#	1987	Landings	MT	117.2	36.1	10.5	82
124.4	414.2	165.3 129.3	#	1988	Landings	MT	111.2	34.3	9.5	129
43.3	369.6 369.6	119.4 165.3	#	1989	Landings	MT	105.2	32.6	8.5	124.4
46.2	387.2	83.4	#	1990	Landings	MT	99.2	30.8	7.5	43.3
71.4	387.2 332.3 332.3	119.4 132.3 83.4	#	1991	Landings	MT	93.2	29.1	6.5	46.2
46.8	342.9 342.9	88.4 132.3	#	1992	Landings	MT	87.2	27.3	5.5	71.4
1	316.9	106.3	#	1993	Landings	MT	81.2	25.6	4.5	46.8
3.3	316.9 358.6	88.4	#	1994	Landings	MT	75.2	23.8	3.5	1
0	358.6 264.8	106.3	#	1995	Landings	MT	69.2	22.1	2.5	3.3
9	264.8	65.8 15	#	1996	Landings	MT	63.2	20.3	1.5	0
73.1	264.2	8.6 4.8	#	1997	Landings	MT	57.2	18.6	0	9
	234.1	15								

```
259.4 4.3
0
                       #
                               1998
                                       Landings
                                                       MT
                                                               51.2
                                                                       16.8
                                                                               0
        259.4
               4.8
0
        221.6
               1.2
                               1999
                                       Landings
                                                       MT
                                                               5.2
                                                                       15.1
                                                                               0
       221.6
               4.3
                               2000
                                       Landings
                                                               39.2
0
       224.8
               1.1
                       #
                                                       MT
                                                                       13.3
                                                                               0
        224.8
               1.2
0.2
       188.7
               1.5
                               2001
                                       Landings
                                                       МТ
                                                               33.2
                                                                       11.6
                                                                               0
       188.7
               1.1
       238.9
                               2002
0.1
               0.2
                       #
                                       Landings
                                                       МТ
                                                               27.2
                                                                       9.8
                                                                               Ω
       232.9
               1.5
       237.1
0.6
               0.7
                               2003
                                       Landings
                                                       МТ
                                                               21.2
                                                                       8.1
                                                                               0
       232.2
               0.2
0
       268
               0.9
                               2004
                                       Landings
                                                               15.2
                                                                               0
                                                       MT
                                                                       6.3
       257.6
               0.7
       331.7
                               2005
                                       Landings
                                                               9.2
1.4
               1.2
                                                       MT
                                                                       4.6
                                                                               0
       315.5
               0.9
1.4
        321.5
               0.4
                       #
                               2006
                                       Landings
                                                       MT
                                                               3.2
                                                                       2.8
                                                                               0
       312
               1.2
#
       1.4
               312
                       0.4
       Surveys
#CPUE_from_Area_2_Raw_Means
                               ln(1+cv)
#Year
       Season Type
                       Value
28
                       5.73
                               0.728959186
1990
       1
               4
1991
       1
               4
                       5.426
                               0.703659282
1992
       1
               4
                       4.768
                               0.695933036
1993
       1
                       4.242
                               0.759157379
               4
1994
       1
               4
                       4.426
                               0.740246527
1995
       1
               4
                       4.069
                               0.705679139
#1996
       1
               4
                       4.569
                               0.646320543
#1997
                               0.699568754
                       3.932
       1
               4
#1998
       1
               4
                       4.805
                               0.622019705
#1999
       1
               4
                       4.856
                               0.620031093
#2000
       1
               4
                       5.028
                               0.604528452
#2001
       1
               4
                       4.288
                               0.673016624
                               0.607570313
#2002
               4
                       5.01
       1
#2003
       1
               4
                       4.946
                               0.607124035
                       5.571
                               0.553122333
#2004
       1
               4
#2005
       1
               4
                       5.355
                               0.562373981
#2006
       1
               4
                       5.201
                               0.586151481
#Tag_Abundance_from_Area_2_Raw_Means
       Season Type
#Year
                       Value
                               ln(1+cv)
2000
                               0.0854
               5
                       1389
       1
2001
       1
               5
                       2997
                               0.1157
2002
                               0.0806
                       1944
       1
               5
2003
       1
               5
                       2119
                               0.0624
2004
       1
               5
                       2996
                               0.1107
2005
       1
               5
                       5015
                               0.1276
                       3464
2006
       1
                               0.08
#TagCPUE_from_Area_2_Raw_Means
#Year
       Season Type
                       Value
                               ln(1+cv)
1981
                       4.75
                               0.666
       1
               6
1986
       1
               6
                       2.337
                               0.5993
1987
                               0.6344
       1
               6
                       1.172
1988
                       0.826
                               0.5539
       1
               6
1989
                               0.9771
       1
               6
                       1.236
1990
                       0.991
                               0.8439
       1
               6
1998
       1
               6
                       2.46
                               0.813
1999
                       3.061
                               0.7407
       1
               6
2000
               6
                       2.203
                               0.5684
2001
               6
                       4.657
                               0.6076
       1
2002
       1
               6
                       5.486
                               0.5034
```

73.1

0

0

0

0.2

0.1

0.6

1.4

0

0.5913

6.245

2003

1

```
1 6
1 6
                    9.414 0.5149
10.192 0.7579
10.543 0.4205
2004
2005
      1
2006
      1
             6
#
      Discards
#
2
-1
#
      Mean
             Body
                    Weight
#
0
#
#
      Composition
                    Conditioners
0.0001
0.0001
#
#
      Length Composition
#
#Yr
             Flt/Svy Gender Part Nsamp datavector(female-male)
      Seas
19
20
      22
             24
                    26
                           28
                                  30
                                         32
                                                34
                                                     36
                                                              38
                                                                     40
                                                                            42
             46
                    48
                           50
                                  52
                                         54
                                                56
#
67
      1 1 3 0
0 0 0 0 0.0051
0.0358 0.0396 0 0
1976
                                  28
                                               0
                                                      0
                           0.0051 0.0102 0.0332 0.0652 0.0985 0.0806 0.0793 0.0678
                                  0
                                         0
                                                0
                                                       0.0013 0 0
      0.0026 \ 0.0307 \ 0.0729 \ 0.1036 \ 0.1189 \ 0.0844 \ 0.0473 \ 0.0179 \ 0.0026 \ 0.0026
1980
                                  14 0
                                                             0 0
      1
             1
                    3 0
                                                0
                                                      0
             0.0049 \ 0.0049 \ 0.0388 \ 0.0194 \ 0.034 \ 0.0485 \ 0.0631 \ 0.0825 \ 0.0194 \ 0.0146
      0.0097 0.0097 0
                                                                            0.0049
                         0
                                 0
                                         0
                                                0
                                                      0
                                                              0
                                                                   0
      0.0243 0.034 0.0922 0.1165 0.1359 0.1165 0.0631 0.034
                                                             0.0243 0.0049
1981
             1
                    3
                           0
                                  28
                                         0
                                                0
                                                      0
                                                              0
                                                                     0
      1
                                                              0.0875 0.055
      0.0025 0
                    0.0025 0.0075 0.0225 0.035
                                                                            0.0475
                                                0.0625 0.08
                                                                     0.0025 0.005
      0.0175 0.015 0
                           0
                                  0
                                         0
                                                0
                                                       0
                                                              0
```

0.1

0.0625 0.035

0.01

0.005

0.12

0.0175 0.0575 0.05

0.1

1982	1	1	3	0	28	0	0	0	0	0	0
	0 0.0025	0.0025	0.01	0.025 0	0.0325 0	0.075 0	0.08	0.0625 0	0.0575 0.005	0.0225	0.025 0.01
1983	0.0275 1	0.0875 1	0.1475 3	0.1175 0	0.1 56	0.0625 0	0.0275 0	0.0125 0	0	0	0
1000	0	0	0.0063	0.015	0.0363	0.0625	0.0788	0.0675	0.0738	0.05	0.0388
	0.01 0.0163	0.0038 0.07	0 0.1138	0 0.13	0 0.0963	0 0.0763	0 0.0338	0 0.0075	0.0025 0	0.0025 0.0013	0.0075
1984	1 0	1	3 0.0033	0 0.02	21 0.0367	0 0.0367	0 0.0733	0 0.1	0 0.0633	0 0.0333	0 0.02
	0.0167 0.0433	0 0.0767	0 0.12	0 0.1267	0 0.09	0 0.0567	0 0.0333	0 0.01	0 0.0067	0.0133 0	0.02
1986	1	1	3	0	57	0	0	0	0	0	0
	0 0.0093	0.0031 0.0155	0.0062 0	0.0093	0.059 0	0.0497 0	0.087 0	0.059 0	0.0932 0	0.0994	0.0559 0.0124
1987	0.0217 1	0.0683 1	0.0963	0.1056 0	0.0994 71	0.0311	0.0124 0	0.0031	0	0	0
	0 0.0399	0 0.0299	0	0.0025	0.01	0.0299	0.0623	0.0823	0.1471 0	0.0873 0	0.0599
1000	0.005	0.0249	0.0698	0.1446	0.1172	0.0499	0.0224	0.0125	0	0	
1988	1 0	1 0	3 0	0	18 0.02	0 0.02	0 0.08	0 0.1	0 0.12	0 0.12	0
	0.01 0	0.01	0 0.03	0 0.08	0 0.19	0 0.09	0.01	0 0.03	0	0 0.01	0.01
1989	1	1 0	3	0 0.0044	40 0	0 0.0311	0 0.0489	0.08	0 0.1067	0 0.12	0 0.1022
	0.0889	0.0578 0.0178	0 0.0222	0 0.0533	0	0	0	0 0.0178	0	0.0044	0
1990	1	1	3	0	44	0	0	0	0	0	0
	0 0.0482	0 0.0281	0.004 0	0.008	0.0241	0.0402 0	0.0281 0	0.0723 0	0.0723 0	0.0803	0.0562 0.0161
1991	0.0201 1	0.0442	0.0683	0.0803	0.1044 54	0.1365 0	0.0522 0	0.0161	0	0	0
	0 0.0464	0.0033 0.0166	0.0066 0	0.0066	0.0132	0.0728 0	0.0695 0	0.0861 0	0.0695 0	0.0662	0.0232
	0.0099	0.0397	0.0695	0.0662	0.1026	0.0795	0.0828	0.0464	0.0199	0	
1992	1 0	1 0	3 0	0 0.01	36 0.025	0 0.055	0 0.06	0 0.085	0 0.09	0 0.075	0 0.035
	0.03 0.01	0.03 0.065	0 0.065	0 0.16	0 0.085	0 0.07	0 0.035	0	0	0.005 0	0.01
1993	1	1	3	0 0.008	22 0.024	0 0.048	0 0.032	0 0.064	0 0.056	0 0.064	0 0.08
	0.048	0.008	0	0	0	0	0	0	0	0	0
#1994	0.056 1	0.024 1	0.072 3	0.088	0.128 9	0.104 0	0.08	0.008 0	0.008	0	0
	0 0.0408	0	0.0612	0	0	0.0612 0	0.1224 0	0	0.0408	0.0612 0	0.0204 0
#1995	0 1	0.0408	0.102	0.102 0	0.1224 9	0.102 0	0.102 0	0.0204 0	0	0	0
	0	0	0	0.04	0	0.06	0.02	0.04	0.02	0.04	0
	0.02	0.02	0.06	0.2	0.14	0.12	0.1	0.04	0.02	0	
#1998	1 0	1 0	3 0	0 0.0235	14 0.0235	0 0.0941	0 0.1059	0 0.0941	0 0.0588		0
	0 0.0235	0 0.1294	0 0.1765	0 0.1412	0 0.0588	0 0.0235	0 0	0	0	0	0.0353
#2002	1	1	3	0	8	0	0 0.02	0 0.14	0 0.16	0 0.18	0 0.08
	0.02	0	0	0	0	0	0	0	0	0	0
#1980	0.02	0	0.04	0.12	0.16	0.06	0	0	0	0	0
	0.002 0.0276	0	0	0	0	0	0.1575 0	0.1594 0	0.25 0.002		0.0413 0.0236
1980	0.0394 1	0.0571 2	0.0886	0.1575 0	0.1594 10	0.25 0	0.1496 0	0.0413	0.0276 0	0	0.0014
	0		0.0085				0.0284		0.0512	0.027	
1001	0.0341	0.0612	0.0512	0.1124	0.1607	0.1579	0.0811	0.0256	0	0	
1981	1 0.0206	2 0.0206	3 0	0 0.0206	30 0.0103	0	0 0.1031		0.0103	0.0103 0.0103	

	0	0	0	0	0	0	0	0	0 0206	0.0309	0 0309
	0.0309	0.0722			0.1443			0.0206		0.0305	0.0307
#1982	1	2	0	0	22	0.0088	0.0109	0.0044		0.0073	
	0.0146	0.0248	0.0518	0.0904	0.1196	0.1276	0.1488	0.1145	0.1371	0.0795	0.0343
	0.0904	0.1196	0.1276	0.1488	0.1145	0.1371	0.0795	0.0343		0.0022	0.0310
1982	1	2	3	0	12	0	0	0	0	0	0
	0 0	0.0177	0.0177	0.0442	0.0796 0	0.0619	0.0796	0.0531	0.0265	0.0088	0 0.0354
	0.115	0.0973	0.0619	0.115	0.0973	0.0531	0.0177	0	0	0	0.0331
#1983	1	2	0	0	1	0	0	0	0	0	0.04
	0.0733	0.12	0.1133	0.1733	0.1733	0.12	0.08	0.0467 0.04	0.0133	0.0267 0.12	0.02 0.1133
	0.1733	0.1733	0.12	0.08	0.0467	0.0133	0.0267		0	0	
#1984	1	2	0	0	2	0	0	0	0.1	0	0
	0.2	0.1	0.2	0	0	0.4	0	0	0 0.2	0 0.1	0 0.2
	0	0	0.4	0	0	0	0	0	0	0	
1984	1	2	3	0	25	0	0	0	0.0029		
	0.033	0.0445	0.0603	0.0848	0.0905	0.0876		0.0546	0.0417	0.0158 0.0172	0.0129
	0.0388	0.0704	0.0718	0.0503	0.0431		0.023	0.0086		0	
#1985	1	2	0	0	1	0.1884	0.0072	0	0.0072	0.0217	
	0.1594	0.1377	0.2391	0.1014	0.058 0	0.0145	0.0072	0 0.0435	0.0072 0.1594	0 0.1377	0 0.2391
	0.1014		0.0145	0.0072	0	0.0072	0	0	0	0.0072	
1985	1	2	3	0 0633	3 0.0759	0 0.0443	0 0.0759	0 0.0633	0	0 0.0127	0 0.019
	0.0127	0.019	0.0316	0.0633	0.0759	0.0443	0.0759	0.0633	0.0443	0.0127	0.019
	0.0759	0.0633	0.0759	0.0886	0.1076	0.038	0.019	0.0127	0	0	
1986	1 0.0098	2 0.0117	3 0.0391	0 0.0469	17 0.0723	0 0.0801	0.0938	0 0.0488	0 0.0313	0.002	0.0039
	0.0098	0.0117	0.0391	0.0409	0.0723	0.0001	0.0530	0.0039	0.0313	0.0234	0.0547
	0.0957	0.0957	0.1055	0.0566	0.043	0.0176	0.0078	0.0059	0	0	
1987	1 0.014	2 0.0171	3 0.0295	0 0.0481	21 0.0543	0 0.0698	0 0.0837	0 0713	0 0.0543	0 0341	0.0047
		0.0016	0.0293	0.0401	0.0343	0.0098	0.0037	0.014	0.014	0.0155	0.0202
	0.0326	0.0651	0.0775	0.0713	0.0605	0.0682	0.0155	0.0124		0	
1988	1	2 0.0288	3 0.0421	0.0643	15 0.0576	0.0909	0.0909	0 0.0576	0 0.0443	0.0022	0.0177 0.0089
	0	0.0022	0	0	0	0.0022	0.0022	0.0067	0.0222	0.0111	0.0377
1000	0.0665	0.0798	0.0865	0.0488	0.0421	0.0288	0.0222	0.0022	0.0022	0	0 005
1989	1 0.0025	2 0.0126	3 0.0176	0 0.0579	13 0.0806	0 0.0957	0 0.0957	0 0.0705	0 0.0605	0 0.0403	0.005 0.0076
	0	0	0	0	0	0	0	0.0025	0.0101	0.0327	0.0428
1000	0.0705	0.1083	0.0806	0.0479	0.0378	0.0151	0.005	0	0	0	0 0024
1990	1 0.0138	2 0.0103	3 0.0379	0 0.0828	9 0.1241	0.0828	0 0.0966	0 0.0724	0 0.0138	0.0034	0.0034
	0	0	0	0	0.0034	0	0.0034	0	0	0.0276	0.0483
1991	0.0621 1	0.0897 2	0.0759 3	0.0517	0.0345	0.031	0.0276	0	0	0 0125	0.0083
1991		0.0403								0.0123	
		0.0014		0	0					0.0306	0.0542
1992	0.0708	0.0833	0.0597	0.0556	0.0278 29	0.0236	0.0097			0.0023	0 0001
1002		0.0272									
		0.0011		0	0		0.0045				0.0522
1993	0.059 1	0.0942	0.0749	0.0681	0.042 28	0.0193	0.0125	0.0057	0.0011		0.0163
1993		0.0431								0.0023	
		0.0012		0	0	0	0	0.0116			0.0547
1994	0.0664 1	0.071 2	0.0559	0.0547	0.0233	0.0116	0.0047 0	0.0035	0.0012		0.0035
エノシオ	0.0208		0.0428				0.0567			0.0023	
	0.0046	0	0	0	0	0	0.0012	0.0093	0.0162	0.0579	
1995	0.0752 1	0.0729	0.0706	0.0451	0.0289 26	0.022 0	0.0116	0.0058	0	0 0.0012	0.0037
100		0.0333								0.0012	0.0062
	0.0037		0	0	0	0	0.0012	0.0049	0.0185	0.0357	0.0776
	0.0936	0.0764	0.0665	0.0468	0.0222	0.0086	0.0025	0.0012	0.0025	0	

1996	1	2	3	0	27	0	0	0	0.0012	0.0012	0.006
	0.0193	0.0398	0.0736	0.1049	0.1049	0.0676	0.047	0.0277	0.0181	0.006	0.0012
	0	0	0	0	0	0.0024	0.0024	0.0024	0.0205	0.0507	0.0881
	0.0929	0.1025	0.0651	0.0241	0.0205	0.0084	0	0	0	0.0012	
1997	1	2	3	0	29	0	0	0	0	0	0.0022
	0.0189	0.04	0.0633	0.0756	0.0867	0.0756	0.0533	0.0344	0.02	0.02	0.0056
	0.0044	0.0022	0	0	0	0.0033	0.0022	0.0044	0.0233	0.0356	0.0678
	0.0889	0.11	0.0633	0.0467	0.0289	0.0122	0.0089	0.0022	0	0	
1998	1	2	3	0	40	0	0	0	0	0.0053	0.0106
	0.028	0.0355	0.0559	0.0854	0.0824	0.0695	0.0544	0.028	0.0144	0.0136	0.0008
	0	0.0008	0	0	0	0	0.0023	0.0136	0.0333	0.0438	0.0673
	0.0907	0.0998	0.0703	0.0537	0.0242	0.0128	0.0023	0.0008	0.0008	0	
1999	1	2	3	0	44	0	0	0.0006	0.0012	0.0042	0.0126
	0.0263	0.04	0.0592	0.0813	0.0843	0.0699	0.0562	0.0359	0.0132	0.0036	0.0012
	0.0012	0.0006	0	0	0	0.0006	0.0012	0.0185	0.0281	0.0466	0.0831
	0.095	0.0932	0.0652	0.046	0.0185	0.0048	0.0036	0.0024	0.0012	0.0006	
2000	1	2	3	0	43	0	0	0	0.0012	0.0042	0.0079
	0.0176	0.0382	0.0576	0.0885	0.1091	0.0873	0.0812	0.0303	0.0212	0.003	0.0018
	0.0018	0	0	0	0	0	0.0036	0.0091	0.0188	0.0479	0.0661
	0.08	0.0939	0.0739	0.0315	0.0164	0.0055	0.0018	0	0.0006	0	
2001	1	2	3	0	47	0	0	0	0	0	0.0028
	0.013	0.0355	0.0558	0.0986	0.1189	0.1048	0.0614	0.0271	0.018	0.009	0.0023
	0.0017	0	0	0	0	0	0.0011	0.0056	0.0152	0.0355	0.0716
	0.1071	0.0891	0.0705	0.031	0.0147	0.0056	0	0.0023	0.0011	0.0006	
#2002	1	2	3	0	72	0	0	0.0005	0.0005	0.0005	0
	0.0049	0.0168	0.0481	0.0898	0.1028	0.1141	0.0984	0.0606	0.0292	0.0059	0.0027
	0.0011	0.0005	0	0	0	0.0011	0.0016	0.0027	0.0087	0.0276	0.0573
	0.0957	0.1001	0.0757	0.0335	0.013	0.0043	0.0016	0.0005	0	0	
#2003	1	2	0	0	9	0	0	0	0.0013	0.0077	0.0103
	0.0333	0.0833	0.1308	0.1487	0.2064	0.1603	0.1038	0.0641	0.0282	0.0128	0.0051
	0.0038	0	0	0	0	0.0013	0.0077	0.0103	0.0333	0.0833	0.1308
	0.1487	0.2064	0.1603	0.1038	0.0641	0.0282	0.0128	0.0051	0.0038	0	
2003	1	2	3	0	70	0	0	0	0	0.0005	0.0011
	0.0059	0.0184	0.0394	0.0691	0.1021	0.128	0.0977	0.0551	0.0232	0.0151	0.0059
	0.0016	0	0	0	0.0005	0	0.0005	0.0016	0.0108	0.0286	0.0524
	0.0945	0.1037	0.0729	0.0481	0.0151	0.0038	0.0038	0.0005	0	0	
#2004	1	2	0	0	9	0	0	0	0.0021	0.0042	0.0127
	0.0276	0.0467	0.1083	0.1826	0.1868	0.1571	0.1253	0.0425	0.034	0.0127	0.0212
	0.0127	0.0234	0	0	0	0.0021	0.0042	0.0127	0.0276	0.0467	0.1083
0004	0.1826	0.1868	0.1571	0.1253	0.0425	0.034	0.0127	0.0212	0.0127	0.0234	0 0040
2004	1	2	3	0	57	0	0	0 0 0 0 1	0.0012	0.0018	0.0048
	0.0133	0.023	0.0387	0.0762	0.0992	0.1168	0.0768	0.0581	0.0302	0.0097	0.0054
	0.0018	0.0006	0 0.0762	0.0532	0 0.0169	0.0073	0.0048	0.003	0.0006	0.0254	0.0387
#2005	0.0907 1	0.1071	0.0702	0.0332	9	0.0073	0.0048	0.003	0.0000	0.0012	0.0259
#2003	0.049	0.1037	0.1153	0.1441	0.2046	0.1383	0.0951	0.0548	0.0023	0.0038	0.0239
	0.049	0.0086	0.0029	0.1441	0.2040	0.1383	0.0058	0.0348	0.0231	0.1037	0.1153
	0.1441	0.2046	0.1383		0.0548	0.0023	0.0036	0.0233	0.0029	0.0086	0.1133
2005	1	2	3	0.0551	67	0.0231	0.0113	0.0113	0.0025		0.0042
2003			0.0451				0.1052				
	0.0042		0	0	0	0	0	0.0006			0.0325
	0.08	0.1046					0.0012				
#2006	1	2	0	0	9	0	0.0018		0		0.0159
				0.1752				0.0726			
		0.0018		0.0018		0		0.0159			
		0.1416					0.0142				
2006	1	2	3	0	100	0	0.0006			0.0012	0.0075
	0.01						0.0715				
	0	0.0006		0	0		0.0019				
	0.0858	0.1132	0.1007		0.0261		0.0012			0	
#1980	1	3	0	0	14	0	0	0	0	0	0
	0	0.0104	0.0208	0.0313	0.0938	0.1146	0.1979	0.1563	0.1771	0.1354	0.0521
	0	0.0104		0	0	0	0	0	0		0.0208
							0.1354			0.0104	
#1982	1	3	0	0	5	0	0	0	0	0	0.0345
	0.0345						0.1379				0
	0	0	0	0	0	0	0		0.0345		0.2414
						0.0345		0	0	0	
#1983	1	3	0	0	19	0	0	0	0	0	0.0833
	0.0417	0.0833	0.2083	0.2083	0.2083	0.0833	0.0417	0	0.0417	0	0

	0	0	0	0	0	0	0	0.0833	0.0417	0.0833	0.2083
#1983	0.2083	0.2083	0.0833	0.0417	0 5	0.0417	0	0	0	0	0
	0.03	0.02	0.09	0.06	0.07	0.07	0.02	0.07	0	0	0.01
#1984	0.07 1	0.03	0.07 3	0.12	0.04 7	0.14 0	0.01	0	0	0	0
	0.02 0	0.02 0	0.05 0	0.07	0.07 0	0.09 0	0.03	0.03	0.02	0.01	0 0.05
1986	0.1	0.13	0.1	0.04	0.08 100	0.03	0	0	0	0 0.0019	0
	0.0076 0.0114	0.0133	0.0228	0.038	0.0626	0.0569	0.0911	0.1044		0.0759	0.055 0.0266
4000	0.0361	0.055	0.0778	0.0721	0.0323	0.0285	0.0114	0.0038	0.0019	0.0019	
1987	1 0.0111	3 0.025	3 0.0305	0 0.0319	125 0.0555	0 0.0638	0 0.0749	0 0.0652	0 0.043	0.0014 0.0485	0.0042 0.0236
	0.0097 0.0527	0.0069	0 0.0777	0 0.0707	0 0.0721	0 0.0416	0.0028 0.018	0.0028 0.0125	0.0097	0.0222	0.0458
1988	1 0.0118	3 0.0189	3 0.0142	0 0.0307	76 0.059	0 0.092	0 0.1085	0 0.0896	0 0.0495	0.0024 0.0425	0 0.0165
	0.0142	0.0071	0	0	0	0	0	0	0.0118	0.0142	0.0236
1989	0.0542 1	0.0755	0.0755	0.0495	0.059 53	0.0425	0.0189	0.0165 0	0.0024	0	0.0033
	0 0.0201	0.01	0.0334	0.0301	0.0502 0	0.0301	0.0502 0	0.1137 0	0.087 0	0.0368	0.0368 0.0268
1990	0.0602 1	0.0401	0.0702	0.1104	0.0936 21	0.0502	0.01	0.0134	0.0033	0 0.064	0.088
	0.04	0.032	0.056 0	0.048	0.072 0.024	0.032 0.016	0.024	0.024	0.024	0.032	0.016 0.024
1001	0.04	0.048	0.048	0.024	0.008	0.008	0	0	0.008	0	0.0126
1991	1 0.0358	0.04	0.0589	0.0737	0.08	0.0547	0.0568	0.0379	0.0232	0.0063	0.0063
	0 0.0611	0.0021	0 0.0695	0 0.0337	0 0.0316	0 0.0189	0.0042 0.0105	0.0168 0.0042	0.0295	0.0568 0	0.0589
1992	1 0.0037	3 0.0733	3 0.0989	0 0.0696	49 0.0989	0 0.0586	0 0.0586	0 0.0366	0 0.0293	0.011 0.0073	0.011 0.0073
	0 0.0733	0 0.0513	0 0.0659	0 0.0696	0 0.0256	0 0.011	0.0073 0	0.011 0.0037	0.022 0	0.044 0	0.0513
1993	1 0.0031	3	3	0 0.0772	58	0 0.0525	0 0.071	0	0.0031 0.0247	0.0031 0.0216	0.0123 0
	0	0	0	0	0	0	0.0031	0.0432	0.0123	0.0494	0.0494
1994	0.0648 1	0.0864 3	0.071 3	0.0772 0	0.037 44	0.0031	0	0	0 0.004	0 0.012	0.044
	0.068 0	0.068 0	0.068 0	0.08	0.08 0	0.048 0	0.04 0.016	0.008 0.016	0.008	0.004 0.052	0 0.072
1995	0.056 1	0.072 3	0.068 3	0.052 0	0.012 40	0.02	0	0	0	0	0.0089
	0.0268 0	0.0446 0	0.067 0	0.0714		0.0759 0		0.0268	0.0134	0.0045 0.0357	
1981		0.0804		0.0536	0.0268	0.0089	0	0	0	0 0.0006	
1701	0.0159	0.0416	0.0855	0.1166	0.1255	0.1247	0.137	0.1569	0.117	0.0492	0.0142
	0.1166	0.0004 0.1255	0.1247		0.1569	0.117	0.0492	0.0142	0.0023		
1982	1 0.0142	6 0.0305	0 0.0665	0 0.1322	24 0.1535	0 0.1516	0 0.1504			0.0044 0.0372	
		0.0004 0.1535		0 0.1504						0.0305	0.0665
1983	1 0.06	6	0	0	29	0	0	0.0005	0.0015	0.0045 0.0183	
	0.0005	0	0	0	0.0005	0.0015	0.0045	0.0198	0.06	0.1011	0.1269
1984	1	0.1472 6	0	0	24	0	0.0015	0.0089		0.0297	
	0	0.1441 0	0	0.0015	0.0089	0.0193	0.0297			0.003 0.1441	
#1985	0.1441 1	0.1441 6	0.0966 0	0.0609	0.0416 64			0.003 0.0031	0 0.0025	0 0.006	0.0151
		0.1035	0.166	0.1766	0.1708	0.1387	0.0987	0.0735	0.036		0.0029
		0.1708						0.0131			0.100

```
1986
                             0
                                     103
                                           0.0002 0.0002 0.0007 0.0017 0.005 0.0133
       0.0302 \ 0.067 \ 0.1064 \ 0.1577 \ 0.1761 \ 0.1616 \ 0.1546 \ 0.1116 \ 0.0395 \ 0.0135 \ 0.0041
       0.0007 0
                       0.0002 0.0002 0.0007 0.0017 0.005 0.0133 0.0302 0.067
                                                                                     0.1064
       0.1577 0.1761 0.1616 0.1546 0.1116 0.0395 0.0135 0.0041 0.0007 0
                                                      0.0009 0.0025 0.007 0.0101 0.0216
1987
               6
                      Ο
                              Ω
                                      122
                                            0
       0.0363 0.0877 0.1338 0.1631 0.1739 0.1853 0.134 0.0723 0.018 0.0059 0.0008 0.0002 0 0.0009 0.0025 0.007 0.0101 0.0216 0.0363 0.0877
                                                                                     0.1338
       0.1631 0.1739 0.1853 0.134 0.0723 0.018
                                                      0.0059 0.0021 0.0008 0.0002
                                              0.0003 0.0004 0.0051 0.0105 0.016
1988
                      Ω
                              Ω
                                      103
                                                                                      0.0326
       0.0465 0.0603 0.0869 0.1227 0.1433 0.1745 0.1622 0.1071 0.0416 0.0131
                                                                                     0.0045
       0.0023 0.0006 0.0003 0.0004 0.0051 0.0105 0.016 0.0326 0.0465 0.0603 0.0869
       0.1227 \quad 0.1433 \quad 0.1745 \quad 0.1622 \quad 0.1071 \quad 0.0416 \quad 0.0131 \quad 0.0045 \quad 0.0023 \quad 0.0006
                                                      0.0006 0.0033 0.0081 0.0215
1989
                                      103
                                              Ω
       0.0695 \ \ 0.0993 \ \ 0.1085 \ \ 0.1265 \ \ 0.1362 \ \ 0.1476 \ \ 0.1311 \ \ 0.095 \ \ \ 0.0288 \ \ 0.0105 \ \ 0.0018
       0.0007 0.0006 0 0.0006 0.0033 0.0081 0.0215 0.047 0.0695 0.0993
       0.1265 \quad 0.1362 \quad 0.1476 \quad 0.1311 \quad 0.095 \quad 0.0288 \quad 0.0105 \quad 0.0018 \quad 0.0007 \quad 0.0006
1990
                                      108
                                              0.0004 0.0026 0.0116 0.0211 0.026
                       0
                              0
       0.0766 0.1349 0.1533 0.1321 0.126 0.1061 0.093 0.0684 0.0268 0.0099 0.0026
       0.0012 \ 0.0007 \ 0.0004 \ 0.0026 \ 0.0116 \ 0.0211 \ 0.026 \ 0.0464 \ 0.0766 \ 0.1349 \ 0.1533
                      0.1061 0.093 0.0684 0.0268 0.0099 0.0026 0.0012 0.0007
       0.1321 0.126
                                                      0.0019 0.0023 0.0034 0.0129 0.0278
1998
       1
               6
                       Λ
                              Λ
                                      83
                                           0
                       0.2034 0.2224 0.171 0.107
                                                      0.0468 0.0129 0.0038 0.0008 0
       0.0636 0.12
                              0.0019 0.0023 0.0034 0.0129 0.0278 0.0636 0.12
       Ω
               Ω
                       Ω
                                                                                      0.2034
       0.2224 0.171
                       0.107
                              0.0468 0.0129 0.0038 0.0008 0
                                                                      Ω
                                                                              Ω
                                                      0.0003 0.0003 0.0029 0.0063 0.0173
1999
                                      93
                      0
                              0
                                              0
               6
       0.0434 \ \ 0.0811 \ \ 0.157 \ \ \ 0.2105 \ \ 0.1915 \ \ 0.1432 \ \ 0.0894 \ \ 0.0408 \ \ 0.0109 \ \ 0.004 \ \ \ 0.0009
               0.0003 0
                              0.0003 0.0003 0.0029 0.0063 0.0173 0.0434 0.0811 0.157
       0
       0.2105 \ 0.1915 \ 0.1432 \ 0.0894 \ 0.0408 \ 0.0109 \ 0.004 \ 0.0009 \ 0
                                                                              0.0003
                                                      0.0007 0.0011 0.0011 0.0093 0.0237
2000
                      0
                              0
                                      78
                                            0
       0.0714 0.1104 0.166 0.2302 0.1775 0.1233 0.0567 0.0183 0.0093 0.0011 0
                               0.0007 \ 0.0011 \ 0.0011 \ 0.0093 \ 0.0237 \ 0.0714 \ 0.1104 \ 0.166
               0
                       0
       0.2302 0.1775 0.1233 0.0567 0.0183 0.0093 0.0011 0
                                                             0.0003 0.0016 0.0041 0.0062
2001
               6
                      0
                             0
                                    78
                                            0.0003 0
       0.0212 0.0614 0.1156 0.1911 0.2347 0.1911 0.1141 0.0396 0.0128 0.0041 0.0016
                                      0.0003 0.0016 0.0041 0.0062 0.0212 0.0614 0.1156
                      0.0003 0
       0.0003 0
       0.1911 \ \ 0.2347 \ \ 0.1911 \ \ 0.1141 \ \ 0.0396 \ \ 0.0128 \ \ 0.0041 \ \ 0.0016 \ \ 0.0003 \ \ 0
2002
                                                                     0.0012 0.0017 0.0113
                              Ω
                                      49
                                              Ω
                                                      Ω
                                                              Ω
                      Ω
       0.0237 0.0614 0.1177 0.1955 0.2214 0.1781 0.115 0.0521 0.0135 0.0049 0.0015
       0.0007 0.0005 0 0 0 0.0012 0.0017 0.0113 0.0237 0.0614 0.1177
       0.1955 0.2214 0.1781 0.115
                                      0.0521 \ 0.0135 \ 0.0049 \ 0.0015 \ 0.0007 \ 0.0005
2003
       0.0007 0
                                      0.0006 0.001
                                                      0.0013 0.0043 0.0196 0.0444 0.1013
       0.1739 \quad 0.2486 \quad 0.221 \quad 0.1182 \quad 0.0505 \quad 0.0123 \quad 0.0015 \quad 0.0004 \quad 0.0003 \quad 0
2004
                              0
                                      68
                                              0.0005 0
                                                              0.0005 0.0028 0.0065 0.0138
       0.0242 0.0615 0.136 0.2066 0.2167 0.1753 0.0969 0.0399 0.0137 0.0036 0.0013
       0.0002 0.0002 0.0005 0
                                      0.0005 0.0028 0.0065 0.0138 0.0242 0.0615 0.136
       0.2066 \quad 0.2167 \quad 0.1753 \quad 0.0969 \quad 0.0399 \quad 0.0137 \quad 0.0036 \quad 0.0013 \quad 0.0002 \quad 0.0002
                                              0.0005 0.0005 0.0003 0.001 0.0043 0.0205
2005
                      Λ
                              0
                                      49
       0.0352 \ 0.0777 \ 0.1372 \ 0.197 \ 0.2051 \ 0.1752 \ 0.0972 \ 0.0337 \ 0.0104 \ 0.003 \ 0.0005
                      0.0005 0.0005 0.0003 0.001 0.0043 0.0205 0.0352 0.0777 0.1372
       0.0005 0
               0.2051 \ 0.1752 \ 0.0972 \ 0.0337 \ 0.0104 \ 0.003 \ 0.0005 \ 0.0005 \ 0
       0.197
2006
                                       64
                                                      0.0005 0.0017 0.0025 0.0035 0.0153
                               0
                                              0
       0.038
               0.0824 0.1454 0.1829 0.2063 0.1624 0.0953 0.0445 0.0146 0.003
       0.0003 0.0002 0
                              0.0005 0.0017 0.0025 0.0035 0.0153 0.038 0.0824 0.1454
       0.1829 \quad 0.2063 \quad 0.1624 \quad 0.0953 \quad 0.0445 \quad 0.0146 \quad 0.003 \quad 0.001 \quad 0.0003 \quad 0.0002
```

# Age Composition

#

#Yr Seas Flt/Svy Gender Part Ageerr Lbin\_lo Lbin\_hi Nsamp datavector(femalemale)

```
24
```

```
10
3
                 6
                        7
                                                   11
                                                          12
                                                             13 14
      15
            16
                   17
                         18
                                19
                                      20
                                             21
                                                    22
                                                                       25
                                                          23
      30
      number of
                   unique ageing error matrices
                                                          generate
                                                   to
1
0.5
      1.5
             2.5
                   3.5
                         4.5
                                5.5
                                      6.5
                                             7.5
                                                    8.5
                                                          9.5
                                                                10.5
                                                                       11.5
      12.5
            13.5
                   14.5
                         15.5
                                16.5
                                      17.5
                                             18.5
                                                   19.5
                                                          20.5
                                                                 21.5
                                                                       22.5
                                27.5
      23.5
             24.5
                   25.5
                         26.5
                                      28.5
                                             29.5
                                                    30.5
                                                          31.5
                                                                 32.5
                                                                       33.5
      34.5
             35.5
                   36.5
                         37.5
                                38.5
                                      39.5
                                             40.5
0.4817 \ \ 0.5149 \ \ 0.5481 \ \ 0.5813 \ \ 0.6145 \ \ 0.6477 \ \ 0.6809 \ \ 0.7141 \ \ 0.7473 \ \ 0.7805 \ \ 0.8137 \ \ 0.8469
      0.8801 \ 0.9133 \ 0.9465 \ 0.9797 \ 1.0129 \ 1.0461 \ 1.0793 \ 1.1125 \ 1.1457 \ 1.1789 \ 1.2121
      1.2453 \ 1.2785 \ 1.3117 \ 1.3449 \ 1.3781 \ 1.4113 \ 1.4445 \ 1.4777 \ 1.5109 \ 1.5441 \ 1.5773
      1.6105 1.6437 1.6769 1.7101 1.7433 1.7765 1.7765
#Sampson Below
            2.5
                   3.5
                                5.5
                                             7.5
#0.5
     1.5
                         4.5
                                      6.5
                                                    8.5
                                                          9.5
                                                                 10.5
                                                                       11.5
            13.5
                                16.5
                                             18.5
      12.5
                   14.5
                         15.5
                                      17.5
                                                    19.5
                                                                 21.5
                                                                       22.5
                                                          20.5
      23.5
             24.5
                   25.5
                         26.5
                                27.5
                                      28.5
                                             29.5
                                                    30.5
                                                          31.5
                                                                 32.5
                                                                       33.5
                         37.5
            35.5
                   36.5
                                38.5
                                      39.5
                                             40.5
      34.5
#0.062 0.186 0.310
                  0.435 0.559
                               0.683
                                      0.807
                                             0.931
                                                   1.056 1.180 1.304 1.428
      1.552 1.676 1.801 1.925 2.049 2.173
                                             2.297 2.422 2.546 2.670 2.794
      3.663
                                                  3.788 3.912 4.036 4.160
#
#
53
1976
                              1
                                     -1 -1 14 0 0
      0.0084 0.021 0.0924 0.0504 0.0714 0.0672 0.0588 0.0336 0.0336 0.021 0.0126
      0.0126 0
      0.0168 0.0126 0.0084 0.0042 0.0042 0
                                                               0.0042 0.0126
      0
1980
                                                  14
      1
                   3
                         0
                              1
                                      -1 -1
                                                        0
      0.0205 \ \ 0.0256 \ \ 0.041 \ \ \ 0.0462 \ \ 0.0051 \ \ 0.0359 \ \ 0.041 \ \ \ 0.0513 \ \ 0.0359 \ \ 0.0051 \ \ 0
      1 1 3 0 1 -1 -1 28 0 0 0.0025
0.0127 0.0406 0.0457 0.0457 0.066 0.0635 0.0457 0.0355 0.0178 0.0228 0.0127
1981
      0.0102 \ 0.0051 \ 0.0025 \ 0 \ 0.0025 \ 0.0025 \ 0 \ 0 \ 0 \ 0
```

```
0.0025 0.0152 0.0508 0.0787 0.0457 0.0558 0.0279 0.0381 0.0635 0.0533
      0.0152 0.0152 0.0228 0.0178 0.0076 0.0051 0.0051 0.0102 0.0025 0.0127 0.0127
      0.0076
1982
                                    -1
                                          -1
                                                2.1
      1
      0.0576 \ 0.061 \ 0.0814 \ 0.0373 \ 0.0373 \ 0.0169 \ 0.0305 \ 0.0305 \ 0.0102 \ 0.0034 \ 0.0169
      0.0034 0.0102 0 0 0.0034 0 0 0
           0.0034 0.0814 0.0814 0.1186 0.061 0.0339 0.0305 0.0203 0.0305 0.0305
      0.0237 0.0169 0.0068 0.0102 0.0169 0
                                          0 0.0068 0.0068 0.0034 0.0034
      0.0034
1983
                                    -1
                                          -1
                                                56
                                                       0.0013 0
      1
      0.0277 0.0806 0.0529 0.0856 0.0516 0.0428 0.029 0.0189 0.0151 0.0101 0.0063
      0.0025 0 0 0.0013 0 0 0.0025 0.0038 0 0.0013 0
           0.0126 0.0416 0.0957 0.0642 0.0844 0.0592 0.0302 0.0327 0.0277 0.0227
      0.0189 0.0176 0.0113 0.0063 0.0038 0.005 0.005 0.0038 0.0038 0.0013 0.0038
      1984
      0.0269 0.0067 0.0034 0.0034 0.0034 0
                                          0
                                                0 0 0 0
              0.0101 0.037 0.0606 0.0539 0.0842 0.0673 0.0404 0.037 0.0303
      0.0168 \ \ 0.0202 \ \ 0.0269 \ \ 0.0135 \ \ 0.0135 \ \ 0.0101 \ \ 0.0269 \ \ 0.0034 \ \ 0 \\ \\ 0.0101 \ \ 0.0168
      0.0168
1986
                                    -1
                                          -1
                                                57
      0.0031 0.0031 0.0031 0.0031 0.0031 0
      0.0218 0.0031 0
                                                                   0.0031
      0
1987
      1
                        Ω
                              1
                                    -1
                                          -1
                                                71
                                                      0
                                                             0
      0.0075 \ \ 0.0249 \ \ 0.0723 \ \ 0.0848 \ \ 0.0574 \ \ 0.0673 \ \ 0.0648 \ \ 0.0499 \ \ 0.0324 \ \ 0.0349 \ \ 0.0125
      0.0698 0.0873 0.0499 0.0374 0.0549 0.0324 0.0175
            0.015 0.01 0.0025 0.005 0.0025 0.0025 0
      0.02
                                                           0.0025 0.0075
                                                      0
      0.0025
      1
n
1988
                        Ω
                              1
                                    -1
                                          -1 18
                                                      Ω
                                                             Ω
            0.0202 0.0202 0.1212 0.0505 0.0808 0.0505 0.0303 0.0404 0.0303 0.0202
      0.0202 0 0.0202 0 0 0.0101 0 0
                 0.0101 0.0101 0.0404 0.0707 0.0707 0.0303 0.0808 0.0303 0.0101
      0 0
      0.0404 0.0101 0.0202 0 0.0101 0.0101 0.0202 0
                                                       0.0202 0
      0
1989
                        0
                                    -1
                                          -1
                                                 40
      0.0179 0.0268 0.0848 0.0938 0.0982 0.1027 0.067 0.0625 0.0357 0.0179 0.0179
      0.0045 0 0.0089 0.0045 0 0 0 0 0
                                                                   Ω
      0 0.0089 0.0089 0.0357 0.0179 0.0893 0.0804 0.0357 0.0134 0.0313 0.0134
      0.0134 0 0.0089 0
                             Ω
                                    Ω
                                          Λ
                                                 Ω
                                                       Ω
                                                             Λ
      0
1990
                                                44
      1
                                    - 1
                                          - 1
      0.004 \quad 0.0361 \quad 0.0482 \quad 0.0482 \quad 0.0562 \quad 0.0602 \quad 0.0643 \quad 0.0402 \quad 0.0241 \quad 0.012 \quad 0.0281
      0.0161 0
      0.0241 0.0321 0.008 0.012 0
                                   Ο
      Ω
1991
      1
                                     -1
                                           -1
                                                 54
      0.0133 0.0365 0.0664 0.0797 0.0565 0.0532 0.0565 0.0365 0.0266 0.0199 0.0133
                  0 0.0033 0 0.0033 0 0 0
                 0.0066 0.0299 0.0631 0.0399 0.0532 0.0399 0.0465 0.0399 0.01
      Λ
           Ω
      0.0199 \ \ 0.0332 \ \ 0.0199 \ \ 0.0133 \ \ 0.01 \quad \  \  0.0299 \ \ 0.0166 \ \ 0.0166 \ \ 0.01 \quad \  \  0.0066 \ \ 0.0166
      0
1992
                  3
                                    -1
      1
                        Ω
                              1
                                           _ 1
                                                 36
                                                       Ω
                                                             Λ
                                                     0.015
      0.015
            0.055
                 0.08
                        0.135
                              0.05
                                    0.04
                                           0.025
                                                0.05
                                                            0.005
                                                                   0.005
                        0
                                    0
      0.01
            0.01
                  0
                              Ω
                                           0
                                                 0
                                                       0
                                                             0
                                                                   Ω
      Ω
            Ω
                  0.02
                        0.03
                              0.11
                                     0.115
                                          0.06
                                                 0.015 0.04
                                                             0.03
                                                                   0.045
      0.015
           0.01
                  0.015
                       0
                              Ω
                                    Ω
                                           Ω
                                                 0
                                                       Ω
                                                             Ω
      Λ
1993
      1
                                     -1
                                           -1
                                                 2.2
      0.016
                                                             0.016
            0.056
                  0.04
                        0.072
                              0.024
                                    0.056
                                                0.048
                                          0.08
                                                      0.008
                                                                   0.008
      0.008
            0
                  0
                        0
                              0
                                     0
                                           0
                                                 0
                                                       0
                                                             0
                        0.024
                              0.112
                                    0.088
                                           0.088 0.048
                                                      0.04
                                                             0.048
                                                                   0.024
      0
            0
                  0
      0.024
            0.032
                  0.016
                                     0
                                           0.016 0
                                                       0.008 0
                        0
                              0
      0
1994
      1
                                     -1
                                           -1
      0.0208 0.0625 0.0208 0.0208 0.0417 0.0625 0.0625 0.0417 0 0.0208 0
```

```
    0.0208
    0.0208
    0
    0.0208
    0
    0
    0
    0
    0
    0
    0
    0

    0
    0
    0
    0
    0.0417
    0.0833
    0.0625
    0
    0.0625
    0.0625
    0.0625
    0.0417

    0.0208
    0.0625
    0.0625
    0
    0
    0
    0.0208
    0.0208
    0.0417

       0.0208
       1995
                     0.0204 0.0408 0.0204 0 0.0204 0 0
                                                                        0.0204 0
       0.0612 0
       Ω
       0.0192 0.022 0.0275 0.0412 0.022 0.0165 0.0467 0.0192 0.011 0.0027 0.0027 0.0011 0.0082 0.0055 0.0055 0.0027 0.0027 0.0027 0.0027 0.0027 0.0027
1980
       0.011 0.0082 0.0055 0.0055 0.0027 0.0027 0.0027 0.0027 0.0055 0.0027 0.011 0.0055 0.0247 0.033 0.0687 0.0412 0.033 0.0412 0.0467 0.0357 0.011
       0.033 \quad 0.0495 \quad 0.0275 \quad 0.0275 \quad 0.033 \quad 0.0302 \quad 0.0165 \quad 0.0247 \quad 0.011 \quad 0.011 \quad 0.0522
       0.0275
1981
                                            -1
                                                           12
                                                                  0.0141 0.0141 0
                                                    -1
       0.0423 0.0141 0.0141 0.0282 0.0423 0.0563 0.0282 0.0141 0.0282 0
              0 0 0 0 0 0 0 0 0
              0.0282 \ 0.0282 \ 0.0141 \ 0.1408 \ 0.0986 \ 0.0423 \ 0.1268 \ 0.0282 \ 0.0423 \ 0.0141
       0.0423 0.0141 0.0141 0.0282 0 0.0282 0.0141 0
                                                                 0
                                                                          Λ
1984
                                                                         0.0086 0.0375
                             Ω
                                    1
                                            - 1
                                                   - 1
                                                          134
                                                                  Ω
       0.0893 0.1081 0.0634 0.0605 0.0879 0.049 0.0288 0.0245 0.0115 0.013 0.013
       0.0072 0.0072 0.0029 0 0.0014 0.0029 0 0
                                                                 0
                                                                         Ω
       0.0029 \ \ 0.0029 \ \ 0.0303 \ \ 0.0346 \ \ 0.0331 \ \ 0.0418 \ \ 0.049 \ \ \ 0.0346 \ \ 0.0173 \ \ 0.0159 \ \ 0.0144
       0.0202 \ 0.013 \ 0.0173 \ 0.0029 \ 0.0043 \ 0.0029 \ 0.0086 \ 0.0043 \ 0.0086 \ 0.0088 \ 0.0013
       0.0058
1985
                                            -1
                                                   -1
                                                          14
       0.0633 \quad 0.038 \quad 0.0886 \quad 0.0443 \quad 0.0316 \quad 0.0316 \quad 0.0506 \quad 0.0316 \quad 0.0253 \quad 0.0127 \quad 0
       0.0063 0
                     0.0063 0.0063 0.0063 0 0
                                                          0
                                                                 0
                                                                          0
              0.0127 \ 0.0316 \ 0.0633 \ 0.0759 \ 0.0316 \ 0.0253 \ 0.0253 \ 0.0253 \ 0.0633 \ 0.038
       Λ
              0.0316 0.0316 0.0063 0.0127 0.038 0
                                                          0.0063 0 0 0.0063
       Ω
                                                                        0.0079 0.0237
1986
                                   1
                                            -1
                                                   -1
                                                          91
                                                                 0
       0.0632 \ 0.0731 \ 0.0751 \ 0.0593 \ 0.0514 \ 0.0296 \ 0.0296 \ 0.0217 \ 0.0119 \ 0.0079 \ 0.0059
       0.004 0.0059 0.002 0 0 0 0 0.002 0 0 0 0.0079 0.0356 0.0711 0.0909 0.0889 0.0534 0.0356 0.0237 0.0138 0.0198 0.0138
       0.0099 0.0138 0.0119 0.0138 0.002 0.0059 0.0059 0
                                                                 0.002 0.002 0.004
       1 2 3 0 1 -1 -1 112 0 0.0093 0.0125 0.0327 0.0545 0.0654 0.0607 0.0607 0.0327 0.0452 0.028 0.0405 0.0171 0.0156
1987
       0.0109 0.0078 0.0125 0.0016 0.0016 0.0016 0 0 0.0031 0
       0.0078 0.0171 0.0234 0.0498 0.0452 0.0374 0.0312 0.0405 0.0249 0.0249 0.0389 0.0296 0.0093 0.0156 0.0171 0.0171 0.014 0.0125 0.0016 0.0031 0.0031 0.0156
       0.0062
                                                                 0 0.0067 0.0201
       0
                                   1 -1 -1
1988
       0.0022 0.0223 0.0424 0.067 0.0513 0.0469 0.058 0.0424 0.0201 0.0179 0.0156
       0.029 0.0045 0.0089 0.0045 0.0067 0.0067 0.0045 0.0045 0.0022 0
                                                                                0.0045
                                                   -1
                                                          71
1989
                            0
                                           -1
                                                                         0
       0.0658 \ \ 0.0759 \ \ 0.0911 \ \ 0.0709 \ \ 0.0608 \ \ 0.0405 \ \ 0.0456 \ \ 0.0228 \ \ 0.0177 \ \ 0.0228 \ \ 0.0076
       0.0101 \ 0 \qquad 0.0076 \ 0.0051 \ 0.0025 \ 0.0076 \ 0.0025 \ 0.0025 \ 0.0025 \ 0
       Ω
1990
       1
                                            -1
                                                   -1
                                                          51
       0 0 0 0 0 0 0 0 0 0 0 0 0.0035
              0.0208 0.0554 0.0623 0.0761 0.0415 0.0484 0.0242 0.0242 0.0208 0.0173
       Ω
       0.0104 0.0104 0.0104 0 0.0069 0
                                                   0.0104 0.0035 0.0035 0 0.0069
       Ω
                                                          121 0.0042 0.0195 0.0432
1991
                     3
                             0
                                    1
                                           -1
                                                   -1
       0.0669 \quad 0.0628 \quad 0.1144 \quad 0.0753 \quad 0.0474 \quad 0.0223 \quad 0.0237 \quad 0.0237 \quad 0.0153 \quad 0.0112 \quad 0.0056
       0.0139 \ \ 0.0056 \ \ 0.0126 \ \ 0.0084 \ \ 0.0056 \ \ 0.0014 \ \ 0.0042 \ \ 0.0042 \ \ 0.0042 \ \ 0.0014 \ \ 0.0028
       0.0028
```

```
1992
                                   1
                                           -1
                                                   -1
                                                          157 0.0011 0.0125 0.0227
       0.0409 \ 0.0727 \ 0.0864 \ 0.0841 \ 0.067 \ 0.033 \ 0.0307 \ 0.0216 \ 0.0136 \ 0.0148 \ 0.0045
       0.0034 0.0057 0.0011 0.0011 0
                                            0.0011 0.0011 0 0
                                                                          0
       0.0114 \ \ 0.0318 \ \ 0.0295 \ \ 0.067 \ \ \ 0.0693 \ \ 0.0614 \ \ 0.0398 \ \ 0.0295 \ \ 0.0227 \ \ 0.025 \ \ \ 0.0182
       0.0182 \ 0.0091 \ 0.008 \ 0.0125 \ 0.0034 \ 0.0034 \ 0.0011 \ 0.0023 \ 0.0023 \ 0.0057 \ 0.0068
1993
                                                          154
                                                                         0.0082 0.0397
                                            -1
                                                    -1
       0.0806 \ 0.0771 \ 0.0993 \ 0.0993 \ 0.0561 \ 0.0409 \ 0.0129 \ 0.0175 \ 0.0117 \ 0.014 \ 0.007
       0.0023 0.0023 0.0047 0.0012 0.0012 0 0 0 0.0012 0 0
0.0082 0.0234 0.0666 0.0537 0.0806 0.0584 0.035 0.0199 0.0187 0.0129 0.0023
       0.0058 0.007 0.0117 0.0023 0.0035 0.0035 0.0035 0
                                                                 0
                                                                          0.0012 0.0035
       0.0012
1994
                                     1
                                            -1
                                                    -1 155 0
       0.0615 0.0893 0.08 0.058 0.0545 0.0394 0.0244 0.0174 0.0162 0.0128 0.0151
       0.0081 0.0058 0.0046 0.0035 0.0012 0 0 0 0.0012 0.0023 0.0012
       0.007 0.0383 0.0638 0.0615 0.0464 0.0487 0.0441 0.0302 0.022 0.0232 0.022
       0.0128 \ 0.0058 \ 0.0046 \ 0.0023 \ 0.0093 \ 0.0023 \ 0.0046 \ 0.0046 \ 0.0023 \ 0
       0.0023
                                                                 0.0012 0.0025 0.0395
       1995
       0.0012 \ 0.0012 \ 0.0025 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0
       0.0049 \ \ 0.0456 \ \ 0.0691 \ \ 0.0937 \ \ 0.0654 \ \ 0.0469 \ \ 0.0358 \ \ 0.0222 \ \ 0.0222 \ \ 0.0037 \ \ 0.0086
       0.0062 0.0086 0.0062 0.0049 0.0037 0.0025 0.0037 0 0.0012 0.0012 0
       0.0025
1996
                                                           147
                                                                 0
                             Ω
                                            -1
                                                    -1
       0.0836 \quad 0.1358 \quad 0.0727 \quad 0.0764 \quad 0.0364 \quad 0.0255 \quad 0.0255 \quad 0.0097 \quad 0.0024 \quad 0.0012 \quad 0.0048
       0.0073 0.0061 0.0097 0.0048 0.0024 0.0024 0
                                                           0.0012 0.0012 0.0012 0.0036
       0 0012
1997
                                             -1
                                                    -1
                                                           159
       0.047 \quad 0.0761 \quad 0.0873 \quad 0.0672 \quad 0.0571 \quad 0.0493 \quad 0.0246 \quad 0.0202 \quad 0.0112 \quad 0.0123 \quad 0.0056
       0.0056 \ 0.0056 \ 0.0022 \ 0 \ 0.0022 \ 0 \ 0 \ 0.0011 \ 0.0022 \ 0
       1998
                                                           220
                      3
                             Ω
                                            - 1
                                                    -1
                                                                   Ω
                                                                          0.0182 0.0174
                                     1
       0.0477 \ \ 0.0795 \ \ 0.0811 \ \ 0.0795 \ \ 0.0462 \ \ 0.0311 \ \ 0.0364 \ \ 0.0159 \ \ 0.0106 \ \ 0.0061 \ \ 0.0038
       0.0038 0.0015 0 0.0015 0.0008 0.0008 0.0008 0 0.0008 0
       0.0159 \ \ 0.0265 \ \ 0.0386 \ \ 0.0644 \ \ 0.0909 \ \ 0.0742 \ \ 0.0508 \ \ 0.0326 \ \ 0.0197 \ \ 0.0189 \ \ 0.0129
       0.0167 \ 0.0083 \ 0.0076 \ 0.0076 \ 0.0076 \ 0.0053 \ 0.0045 \ 0.0053 \ 0.0015 \ 0.0023 \ 0.0045
       Ω
1999
                                                    -1
                                                           240
                                                                  0.0024 0.0127 0.0453
       0.0375 \ \ 0.0689 \ \ 0.0943 \ \ 0.0647 \ \ 0.058 \ \ \ 0.029 \ \ \ 0.0314 \ \ 0.0169 \ \ 0.0115 \ \ 0.0085 \ \ 0.0048
       0.0006 0.0018 0.0012 0 0.0012 0.0006 0.0006 0.0006 0
                                                                          0
       0.0133 \ 0.0538 \ 0.0417 \ 0.0749 \ 0.0725 \ 0.0616 \ 0.0508 \ 0.0302 \ 0.0242 \ 0.0169 \ 0.0139
       0.0109 0.006 0.0066 0.0018 0.006 0.006 0.0048 0.0036 0.0006 0.0018 0.0036
                                                                 0.0006 0.003 0.0231
2000
       1
                                   1
                                        -1
                                                    -1
                                                          233
       0.0669 0.0681 0.0748 0.0809 0.0596 0.0663 0.0383 0.0255 0.0152 0.0085 0.0043
       0.0061 0.003 0.003 0.0006 0 0.0018 0 0 0.0006 0 0.0006
0.0043 0.017 0.062 0.0669 0.0724 0.0408 0.0408 0.0347 0.028 0.0152 0.0158
0.0073 0.0091 0.0049 0.0073 0.0036 0.0024 0.0043 0.0006 0.0043 0.0024 0.0036
2001
                             0
                                     1
                                            -1
                                                    -1 254
                                                                 0
                                                                          0.0034 0.0141
       0.0401 0.1073 0.0915 0.0554 0.0446 0.0542 0.0475 0.0362 0.0158 0.0136 0.0073
       0.0068 0.0045 0.0023 0.0023 0.0017 0.0006 0 0 0
       0.0028
                             0
                                            -1
2002
                                     1
                                                    -1
                                                          268
                                                                 0.0005 0.0022 0.0146
       0.0304 \ 0.0781 \ 0.1204 \ 0.0765 \ 0.0613 \ 0.051 \ 0.0418 \ 0.0385 \ 0.0211 \ 0.0125 \ 0.0108
       0.0049 0.0043 0.0016 0.0033 0.0005 0
                                                    0.0011 0 0.0005 0 0.0011
       0.0016 \ 0.0179 \ 0.0396 \ 0.0667 \ 0.0743 \ 0.0521 \ 0.0445 \ 0.0233 \ 0.0222 \ 0.0184 \ 0.0152
       0.0119 \ 0.0081 \ 0.0065 \ 0.0054 \ 0.0016 \ 0.0022 \ 0.0016 \ 0.0016 \ 0.0011 \ 0.0016 \ 0.0033
                                                                0
2003
                                            -1
                                                    -1
                                                          261
                                                                        0.006 0.0136
       0.0462 \ 0.062 \ 0.1011 \ 0.0989 \ 0.0609 \ 0.0527 \ 0.0326 \ 0.0304 \ 0.0207 \ 0.0141 \ 0.0092
       0.0038 \ 0.0033 \ 0.0022 \ 0.0011 \ 0.0022 \ 0 \qquad 0.0005 \ 0 \qquad 0 \qquad 0 \quad 0.0016
       0.0168 \ \ 0.0125 \ \ 0.0543 \ \ 0.0554 \ \ 0.0723 \ \ 0.0609 \ \ 0.0321 \ \ 0.0321 \ \ 0.0201 \ \ 0.0168 \ \ 0.0158
```

```
0.0109 \ \ 0.0071 \ \ 0.0043 \ \ 0.0065 \ \ 0.0016 \ \ 0.0043 \ \ 0.0016 \ \ 0.0016 \ \ 0.0027 \ \ 0.0016 \ \ 0.0038
       0.0016
2004
                              0
                                               -1
                                                      -1
                                                              233
                                                                     0.0006 0.0116 0.0469
       0.0451 \ \ 0.084 \ \ \ 0.0572 \ \ 0.0767 \ \ 0.0682 \ \ 0.0627 \ \ 0.0329 \ \ 0.0238 \ \ 0.0171 \ \ 0.0122 \ \ 0.0073
       0.0043 0.0024 0.0018 0.0006 0 0
                                                                   0.0012 0
                                                      0.0006 0
        0.0079 \ \ 0.0359 \ \ 0.0438 \ \ 0.0688 \ \ 0.0542 \ \ 0.0487 \ \ 0.039 \ \ \ 0.0451 \ \ 0.0183 \ \ 0.0164 \ \ 0.0152
       0.014 \quad 0.0091 \quad 0.0049 \quad 0.0037 \quad 0.0024 \quad 0.0024 \quad 0.0018 \quad 0.0024 \quad 0.003 \quad 0.0006 \quad 0.0018
        0.0024
2005
                              Ω
                                                              233
                                                                      Ω
                                                                              0.0037 0.0331
        1
                                      1
                                              - 1
                                                      - 1
        0.0893 0.06
                       0.1118 0.055 0.0687 0.0537 0.0331 0.0175 0.0162 0.0156 0.0119
        0.0094 0.0056 0.0019 0 0.0012 0.0012 0.0006 0
                                                                    0.0006 0.0006 0
        0.0012 0.0131 0.0562 0.05
                                     0.0687 0.045 0.0294 0.0437 0.0231 0.0131 0.0144
        0.0081 \ 0.0112 \ 0.0056 \ 0.0044 \ 0.0025 \ 0.0044 \ 0.0031 \ 0.0012 \ 0.0025 \ 0.0012 \ 0.005
       0.0019
2006
                                               -1
                                                       -1
                                                              212
        0.0647 \ \ 0.1005 \ \ 0.0742 \ \ 0.062 \ \ \ 0.0432 \ \ 0.0512 \ \ 0.0297 \ \ 0.0169 \ \ 0.0088 \ \ 0.0108 \ \ 0.0054
               0.0047 0
                               0.0034 0
                                               0.0007 0
                                                              0
                                                                      0
                                                                              0
        0.0027 0.0121 0.0512 0.0829 0.0722 0.0762 0.0418 0.0384 0.0236 0.0236 0.0128
        0.0108 \ \ 0.0088 \ \ 0.0088 \ \ 0.0061 \ \ 0.0067 \ \ 0.0061 \ \ 0.0027 \ \ 0.0013 \ \ 0.0007 \ \ 0.0007 \ \ 0.0047
        0.002
1983
       1
                       3
                               Λ
                                       1
                                               -1
                                                       -1
                                                              7
                                                                      Ω
                                                                              Λ
                                                                                      0.03
                               0.09
        0.03
                      0.07
                                       0.03
                                               0.03
                                                       0.01
                                                              0.01
                                                                      0.02
                                                                              0.01
               0.1
               0.01
                               0
        Ω
                                       0
                                               0
                                                      0
                                                              0
                                                                      0
                                                                                      Ω
                       0
                                                                              Ω
        n
               0.03
                       0.03
                               0
                                       0.05
                                               0.03
                                                       0.02
                                                              0.07
                                                                      0.05
                                                                              0.02
                                                                                      0.01
        0.04
               0.02
                      0.04
                               0.03
                                       0.02
                                               0.03
                                                      0.01
                                                              0
                                                                      0.02
                                                                                      0.02
                                                                              Ω
        0.04
1984
       1
                               0
                                               -1
                                                       -1
                                                                      0
                                                                              0
                                                                                      0.0101
                       3
                                       1
        0.0404 \ \ 0.0505 \ \ 0.0808 \ \ 0.0404 \ \ 0.0404 \ \ 0.0202 \ \ 0.0505 \ \ 0.0303 \ \ 0.0101 \ \ 0.0202 \ \ 0.0101
               0.0101 0 0 0 0
                                                      0 0 0 0
               0.0202 \ 0.0303 \ 0.0404 \ 0.0303 \ 0.0808 \ 0.0808 \ 0.0404 \ 0.0202 \ 0.0505 \ 0.0303
        Λ
        0.0303 0.0303 0.0101 0
                                       0
                                               0.0101 0.0202 0.0202 0.0101 0.0202 0.0101
        0.0101
       1 3 3 0 1
0.0267 0.0552 0.0629 0.059 0.04
1986
                                              -1
                                                      -1
                                                             99
                                                                     Ω
                                                                              0.0057 0.0133
                                              0.019 0.0686 0.0838 0.061 0.0457 0.0362
                                              0 0 0 0.0019 0.0057 0
              0.0229 0.0114 0.0019 0
        0.019
               0.0152 \ 0.0171 \ 0.04 \ 0.04 \ 0.0362 \ 0.0343 \ 0.0057 \ 0.0419 \ 0.019 \ 0.021
        0.0114 0.0152 0.0057 0.0095 0.0114 0.0038 0.0057 0.0038 0.0038 0.0019 0.0114
        0.0114
1987
                                               -1
                                                      -1
                                                             124
                                                                     0
                                                                              0.0042 0.0139
       0.0473 \quad 0.0445 \quad 0.0612 \quad 0.0695 \quad 0.0376 \quad 0.0362 \quad 0.0403 \quad 0.0292 \quad 0.0389 \quad 0.0209 \quad 0.0223
       0.0236 0.0056 0.0139 0.0181 0.0056 0.007 0.007 0.0028 0.0028 0
       0.0111
1988
                                               -1
                                                       -1
       0.0264 \ \ 0.0553 \ \ 0.0625 \ \ 0.0721 \ \ 0.0673 \ \ 0.0769 \ \ 0.0505 \ \ 0.0337 \ \ 0.024 \ \ \ 0.024 \ \ \ 0.0144
        0.0096 0.0048 0.0024 0
                                       0 0.0024 0 0 0.0024 0
       0.0024 0.0264 0.0216 0.0481 0.0457 0.0457 0.0505 0.0361 0.0313 0.0216 0.0192 0.0168 0.0096 0.012 0.0048 0.0096 0.0144 0.0024 0.0024 0.0024 0.0072 0.0048
1989
                              Ω
                                              - 1
                                                             53
                                                                     Ω
                                                                             0.0034 0.0135
       1
                       3
                                      1
                                                      - 1
              0.0505 0.0471 0.0539 0.0438 0.0606 0.0471 0.0337 0.0168 0.0236 0.0135
       0.0067 0.0101 0.0067 0.0067 0 0.0101 0 0.0034 0
                                                                           0
                                                                                     0
        0.0067 \ \ 0.0168 \ \ 0.0438 \ \ 0.0303 \ \ 0.0505 \ \ 0.0673 \ \ 0.0572 \ \ 0.0236 \ \ 0.0438 \ \ 0.037 \ \ \ 0.0236
        0.0135 0.0101 0.0101 0.0101 0.0101 0.0034 0.0034 0.0067 0.0034 0.0034 0.0101
       0.0034
1990
        1
                                               -1
                                                       -1
                                                              21
                                                                      0.008 0.12
                                       0.048
                                              0.032
                                                      0.008
                                                             0.016 0.008 0.008
        0 032
               0.064
                      0.016 0.032
                                                                                      0 024
                                       0
                                               0
        0
               0
                       0
                               0
                                                       0
                                                              0
                                                                      0
                                                                              0
                                                                                      0.016
        0.032
                      0.064
                               0.048
                                       0.032
                                              0.04
                                                       0.024
                                                                              0.008
               0.112
                                                              Ω
                                                                      Ω
                                                                                      0.008
        0.008
               0
                       Ω
                               Ω
                                       Ω
                                               Ω
                                                       0.008 0
                                                                      Ω
                                                                              Ω
       0.008
                                                                      0.0021 0.0042 0.04
1991
       1
                               0
                                       1
                                               -1
                                                      -1
                                                             88
        0.0989 0.0926 0.0905 0.0463 0.0379 0.0358 0.0232 0.0211 0.0021 0.0042 0.0105
        0.0021 \ 0 \qquad 0 \qquad 0.0021 \ 0 \qquad 0 \qquad 0.0021 \ 0 \qquad 0 \qquad 0.0021 \ 0 \qquad 0.0021
        0.0211 \ \ 0.0337 \ \ 0.0674 \ \ 0.0947 \ \ 0.0905 \ \ 0.0295 \ \ 0.0337 \ \ 0.0147 \ \ 0.0189 \ \ 0.0168 \ \ 0.0168
       0.0105 0.0063 0.0021 0
                                      0.0021 0.0021 0.0042 0.0042 0.0042 0
                                                                                      0.0021
1992
                                                             49
                                                                     0.0037 0.0147 0.0586
                       3
                              0
                                     1
                                              -1
                                                      -1
       0.0476 0.0989 0.1209 0.0879 0.0293 0.022 0.0256 0.0293 0.0037 0.0073 0.0037
                                              0.0037 0 0.0037 0
        0.0037 0 0
                              Ο
                                     0
```

	0.0073 0.0073 0	0.0476 0.0147	0.0623 0.0073	0.0586 0.0073	0.0513	0.0366 0.011	0.033 0.0037	0.0183 0.0037	0.0147 0	0.033 0.0073	0.0073
1993	1 0.1053 0 0.0031 0.0031 0.0031	3 0.096 0.0062 0.0217 0.0031	0.0031 0.0836	0 0.1053 0 0.065	1 0.0402 0 0.1022 0.0031	-1 0.0341 0 0.0433 0.0031	-1 0.0155 0 0.0433 0	0 0.0279	0.0031 0 0.0186	0.0062 0.0031 0 0.0031	0.0341 0 0 0.0217 0.0031
1994	1 0.084 0 0.008 0.012	3 0.16 0 0.064	3 0.052 0 0.048 0.008	0 0.056 0 0.076 0.008	1 0.016 0 0.08	-1 0.016 0 0.048	-1 0.004 0 0.048	44 0.008 0 0.028	0.004 0.008 0 0.02	0.04 0 0 0.008	0.084 0 0.008 0.004
1995 #	1 0.1161 0 0.0179 0.0045	3 0.0938 0.0045 0.0982	3 0.1205 0 0.1116 0.0134	0 0.0536 0 0.067	1 0.0223 0 0.0446 0.0045		-1 0.0045 0 0.0402	40 0.0089 0 0.0268	0 0.0045 0 0 0.0045	0 0 0.0134	0.0625 0 0 0.0045
#											

# Mean Size at Age

#_Year		Type		Partitio		Age-Err		3		5	6	
	7	8	9			12	13	14	15	16	17	
	18	19	20	21	22	23	24	25	30	3	4	
	5	6	7	8	9	10	11	12	13	14	15	
	16	17	18	19	20	21	22	23	24	25	30	
#1986	1	2	3	0	1	1	27.5009	5098	31.2838	9375		
	34.4412	3171	37.0764	3134	39.2758	402	41.1115	2629	42.6436	3953		
	43.9223		44.9896	5626	45.8804	3165	46.6238	9688	47.2444	1306		
	47.7623		48.1945			48.55533566 48.85644408			49.1077	5731		
	49.3175		49.4925			.63868979 49.76064073			49.8624			
	49.9473		50.0182		28.1226401 31.45069993				34.20130247			
	36.4746	4329	38.3535			39.90641319 41.18985			42.2505			
	43.1272	9075	43.8518	6823	44.4507	2304	44.9456	6953	45.35473703			
	45.6928		45.9722		46.2031	967	46.3940	6852	46.5518	2185		
	46.6822		46.7899		46.8790		46.9526		47.0134			
	47.0637	4764										
#	0	0	1	9	17	27	28	27	13	19	8	
	8	5	2	5	5	2	0	0	0	0	0	
	0	0	0	0	3	7	26	32	18	16	8	
	10	7	4	7	1	0	1	1	1	1	1	
	0	0	1	0								
#1986	1	1	3	0	1	1	28	31.5	37	41.2	42.1	
	44.6	46.9		49.6	48.4	48.9	48.1	50.6	52	52.2	50	
	49	51	51	51	51	51	51	51	28	31.5	42	
	40.9	41.4	42.2	43.6	45.1	45.1	47.1	46.6	47	46.7	46	
	47.3	46	49	52	46	50	47.3	47.3	49	47.3		
#	0	0	1	9	17	27	28	27	13	19	8	
	8	5	2	5	5	2	0	0	0	0	0	
	0	0	0	0	3	7	26	32	18	16	8	
	10	7	4	7	1	0	1	1	1	1	1	
	0	0	1	0								
#1987	1	1	3	0	1	1	28	31.5	44	41.3	47.1	
	46.5	47	48	47.8	50.4	50.4	50.4	51.2	51.6	53.8	56	
	51	56	54.5	53	58	60	50	50	28	31.5	36	

	41.3	43.4	43.5	44.3	45.4	44.2	45.8	46.1	46.9	47.1	49
	48.8	48	43.5	51	51	47.3	47.3	53	52	53	49
#	0	0	1	3	10	29	34	23	27	26	20
	13 0	14 0	5 0	5 0	1 0	0 4	4 8	2 28	2 35	1 20	1 15
	22	13	7	8	6	4	0	2	1	1	0
	0	1	3	1							
#1984	1	2	3	0	1	1	28	32.3	34.4	36.5	38.9
	40.6 55.9	42.8 50	43.2 41	45.9 47.1	46 50	48.4 50	48.7 50	49.9 50	48.5 28	52.4 30.3	51.3 36.1
	37.1	39.1	40.2	40.7	42.4	42.6	43.4	44.4	45.8	47.3	46.7
	46.5	45.2	47.7	44.9	51.6	51.1	45.3	49	49.5	50.3	
#	0	6	26	62	75 -	44	42	61	34	20	17
	8 0	9 0	9 0	5 2	5 2	2 21	0 24	1 23	2 29	0 34	0 24
	12	11	10	14	9	12	2	3	2	6	3
	6	4	5	4							
#1993	1	2	3	0	1	1	28	32.6	33.9	36.5	39
	40.6 51.8	41.7 55	44.3 52	44.5 50	47 50	49.7 50	49.8 54	47.2 50	46.8 28	50 35.4	47 33.1
	36.4	38	38.6	40.8	41.5	43.6	44.1	45.3	44.5	46.2	45.2
	45.9	50.5	45.3	48.3	46.7	47.3	47.3	48	48	54	
#	0	7	34	69	66	85	85	48	35	11	15
	10 1	12 0	6 0	2 7	2 20	4 57	1 46	1 69	0 50	0 30	0 17
	16	11	2	5	6	10	2	3	3	3	0
	0	1	3	1							
2001	1 39.7	2 40.2	3 41.1	0 42.6	1 43.8	1 43.8	28 45.3	34.2 45.6	35.8 47.1	36.8 44.1	38.2 48.4
	46	42.5	43.3	44	50	50	50	50	28	32.3	33.5
	36.4	37.5	38.5	39.3	40.3	41.3	42.4	42.3	42.1	41.8	42.9
	43.9	44	44.7	45.5	42.3	44	47.3	46.8	44.5	51	- 4
	0 28	6 24	25 13	71 12	190 8	162 4	98 4	79 3	96 1	84 0	64 0
	0	0	0	10	21	74	164	120	82	72	56
	54	36	21	18	14	8	7	9	8	3	2
2002	4	4 2	6	5	1	1	25	21	25 5	27 6	20 1
2002	1 40.8	41.6	3 42.6	0 43.8	1 44.6	1 45.1	25 45.4	31 46.8	35.5 46.3	37.6 44.4	39.1 46.9
	49.3	49.3	47	50	45	50	48	50	27	30	34.3
	36.4	38.2	39.4	39.5	41.1	41.1	41.6	42.5	42.3	43.4	43.1
	44.4 1	44.7 4	44 27	45.8 56	44 144	43.7 222	41 141	45.7 113	46.3 94	48.5 77	71
	39	23	20	9	8	3	6	1	0	2	0
	1	0	2	3	33	73	123	137	96	82	43
	41	34	28	22	15	12	10	3	4	3	3
#1987	2 1	3 3	6 3	4 0	1	1	28	35.7	35.3	36.6	38.8
111101	41.2	43.2	44.4	45	46.6	47.7	48.6	48.9	51.6		50.8
	50.5	52	50	57	51	50	50	50	28	30.7	34.4
	37.9 46.9	38.6 49.4	40.3 50.3	41.5 47	41.6 47.6	44 49	45 52.5	45.5 47.3	45.9 49.43	44.6 51	48.3
#	0	3	10	34	32	44	52.5	27	26	29	21
	28	15	16	8	5	2	3	0	1	2	0
	0	0	0	3	15	23	45	42	35	31	21
	24 2	32 0	21 6	17 3	4	10	13	4	5	5	2
#1991	1	3	3	0	1	1	29	32	34.1	36.1	39
	41.4	42.7	44.9	46.8	47.5	50.5	52	50.8	49.3	54	50
	50	50	47	50	50	50	51	50	34	30.8	33.8
	35.7 47.5	38.4 48	40.9 51	41.8 49.2	44.5 48.3	45.6 49.3	46.5 49.5	45.6 50	45.8 50.7	45.2 47.3	49.6
#	1	2	19	49	49	48	27	21	24	19	13
	7	5	7	3	0	0	0	1	0	0	0
	1	0	1	10	16 7	33	50	51 1	19 5	27	12
	17 4	14 1	9 3	6 3	7	2	2	1	5	4	6
#		nmental		-							

#