

In order to develop an intuition for the mathematical tools we will put to use in the all-orders calculation of groomed heavy hemisphere mass, let us first compute the first-order (or rather, first *nontrivial* order) distribution of the observable. A simple electron-positron production with the production of two quark jets, $e^+e^- \rightarrow q\bar{q}$, would produce a constant distribution: the mass of the heavy hemisphere would simply be half of the final mass. The lowest-order nontrivial distribution corresponds to an $e^+e^- \rightarrow q\bar{q}g$ event, in which an electron and positron annihilate to produce a quark-antiquark pair, off of which a single gluon is emitted. The location of the gluon in phase space sets the heavy hemisphere mass. This process is depicted in a Feynman diagram in Fig. 1.




Figure 1: Feynman diagram of an $e^+e^- \rightarrow q\bar{q}g$ event.

We will work in the limit that the heavy hemisphere mass ρ is approximately equal to the mMDT cutoff z_{cut} , which is itself small: $\rho \sim z_{\text{cut}} \ll 1$. We will utilize the strategy of regions [1] to compute the distribution. This is necessary because the distribution is essentially singular in the limit of interest **[TODO: is this accurate?]**. The method of regions entails computing the distribution in all singular regions of phase space, then summing the results at the end to generate the full distribution.¹ There are three singular regions of phase space:

1. The **soft region**, in which the gluon is emitted with low energy.
2. The **collinear region**, in which the gluon is emitted collinear to the quark or antiquark.
3. The **soft-collinear region**, in which the gluon is both low-energy and collinear to the quark or antiquark.

If we label these cross sections σ_{soft} , $\sigma_{\text{collinear}}$, and $\sigma_{\text{soft-collinear}}$, respectively, then the full cross distribution will be

$$\frac{d\sigma}{d\rho} = \frac{d\sigma_{\text{soft}}}{d\rho} + \frac{d\sigma_{\text{collinear}}}{d\rho} + \frac{d\sigma_{\text{soft-collinear}}}{d\rho}. \quad (1)$$

Let us calculate these components now.

¹The fact that this works is rather magical.

0.1 Soft gluon

0.1.1 Setup

For now, we assume a soft (i.e., low-energy) gluon. Recall that the (normalized) heavy hemisphere mass is defined to be

$$\rho = \left(\frac{m_h}{E_h} \right)^2 \quad (2)$$

with m_h the mass of the more massive hemisphere and E_h its energy.

We first need to sort out the kinematics of the event. Let us shift our reference frame so that the quark has momentum

$$p_1^\mu = \frac{Q_q}{2}(1, 0, 0, 1) \quad (3)$$

and the antiquark has momentum

$$p_2^\mu = \frac{Q_q}{2}(1, 0, 0, -1), \quad (4)$$

and let the gluon have momentum k^μ . The soft-gluon limit means that the energy of the gluon is $k^0 \ll 1$. In this case, the quarks carry most of the energy, so we can approximate $Q_q \approx Q$, the total energy of the event. Furthermore, let us assume that the gluon is emitted in the hemisphere containing the quark (the problem is symmetric under quark-antiquark exchange, so we will simply multiply the cross section by 2 at the end to account for this assumption). Then the momentum of the heavy hemisphere is

$$p_h = p_1 + k, \quad (5)$$

so the heavy hemisphere has mass

$$\begin{aligned} m_h^2 &= p_h^2 = (p_{1,0} + k_0)^2 - (k_1)^2 - (k_2)^2 - (p_{1,3} + k_3)^2 \\ &= 2p_{1,0}k_0 - 2p_{1,3}k_3 + p_1^2 + k^2 \\ &= Q(k_0 - k_3). \end{aligned} \quad (6)$$

The last line follows because we assume every particle to be massless, such that $p_1^2 = k^2 = 0$.² Now let us introduce the **light-cone coordinates**

$$k^+ \equiv k^0 - k^3 \qquad k^- \equiv k^0 + k^3, \quad (7)$$

this can be written simply as

$$m_h^2 = Qk^+. \quad (8)$$

²This is a reasonably accurate assumption for the energies accessible by colliders, and makes our calculations much easier. The gluon is actually massless regardless of the theoretical assumptions made.

The energy of the heavy hemisphere is

$$E_h = p_{1,0} + k_0 = \frac{Q}{2} + k_0 \approx \frac{Q}{2}, \quad (9)$$

since, for a soft gluon, $k_0 \ll Q/2$. The heavy hemisphere mass is therefore

$$\rho = \frac{Qk^+}{Q^2/4} = \frac{4k^+}{Q}. \quad (10)$$

This means that we will need to insert the measurement function

$$\delta\left(\rho - \frac{4k^+}{Q}\right) \quad (11)$$

into Fermi's Golden Rule.

With the kinematics under our belt, let us think about the effects of an mMDT groomer. For the simple case of only 3 particles, the groomer only keeps pairs of particles i and j for which [2, 3]

$$\frac{\min[E_i, E_j]}{E_i + E_j} > z_{\text{cut}}. \quad (12)$$

This must be true for the quark and gluon; since the gluon has lower energy than the quark, this necessitates that

$$\frac{k_0}{E_h} = \frac{(k^+ + k^-)/2}{Q/2} > z_{\text{cut}}, \quad (13)$$

or

$$k^+ + k^- > Q z_{\text{cut}}. \quad (14)$$

Moreover, since we are assuming that the gluon shares a hemisphere with the quark, we must also have $k_3 > 0$, which requires

$$k^- - k^+ > 0. \quad (15)$$

Equations 14 and 15 generate the phase space constraints

$$\Theta(k^+ + k^- - Q z_{\text{cut}})\Theta(k^- - k^+). \quad (16)$$

When we insert these into Fermi's Golden Rule, the differential cross section takes the form

$$\frac{d\sigma_{\text{soft}}}{d\rho} = 2 \int d\text{LIPS} |\mathcal{M}|^2 \delta\left(\rho - \frac{4k^+}{Q}\right) \Theta(k^+ + k^- - Q z_{\text{cut}}) \Theta(k^- - k^+). \quad (17)$$

Here, $d\text{LIPS}$ is a differential element of Lorentz-invariant phase space, and \mathcal{M} is the matrix element governing the $e^+e^- \rightarrow q\bar{q}g$ process.

Assuming a soft gluon, the matrix element is well-known in the literature (see Eqs. 87 and 88 of [4]):

$$|\mathcal{M}|^2 = 4\pi\alpha_s\sigma_0 C_F \mu^{2\epsilon} \frac{p_1 \cdot p_2}{(p_1 \cdot k)(p_2 \cdot k)}. \quad (18)$$

Here, α_s is the strong coupling constant³, σ_0 is the cross section for $e^+e^- \rightarrow q\bar{q}$, C_F is the quadratic Casimir of the fundamental representation of color (taken to be $C_F = 4/3$ for our purposes [5]), and μ is a mass scale introduced to ensure that the differential cross section will remain dimensionless in $d = 4 - 2\epsilon$ dimensions. Inserting the values of p_1 , p_2 , and k , we have

$$|\mathcal{M}|^2 = 4\pi\alpha_s\sigma_0 C_F \mu^{2\epsilon} \frac{2}{k^+ k^-}. \quad (19)$$

Now we must unpack the Lorentz-invariant phase space element $d\text{LIPS}$. Working in d dimensions, the standard form is

$$d\text{LIPS} = \frac{d^d k}{(2\pi)^{d-1}} \delta(k^2) \Theta(k_0), \quad (20)$$

where the Dirac delta and Heaviside functions ensure that the gluon is on-shell (i.e., real) with positive energy. If $\epsilon = 0$, we would have

$$d^d k = d^4 k = dk_0 dk_1 dk_2 dk_3. \quad (21)$$

When we transform to light-cone coordinates with $(k_0, k_3) \rightarrow (k^+, k^-)$, the Jacobian of the transformation is

$$\frac{\partial(k_0, k_3)}{\partial(k^+, k^-)} = \begin{pmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{pmatrix}, \quad (22)$$

so

$$dk_0 dk_3 = \left| \det \frac{\partial(k_0, k_3)}{\partial(k^+, k^-)} \right| dk^+ dk^- = \frac{1}{2} dk^+ dk^-. \quad (23)$$

Now let $k_\perp = (k_1, k_2)$ be the transverse components of the gluon momentum. For $\epsilon \neq 0$, we imagine that these transverse components are the ones which bleed into the modified dimensions. Therefore, after noticing that

$$\delta(k^2) = \delta(k_0^2 - k_3^2 - k_\perp^2) = \delta(k^+ k^- - k_\perp^2), \quad (24)$$

we have

$$d\text{LIPS} = \frac{dk^+ dk^- d^{d-2} k_\perp}{2(2\pi)^{d-1}} \delta(k^+ k^- - k_\perp^2) \Theta(k^+ + k^-). \quad (25)$$

Now it is convenient to transfer the $2 - 2\epsilon$ dimensions of k_\perp into spherical coordinates, so that

$$d^{d-2} k_\perp = k_\perp^{d-3} dk_\perp d\Omega_{d-2} \quad (26)$$

³Which is not really a constant

with Ω_{d-2} the solid angle of the $(d-2)$ -dimensional unit sphere. Since none of the terms in the cross section of Eq. 17 or matrix element of Eq. 19 have angular dependence, we can go ahead and integrate the solid angle:

$$\int d\Omega_{d-2} = \frac{2\pi^{(d-2)/2}}{\Gamma(\frac{d-2}{2})} \quad (27)$$

with $\Gamma(x)$ the gamma function; this identity comes from Eq. B.28 of [6]. Therefore,

$$d\text{LIPS} = \frac{2\pi^{(d-2)/2}}{\Gamma(\frac{d-2}{2})} \frac{dk^+ dk^- dk_\perp}{2(2\pi)^{d-1}} k_\perp^{d-3} \delta(k^+ k^- - k_\perp^2) \Theta(k^+ + k^-). \quad (28)$$

As a final step, we can resolve this delta function:

$$\delta(k^+ k^- - k_\perp^2) = \frac{1}{2\sqrt{k^+ k^-}} \delta\left(k_\perp - \sqrt{k^+ k^-}\right). \quad (29)$$

Then integrating over k_\perp yields

$$\int dk_\perp \frac{k_\perp^{d-3}}{2\sqrt{k^+ k^-}} \delta\left(k_\perp - \sqrt{k^+ k^-}\right) = \frac{1}{2} (k^+ k^-)^{(d-4)/2}. \quad (30)$$

Putting everything together, inserting $d = 4 - 2\epsilon$, and simplifying, we are left with

$$d\text{LIPS} = \frac{(4\pi)^\epsilon}{\Gamma(1-\epsilon)(4\pi)^2} \frac{dk^+ dk^-}{(k^+ k^-)^\epsilon} \Theta(k^+ + k^-). \quad (31)$$

Notice that a factor of $(k^+ k^-)^\epsilon$ has been introduced — this is what will help us capture divergences as we work and cancel them at the end. Finally, we will work using a convention known as **modified minimal subtraction**, under which we will throw away factors of $(4\pi)^\epsilon$ and set the Euler-Mascheroni constant to be $\gamma_E \rightarrow 0$ when it appears (this does not affect the final result, but will make calculations slightly less unwieldy). Under this scheme, we have

$$d\text{LIPS} = \frac{1}{\Gamma(1-\epsilon)(4\pi)^2} \frac{dk^+ dk^-}{(k^+ k^-)^\epsilon} \Theta(k^+ + k^-). \quad (32)$$

as our final phase space element.

Combining Eqs. 17, 19, and 32 then yields the full cross section:

$$\boxed{\frac{1}{4\pi\alpha_s\sigma_0 C_F} \frac{d\sigma_{\text{soft}}}{d\rho} = \frac{4\mu^{2\epsilon}}{\Gamma(1-\epsilon)(4\pi)^2} \int \frac{dk^+ dk^-}{(k^+ k^-)^{1+\epsilon}} \delta\left(\rho - \frac{4k^+}{Q}\right) \Theta(k^+ + k^-) \times \Theta(k^+ + k^- - Q z_{\text{cut}}) \Theta(k^- - k^+).} \quad (33)$$

0.1.2 Calculation

The integral of Eq. 35 is relatively straightforward to evaluate. The first step is to resolve the Dirac delta:

$$\delta\left(\rho - \frac{4k^+}{Q}\right) = \frac{Q}{4} \delta\left(k^+ - \frac{Q\rho}{4}\right). \quad (34)$$

The integrating over k^+ yields

$$\frac{1}{4\pi\alpha_s\sigma_0 C_F} \frac{d\sigma_{\text{soft}}}{d\rho} = \frac{4\mu^{2\epsilon}}{\Gamma(1-\epsilon)(4\pi)^2} \frac{1}{\rho} \left(\frac{4}{Q\rho}\right)^\epsilon \int \frac{dk^-}{(k^-)^{1+\epsilon}} \Theta\left(\frac{Q\rho}{4} + k^-\right) \times \Theta\left(\frac{Q\rho}{4} + k^- - Q z_{\text{cut}}\right) \Theta\left(k^- - \frac{Q\rho}{4}\right). \quad (35)$$

Now, the integrand is only non-zero when

$$k^- > -\frac{Q\rho}{4} \quad k^- > Q\left(z_{\text{cut}} - \frac{\rho}{4}\right) \quad k^- > \frac{Q\rho}{4}. \quad (36)$$

If the second and third requirements are satisfied, then so is the first, so we can ignore it. To deal with the others, notice that each is stricter for different values of ρ : if $\rho < 2z_{\text{cut}}$, then

$$Q\left(z_{\text{cut}} - \frac{\rho}{4}\right) > \frac{Q\rho}{4}, \quad (37)$$

and the opposite is true if $\rho > 2z_{\text{cut}}$. We can therefore break the integral into two pieces:

$$\begin{aligned} \int \frac{dk^-}{(k^-)^{1+\epsilon}} \Theta\left(\frac{Q\rho}{4} + k^-\right) \Theta\left(\frac{Q\rho}{4} + k^- - Q z_{\text{cut}}\right) \Theta\left(k^- - \frac{Q\rho}{4}\right) \\ = \Theta(\rho - 2z_{\text{cut}}) \int_{Q\rho/4}^{\infty} \frac{dk^-}{(k^-)^{1+\epsilon}} + \Theta(2z_{\text{cut}} - \rho) \int_{Q(z-\rho/4)}^{\infty} \frac{dk^-}{(k^-)^{1+\epsilon}}. \end{aligned} \quad (38)$$

If we take $\epsilon > 0$, then these integrals yield a finite result:

$$\begin{aligned} \frac{1}{4\pi\alpha_s\sigma_0 C_F} \frac{d\sigma_{\text{soft}}}{d\rho} = \frac{4\mu^{2\epsilon}}{\Gamma(1-\epsilon)(4\pi)^2} \frac{1}{\rho} \left(\frac{4}{Q\rho}\right)^\epsilon \frac{1}{\epsilon} \left[\Theta(\rho - 2z_{\text{cut}}) \left(\frac{Q\rho}{4}\right)^{-\epsilon} \right. \\ \left. + \Theta(2z_{\text{cut}} - \rho) \left(Q\left(z_{\text{cut}} - \frac{\rho}{4}\right)\right)^{-\epsilon} \right]. \end{aligned} \quad (39)$$

Notice how dimensional regularization helped us achieve this calculation: without the regulating $(k^-)^\epsilon$, the integrals would have diverged without an upper bound on the value of k^- (which, physically, has no upper bound). Our result is still manifestly divergent if we send $\epsilon \rightarrow 0$, but at least we can *see* the divergence. This is the power of the technique.

We can pull out the divergence even more cleanly if we perform a Laurent expansion⁴ in ϵ . We will send $\epsilon \rightarrow 0$ at the end anyway, so we only care about terms through order $\mathcal{O}(\epsilon^0)$; anything below this order generates divergences, and anything above this order will vanish. Performing the expansion yields

$$\begin{aligned} \frac{1}{4\pi\alpha_s\sigma_0 C_F} \frac{d\sigma_{\text{soft}}}{d\rho} = \frac{1}{4\pi^2\rho} \left[\frac{1}{\epsilon} + \Theta(\rho - 2z_{\text{cut}}) 2 \log\left(\frac{4\mu}{Q\rho}\right) \right. \\ \left. + \Theta(2z_{\text{cut}} - \rho) \left[\log\left(\frac{4\mu^2}{Q\rho}\right) - \log\left(Q z_{\text{cut}} - \frac{Q\rho}{4}\right) \right] \right] + \mathcal{O}(\epsilon). \end{aligned}$$

(40)

⁴Like a Taylor expansion, but possibly including negative exponents

From this expansion, we see that the soft contribution to the cross section diverges as

$$\lim_{\epsilon \rightarrow 0} \frac{d\sigma_{\text{soft}}}{d\rho} \sim \frac{\alpha_s \sigma_0 C_F}{\pi \rho \epsilon}. \quad (41)$$

Stop reading and appreciate this for a minute — it is remarkable! By pushing our calculation out of the standard 4 dimensions, we are able to learn about structure that was inaccessible to us in our 4-dimensional perspective.

This technique, moreover, is not only beautiful from a mathematical point of view; it will be extremely useful to have analytically extracted the divergences in this way. At the end of the calculation, we will find that they all cancel each other out. It is rather magical.

0.2 Collinear gluon

Now that we have computed the contribution from a soft gluon, let us move on to the next singular region of phase space: a gluon collinear to the quark or antiquark.

0.2.1 Setup

For this calculation, we will use a different system of coordinates: the gluon's hemisphere energy fraction and angle from the quark. To derive these coordinates, we first define the phase-space coordinates

$$x_i = \frac{2p_i \cdot Q}{Q^2} \quad (42)$$

where $i = 1, 2, 3$ ranges over the three particles of the event and $Q = p_1 + p_2 + p_3$ is the total four-momentum of the event. Let x_1 be the energy fraction of the quark, x_2 be the energy fraction of the antiquark, and x_3 be the energy fraction of the gluon. Also let $k = p_3$ be the momentum of the gluon. Notice that

$$x_1 + x_2 + x_3 = \frac{2(p_1 + p_2 + p_3) \cdot Q}{Q^2} = 2. \quad (43)$$

In the collinear limit, each hemisphere carries half the momentum and energy (in order to conserve the net-zero initial momentum of the collision). Assume now that the gluon is emitted in the same hemisphere as the quark (we will again multiply the result by a factor of 2 to compensate). Then, in the collinear limit, we have

$$x_1 + x_3 \rightarrow 1. \quad (44)$$

Now we will introduce the gluon's energy fraction

$$z \equiv \frac{x_3}{x_1 + x_3} \approx x_3, \quad (45)$$

where the final step holds in the collinear limit. The quark's hemisphere energy fraction is

$$1 - z = \frac{x_1}{x_1 + x_3} \approx x_1. \quad (46)$$

This is equivalent to the assumption that the quark four-momentum p_1 and the gluon four-momentum k are collinear along some vector \bar{p}_1 :

$$k = z\bar{p}_1 \quad p_1 = (1 - z)\bar{p}_1. \quad (47)$$

Now let θ be the angle between the quark and the gluon. Notice that

$$1 - x_2 = \frac{Q^2 - 2p_2 \cdot Q}{Q^2} = \frac{2p_1 \cdot k}{Q^2} = \frac{x_1 x_3}{2}(1 - \cos \theta). \quad (48)$$

In the collinear limit $\theta \ll 1$, we have $\cos \theta \approx 1 - \theta^2/2$, so this means that

$$\frac{2p_1 \cdot k}{Q^2} = \frac{x_1 x_3}{4}\theta^2 = \frac{z(1 - z)}{4}\theta^2. \quad (49)$$

Then the heavy hemisphere mass is

$$m_h^2 = (p_1 + k)^2 = 2p_1 \cdot k = \frac{z(1 - z)}{4}\theta^2 Q^2, \quad (50)$$

where again we have $p_1^2 = k^2 = 0$. Since the hemisphere energy is half the total energy, $E_h = Q/2$, the observable we are looking for is then

$$\rho = \frac{m_h^2}{E_h^2} = z(1 - z)\theta^2. \quad (51)$$

The measurement function in Fermi's Golden Rule will then be

$$\delta(\rho - z(1 - z)\theta^2) = \frac{1}{z(1 - z)}\delta\left(\theta^2 - \frac{\rho}{z(1 - z)}\right). \quad (52)$$

The quark and gluon only pass the mMDT groomer if [3]

$$\frac{\min[E_1, E_3]}{E_1 + E_3} > z_{\text{cut}}. \quad (53)$$

This means that we require

$$\min[x_1, x_3] = \min[z, 1 - z] > z_{\text{cut}}. \quad (54)$$

Thus, the grooming constraint on the cross section takes the form

$$\Theta(\min[z, 1 - z] - z_{\text{cut}}). \quad (55)$$

In phase space coordinates, the matrix element for $e^+e^- \rightarrow q\bar{q}g$ is [7]

$$|\mathcal{M}|^2 = 4\pi\alpha_s\sigma_0 C_F \frac{x_1^2 + x_2^2}{(1 - x_1)(1 - x_2)}. \quad (56)$$

After performing the the change of variables discussed above and introducing the appropriate Jacobian factor, this reduces to

$$|\mathcal{M}|^2 = 4\pi\alpha_s\sigma_0 C_F \frac{1 + (1 - z)^2}{z\theta^2}. \quad (57)$$

Finally, we must sort out the phase space measure. In $d = 4 - 2\epsilon$ dimensions, the phase space integral with the matrix element is **[TODO: where does this come from? obtained from Andrew's notes but need to either derive or cite]**

$$\frac{4\pi}{\alpha_s C_F} \frac{1}{\sigma_0} \sigma_{\text{collinear}} = \frac{1}{\sqrt{\pi} \Gamma(\frac{1}{2} - \epsilon)} \left(\frac{2\mu}{Q} \right)^{2\epsilon} \int_0^1 dz \int_0^\infty d\theta^2 \int_0^{2\pi} d\phi \sin^{-2\epsilon} \phi \times (\theta^2)^{-1-\epsilon} z^{-2\epsilon} (1-z)^{-2\epsilon} \left(\frac{1 + (1-z)^2}{z} - \epsilon z \right). \quad (58)$$

When we introduce the measurement and grooming terms of Eqs. 52 and 55, we find that the full differential cross section is

$$\boxed{\begin{aligned} \frac{4\pi}{\alpha_s C_F} \frac{1}{\sigma_0} \frac{d\sigma_{\text{collinear}}}{d\rho} &= \frac{1}{\sqrt{\pi} \Gamma(\frac{1}{2} - \epsilon)} \left(\frac{2\mu}{Q} \right)^{2\epsilon} \int_0^1 dz \int_0^\infty d\theta^2 \int_0^{2\pi} d\phi \sin^{-2\epsilon} \phi (\theta^2)^{-1-\epsilon} \\ &\quad \times z^{-2\epsilon} (1-z)^{-2\epsilon} \left(\frac{1 + (1-z)^2}{z} - \epsilon z \right) \\ &\quad \times \frac{1}{z(1-z)} \delta\left(\theta^2 - \frac{\rho}{z(1-z)}\right) \\ &\quad \times \Theta(\min[z, 1-z] - z_{\text{cut}}). \end{aligned}} \quad (59)$$

0.2.2 Calculation

We can immediately perform the integrals in ϕ ,

$$\int_0^{2\pi} d\phi \sin^{-2\epsilon} \phi = \frac{\sqrt{\pi} \Gamma(\frac{1}{2} - \epsilon)}{\Gamma(1 - \epsilon)}, \quad (60)$$

and in θ^2 to find

$$\frac{4\pi}{\alpha_s C_F} \frac{1}{\sigma_0} \frac{d\sigma_{\text{collinear}}}{d\rho} = \frac{1}{\Gamma(1 - \epsilon)} \left(\frac{2\mu}{Q} \right)^{2\epsilon} \frac{1}{\rho^{1+\epsilon}} \int_0^1 dz \frac{1}{z^\epsilon (1-z)^\epsilon} \left(\frac{1 + (1-z)^2}{z} - \epsilon z \right) \times \Theta(\min[z, 1-z] - z_{\text{cut}}). \quad (61)$$

The Heaviside function is satisfied by ensuring that

$$z_{\text{cut}} < z < 1 - z_{\text{cut}}, \quad (62)$$

so

$$\int_0^1 dz \Theta(\min[z, 1-z] - z_{\text{cut}}) = \int_{z_{\text{cut}}}^{1-z_{\text{cut}}} dz. \quad (63)$$

The cross section becomes

$$\frac{4\pi}{\alpha_s C_F} \frac{1}{\sigma_0} \frac{d\sigma_{\text{collinear}}}{d\rho} = \frac{1}{\Gamma(1 - \epsilon)} \left(\frac{2\mu}{Q} \right)^{2\epsilon} \frac{1}{\rho^{1+\epsilon}} \int_{z_{\text{cut}}}^{1-z_{\text{cut}}} dz \frac{1}{z^\epsilon (1-z)^\epsilon} \left(\frac{1 + (1-z)^2}{z} - \epsilon z \right). \quad (64)$$

This integral does not diverge in 4 dimensions, so we can simply set $\epsilon = 0$.⁵ Thus,

$$\frac{4\pi}{\alpha_s C_F \sigma_0} \frac{d\sigma_{\text{collinear}}}{d\rho} = \frac{1}{\rho} \int_{z_{\text{cut}}}^{1-z_{\text{cut}}} dz \frac{1 + (1-z)^2}{z} + \mathcal{O}(\epsilon). \quad (65)$$

This comes out to

$$\boxed{\frac{4\pi}{\alpha_s C_F \sigma_0} \frac{d\sigma_{\text{collinear}}}{d\rho} = \frac{1}{\rho} \left[-\frac{3}{2} + 3z_{\text{cut}} + 2 \log \left(\frac{1 - z_{\text{cut}}}{z_{\text{cut}}} \right) \right] + \mathcal{O}(\epsilon).} \quad (66)$$

0.3 Soft-collinear gluon

0.4 Putting it all together

⁵This is equivalent to computing the $\mathcal{O}(\epsilon^0)$ term in the Taylor expansion.

References

- [1] T. Becher, A. Broggio and A. Ferroglia, *Introduction to Soft-Collinear Effective Theory*, vol. 896 of *Lecture Notes in Physics*, Springer International Publishing, Cham (2015), 10.1007/978-3-319-14848-9.
- [2] M. Dasgupta, A. Fregoso, S. Marzani and G.P. Salam, *Towards an understanding of jet substructure*, *J. High Energ. Phys.* **2013** (2013) 29.
- [3] A. Kardos, A.J. Larkoski and Z. Trócsányi, *Two- and three-loop data for the groomed jet mass*, *Phys. Rev. D* **101** (2020) 114034.
- [4] S. Catani and M. Grazzini, *Infrared factorization of tree-level QCD amplitudes at the next-to-next-to-leading order and beyond*, *Nuclear Physics B* **570** (2000) 287 [[hep-ph/9908523](#)].
- [5] Particle Data Group, *Review of Particle Physics, Progress of Theoretical and Experimental Physics* **2020** (2020) .
- [6] M.D. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge University Press, New York (2014).
- [7] A.J. Larkoski, *Improving the Understanding of Jet Grooming in Perturbation Theory*, *arXiv:2006.14680 [hep-ex, physics:hep-ph]* (2020) [[2006.14680](#)].