

The first step on the path to an all-orders calculation is to derive a factorization formula for the heavy hemisphere mass cross section. The basic process for doing so is laid out in technical detail in Ref. [1], and an example of a similar flavor to our calculation is provided by Frye et al. in Ref. [2].¹ There are two primary steps in developing a factorization formula:

1. **Power counting:** this involves determining the possible radiative modes of an event and their dominant momentum scales. The term ‘power counting’ refers to the fact that for some momentum scale λ , different radiative modes have momenta that scale as different powers of λ .
2. **Factorization and refactorization:** Once the different radiative modes and energy scales are identified, we can use the framework of SCET to split the cross section into a convolution of terms describing different radiative modes. These terms themselves must then be split (refactored) into convolutions of terms, each of which depends, to leading order, only on a single energy scale.

0.1 Setup

Recall that the hemisphere mass is defined to be

$$\rho = \frac{1}{E_J^2} \sum_{i < j} 2p_i \cdot p_j \quad (1)$$

with E_J the jet energy and the sum ranging over all pairs of particles in the jet. Expanding out the dot product, we have

$$\rho = \frac{2}{E_J^2} \sum_{i < j} (E_i E_j - \mathbf{p}_i \cdot \mathbf{p}_j) = \frac{2}{E_J^2} \sum_{i < j} E_i E_j (1 - \cos \theta_{ij}) = \sum_{i < j} 2z_i z_j (1 - \cos \theta_{ij}). \quad (2)$$

Here, z_i and z_j are the relative energy fractions of each particle and θ_{ij} is the angle between particles i and j .

Throughout the following discussion, with n^μ the jet direction and \bar{n}^μ the direction opposite the jet, we will describe momenta in light-cone coordinates

$$p^\mu = (p^-, p^+, p_\perp) \quad (3)$$

with

$$p^- = \bar{n} \cdot p \quad p^+ = n \cdot p \quad (4)$$

¹Indeed, the calculation of Frye et al. is a more general factorization of mass-like variables in groomed jets. Setting $\alpha = 2, \beta = 0$ for their two-point energy correlation function $e_2^{(\alpha)}$ under soft drop grooming with angular exponent β yields the mMDT-groomed jet mass ρ . Their factorization is valid in the limit $\rho \ll z_{\text{cut}} \ll 1$, whereas we are interested in the limit $\rho \sim z_{\text{cut}} \ll 1$.

and p_\perp the components of momentum transverse to n . In these coordinates, the energy fraction with respect to total energy $E_J = Q$ is

$$z = \frac{p^+ + p^-}{2Q} \quad (5)$$

and, in the collinear limit, the angle to the jet axis is $\theta \approx p_\perp/p^0$ [2].

In an $e^+e^- \rightarrow \text{jets}$ event, there are two types of emission: resolved and unresolved. The essential difference is that a resolved emission is one which manifests itself as a jet at a particular scale of observation, while an unresolved emission does not. The presence of unresolved emissions can, however, perturb observable values of a resolved emission. **[TODO: check that this is a reasonable description]**

Suppose now that we have applied an mMDT groomer with energy fraction cutoff z_{cut} . Then every *resolved* emission must satisfy

$$z_i > z_{\text{cut}}, \quad (6)$$

while other emissions with $z_i < z_{\text{cut}}$ can only pass the groomer if they are at a sufficiently small angle to a resolved emission.

0.2 Power counting

0.2.1 Resolved soft emission

The primary emission contributing to the jet mass in the limit $\rho \sim z_{\text{cut}} \ll 1$ is a gluon emission z_i sensitive to both ρ and z_{cut} . In the presence of a hard quark (i.e., the jet) with $z_q \sim 1$, leading contributions to the jet mass of Eq. 2 is

$$\rho \approx \sum_i 2z_i(1 - \cos \theta_i) \quad (7)$$

where θ_i is the angle of emission i from the quark. Considering the case with only one such emission,² we have

$$\rho \approx z_i(1 - \cos \theta_i). \quad (8)$$

But since $z_i \sim z_{\text{cut}}$ and $\rho \sim z_{\text{cut}}$, this means that

$$\rho \approx \rho(1 - \cos \theta_i), \quad (9)$$

so

$$\cos \theta_i \ll 1. \quad (10)$$

This means that

$$\theta_i \sim \frac{\pi}{2}. \quad (11)$$

²This is the one-loop contribution **[is this accurate?]**.

Since this emission is sensitive to $z_{\text{cut}} \ll 1$, we can also conclude that $z_i \ll 1$. Thus, we see that the leading contribution to the cusp region comes from a **resolved soft, wide-angle** gluon. Its momentum scales like

$$p_R \sim z_{\text{cut}} Q(1, 1, 1). \quad (12)$$

For future reference, let this gluon have energy fraction z_R and angle θ_R from the quark axis.

0.2.2 Ungroomed extra-soft unresolved radiation




Figure 1: Head-on schematic of a quark jet q and a resolved gluon g . If the angle between the quark and the gluon is θ and the gluon (plus all lower-scale unresolved radiation) passes the groomer, then the mMDT groomer will accept all radiation within an angle θ of both the quark and the resolved gluon.

Unresolved soft emissions can also contribute to the jet mass if they are sufficiently close to the resolved emission or the quark. Suppose there is another emission i with energy fraction z_i at angle θ_i from the jet axis and angle θ_{iR} from the resolved gluon. If $\theta_i < \theta_R$ or $\theta_{iR} < \theta_R$, a situation displayed in Fig. 1, then emission i will pass the groomer.

What does this mean for the hemisphere mass? Well, first notice that since $\theta_R \sim \pi/2$, most soft radiation in the hemisphere passes this cut. The effect of these extra-soft emissions, which have $z_i \ll z_R$, is to perturb the mass of the resolved emission, so that the hemisphere mass is approximately

$$\rho \sim z_R + \sum_i z_i (1 - \cos \theta_i). \quad (13)$$

The dominant contributions come again from the wide-angle emissions with $1 - \cos \theta_i \sim 1$; these must evidently have an energy scale set by

$$\rho - z_R \sim \sum_i z_i. \quad (14)$$

Hence, since $z_R \sim z_{\text{cut}}$, the unresolved extra-soft emissions must scale as

$$p_{S_R} \sim (\rho - z_{\text{cut}}) Q(1, 1, 1). \quad (15)$$

0.2.3 Collinear radiation

Finally, we consider radiation collinear to a jet axis with angle $\theta_i \ll 1$. This radiation has $p^- \gg p^+$, which means from Eq. 5 that

$$z \approx \frac{p^+}{2Q}. \quad (16)$$

Then because the particle must satisfy

$$z_i \theta_i^2 \lesssim \rho, \quad (17)$$

we find that [2]

$$\rho \sim \frac{p^+}{Q}. \quad (18)$$

If $z \sim 1$, we know that $z \gg z_{\text{cut}}$, so the momentum scales independently of z_{cut} . Hence, the scaling of these **collinear** modes is [2]

$$p_c \sim Q(1, \rho, \rho^{1/2}). \quad (19)$$

In a hemisphere where the resolved emission stops the mMDT groomer, this is the only collinear radiation that contributes **[TODO: ask Andrew: is this correct?]**.

If on the other hand $z \sim z_{\text{cut}} \ll 1$, the result is **collinear-soft** radiation with $p^- \sim z_i Q$ and $p^+ \sim \theta_i^2 z_i Q$. From [2], these momenta scale like

$$p_{cs} \sim z_{\text{cut}} Q \left(1, \frac{\rho}{z_{\text{cut}}}, \left(\frac{\rho}{z_{\text{cut}}} \right)^{1/2} \right) \quad (20)$$

and depend on the single energy scale $\sqrt{\rho z_{\text{cut}}}$. This scale matters in the hemisphere which does not contain the resolved soft gluon.

0.3 Factorization

With the power counting in hand, we are now ready to derive a factorization formula describing the hemisphere mass distribution in the limit $\rho \sim z_{\text{cut}} \ll 1$. First, we should note that it is most straightforward to compute the double differential cross section in the masses of the individual hemispheres, then integrate over them to get the heavy hemisphere mass [3]:

$$\frac{d\sigma}{d\rho} = \int \frac{d^2\sigma}{d\rho_1 d\rho_2} [\delta(\rho - \rho_1) \Theta(\rho_1 - \rho_2) + \delta(\rho - \rho_2) \Theta(\rho_2 - \rho_1)]. \quad (21)$$

The integral simply breaks up the two cases $\rho_1 > \rho_2$ and $\rho_2 > \rho_1$ and assigns the correct value of ρ in each case.

Now, in the limit $\rho_1, \rho_2 \ll 1$, we can apply the technology of SCET to factorize the double-differential cross section into a product of hard, soft, and jet contributions [2, 4]. The basic form is

$$\frac{d^2\sigma}{d\rho_1 d\rho_2} = H(Q^2) \otimes S(\rho_1, \rho_2, z_{\text{cut}}) \otimes J_q(\rho_1) \otimes J_g(\rho_1, z_{\text{cut}}) \otimes J_{\bar{q}}(\rho_2). \quad (22)$$

The symbol \otimes represents convolution. Here, Q^2 is the squared center-of-mass energy of the collision. $H(Q^2)$ is hard function representing the cross section of $e^+e^- \rightarrow q\bar{q}$ events, $S(\rho_1, \rho_2, z_{\text{cut}})$ is the function representing soft contributions (which are sensitive to z_{cut}), and $J_i(\rho)$ is a function describing the production of a jet off of particle i (where i is a quark q , antiquark \bar{q} , or gluon g).

As this factorization currently stands, several terms depend on multiple scales and must be refactorized. In the limit $\rho \sim z_{\text{cut}} \ll 1$ after mMDT grooming, the soft function consists of global soft emissions which contribute only to the normalization; a resolved soft, wide-angle emission generated by a fixed-order function; and soft radiation which passes the groomer due to the resolved emission. The first two depend only on the scale z_{cut} and can be combined into one function. Therefore, we can write

$$\begin{aligned} \frac{d^2\sigma}{d\rho_1 d\rho_2} = & H(Q^2) \times R(\rho_1, \rho_2, z_{\text{cut}}) \times S_R(\rho - z_{\text{cut}}) \otimes J_{c,q}(\rho_1, z_{\text{cut}}) \\ & \otimes J_{c,g}(\rho_1, z_{\text{cut}}) \otimes J_{c,\bar{q}}(\rho_2, z_{\text{cut}}). \end{aligned} \quad (23)$$

Here, $R(\rho_1, \rho_2, z_{\text{cut}})$ describes the resolved emission as well as other soft wide-angle radiation (from which the resolved emission emerged), while $S_R(\rho - z_{\text{cut}})$ describes radiation which passes the groomer in the presence of the resolved emission. We have also re-written the jet functions as $J_{c,i}(\rho, z_{\text{cut}})$ to make explicit their dependence on multiple scales.

Now, as we established in Sec. 0.2.3, radiation collinear to the jet with energy fraction of order 1 depends only on ρ , and radiation with energy fraction much less than 1 depends on the scale $\sqrt{\rho z_{\text{cut}}}$. However, this soft-collinear radiation only matters in the absence of a resolved gluon emission (which stops the mMDT groomer), in which case the jet function factorizes as

$$J_{c,\bar{q}}(\rho, z_{\text{cut}}) = J_{\bar{q}}(\rho) \otimes S_C(\sqrt{\rho z_{\text{cut}}}) \quad (24)$$

The result is that we must treat each hemisphere separately, since one contains the resolved emission and the other does not. Moreover, the gluon's jet function J_g only depends on the scale of the gluon's energy, which is a function of ρ and z_{cut} **[TODO: check this]**. Therefore, if we assume, without loss of generality, that $\rho_1 > \rho_2$, the cross section becomes

$$\boxed{\begin{aligned} \frac{d^2\sigma}{d\rho_1 d\rho_2} = & 2H(Q^2) \times R(\rho_1, \rho_2, z_{\text{cut}}) \times S_R(\rho - z_{\text{cut}}) \otimes J_{c,q}(\rho_1) \\ & \otimes J_{c,g}(\rho_1, z_{\text{cut}}) \otimes [J_{\bar{q}}(\rho_2) \otimes S_C(\sqrt{\rho_2 z_{\text{cut}}})]. \end{aligned}} \quad (25)$$

This is our final factorization formula for the double differential cross section in each hemisphere mass. The distribution of the heavy hemisphere mass then given by Eq. 21.

References

- [1] T. Becher, A. Broggio and A. Ferroglia, *Introduction to Soft-Collinear Effective Theory*, vol. 896 of *Lecture Notes in Physics*, Springer International Publishing, Cham (2015), 10.1007/978-3-319-14848-9.
- [2] C. Frye, A.J. Larkoski, M.D. Schwartz and K. Yan, *Factorization for groomed jet substructure beyond the next-to-leading logarithm*, *J. High Energ. Phys.* **2016** (2016) 64.
- [3] Y.-T. Chien and M.D. Schwartz, *Resummation of heavy jet mass and comparison to LEP data*, *J. High Energ. Phys.* **2010** (2010) 58.
- [4] S.D. Ellis, C.K. Vermilion, J.R. Walsh, A. Hornig and C. Lee, *Jet shapes and jet algorithms in SCET*, *J. High Energ. Phys.* **2010** (2010) 101.