

Gibbs point processes and number-rigidity

Thibaut Vasseur
Université Paris Cité


Hyperuniform structures, rigid point processes
and related topics

Point process Γ : random variable on Config, the locally finite subsets of \mathbb{R}^d

Point process Γ : random variable on Config, the locally finite subsets of \mathbb{R}^d

Γ is *number-rigid* if for every compact set Δ there exists a function $F_\Delta : \text{Config} \rightarrow \mathbb{N}$ such that almost surely


$$\#\Gamma_\Delta = F_\Delta(\Gamma_{\Delta^c}).$$



Point process Γ : random variable on Config , the locally finite subsets of \mathbb{R}^d

Γ is *number-rigid* if for every compact set Δ there exists a function $F_\Delta : \text{Config} \rightarrow \mathbb{N}$ such that almost surely


$$\#\Gamma_\Delta = F_\Delta(\Gamma_{\Delta^c}).$$



Point process Γ : random variable on Config, the locally finite subsets of \mathbb{R}^d

Γ is *number-rigid* if for every compact set Δ there exists a function $F_\Delta : \text{Config} \rightarrow \mathbb{N}$ such that almost surely

$$\#\Gamma_\Delta = F_\Delta(\Gamma_{\Delta^c}).$$



- 1D Coulomb gas (Aizenman, Martin, 1980)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)
- Ginibre (Ghosh, Perez, 2017)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)
- Ginibre (Ghosh, Perez, 2017)
- Perturbed Lattices (Holroyd, Soo 2014, Peres, Sly 2014)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)
- Ginibre (Ghosh, Perez, 2017)
- Perturbed Lattices (Holroyd, Soo 2014, Peres, Sly 2014)
- Stable matching (Klatt, Last, Yogeshwaran, 2020)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)
- Ginibre (Ghosh, Perez, 2017)
- Perturbed Lattices (Holroyd, Soo 2014, Peres, Sly 2014)
- Stable matching (Klatt, Last, Yogeshwaran, 2020)
- Sine_β (1D log-gas) | (Chaïbi, Najnudel, 2018)
(Dereudre, Hardy, Leblé, Maïda, 2020)

- 1D Coulomb gas (Aizenman, Martin, 1980)
- zeros of the Planar GAF (Ghosh, Perez, 2017)
- Ginibre (Ghosh, Perez, 2017)
- Perturbed Lattices (Holroyd, Soo 2014, Peres, Sly 2014)
- Stable matching (Klatt, Last, Yogeshwaran, 2020)
- Sine_β (1D log-gas) | (Chaïbi, Najnudel, 2018)
(Dereudre, Hardy, Leblé, Maïda, 2020)

For Gibbs point processes, dependency is described through an energy function H .

Pairwise interaction

$$H(\gamma) = \sum_{\{x,y\} \subset \gamma} g(y-x)$$

For Gibbs point processes, dependency is described through an energy function H .

Pairwise interaction

$$H(\gamma) = \sum_{\{x,y\} \subset \gamma} g(y-x)$$

The Riesz interaction with parameter $s \in (-2, +\infty)$

$$g(x) = \begin{cases} |x|^{-s} & \text{for } s > 0 \\ -\log|x| & \text{for } s = 0 \\ -|x|^{-s} & \text{for } s \in (-2, 0) \end{cases}$$

For Gibbs point processes, dependency is described through an energy function H .

Pairwise interaction

$$H(\gamma) = \sum_{\{x,y\} \subset \gamma} g(y-x)$$

The Riesz interaction with parameter $s \in (-2, +\infty)$

$$g(x) = \begin{cases} |x|^{-s} & \text{for } s > 0 \\ -\log|x| & \text{for } s = 0 \\ -|x|^{-s} & \text{for } s \in (-2, 0) \end{cases}$$

$s = d - 2$ is the Coulomb interaction

$\Lambda_n = [-n^{1/d}/2, n^{1/d}/2]$: square of volume n

The finite volume canonical Gibbs measure is

$$P_n^\beta(d\gamma) = \frac{1}{Z_n} e^{-\beta H(\gamma)} \text{Bin}_{\Lambda_n, n}(d\gamma).$$

$\Lambda_n = [-n^{1/d}/2, n^{1/d}/2]$: square of volume n

The finite volume canonical Gibbs measure is

$$P_n^\beta(d\gamma) = \frac{1}{Z_n} e^{-\beta H(\gamma)} \text{Bin}_{\Lambda_n, n}(d\gamma).$$

Infinite volume Gibbs point process : an accumulation points P^β of the sequence $(P_n^\beta)_{n \geq 1}$

The periodic setting can be more convenient in the long range case

Modèle périodique

$$H_n(\gamma) = \sum_{\{x,y\} \subset \gamma} g_n(x-y)$$

$$\text{where } g_n(x) = \sum_{u \in \mathbb{Z}^d} \left[g(x + n^{1/d}u) - n^{-1} \int_{\Lambda_n} g(y + n^{1/d}u) dy \right].$$

and $g(x) = |x|^{-s}$ with $s \in (d-1, d)$.

Again, the finite volume canonical Gibbs measure is

$$P_n^\beta(d\gamma) = \frac{1}{Z_n} e^{-\beta H_n(\gamma)} \text{Bin}_{\Lambda_n, n}(d\gamma).$$

$d = 1, s = 0$ ($g(x) = -\log|x|$) one acc. point : Sine process

Law of Γ_Δ is inaccessible

Law of Γ_Δ is inaccessible

Grand canonical DLR equations

$$\Gamma_\Delta \mid \Gamma_{\Delta^c} = \gamma_{\Delta^c} \sim \frac{1}{Z(\gamma_{\Delta^c})} \exp(-\beta(H(\eta) + W(\eta \mid \gamma_{\Delta^c}))) \text{Pois}_\Delta(d\eta),$$

where $W(\eta \mid \gamma_{\Delta^c}) = \sum_{x \in \eta} \sum_{y \in \gamma_{\Delta^c}} g(x - y)$.

Law of Γ_Δ is inaccessible

Grand canonical DLR equations

$$\Gamma_\Delta \mid \Gamma_{\Delta^c} = \gamma_{\Delta^c} \sim \frac{1}{Z(\gamma_{\Delta^c})} \exp(-\beta(H(\eta) + W(\eta \mid \gamma_{\Delta^c}))) \text{Pois}_\Delta(d\eta),$$

where $W(\eta \mid \gamma_{\Delta^c}) = \sum_{x \in \eta} \sum_{y \in \gamma_{\Delta^c}} g(x - y)$.

Adapted to short range ($s > d$) for which W is well defined.

Canonical Setting can be convenient for the long range case
($s < d$)

Canonical Setting can be convenient for the long range case
($s < d$)

Canonical DLR equations

$$\Gamma_{\Delta} \mid \Gamma_{\Delta^c} = \gamma_{\Delta^c}, \# \Gamma_{\Delta} = n \\ \sim \frac{1}{Z(\gamma_{\Delta^c})} \exp(-\beta(H(\eta) + \text{Move}(\eta \mid \gamma_{\Delta^c}))) \text{Bin}_{\Delta,n}(d\eta),$$

where $\text{Move}(\eta \mid \gamma_{\Delta^c}) = W(\eta \mid \gamma_{\Delta^c}) - W(\text{ref} \mid \gamma_{\Delta^c})$

Canonical Setting can be convenient for the long range case
($s < d$)

Canonical DLR equations

$$\Gamma_{\Delta} \mid \Gamma_{\Delta^c} = \gamma_{\Delta^c}, \# \Gamma_{\Delta} = n \\ \sim \frac{1}{Z(\gamma_{\Delta^c})} \exp(-\beta(H(\eta) + \text{Move}(\eta \mid \gamma_{\Delta^c}))) \text{Bin}_{\Delta,n}(d\eta),$$

where $\text{Move}(\eta \mid \gamma_{\Delta^c}) = W(\eta \mid \gamma_{\Delta^c}) - nW(0 \mid \gamma_{\Delta^c})$

Canonical Setting can be convenient for the long range case
($s < d$)

Canonical DLR equations

$$\Gamma_{\Delta} \mid \Gamma_{\Delta^c} = \gamma_{\Delta^c}, \# \Gamma_{\Delta} = n \\ \sim \frac{1}{Z(\gamma_{\Delta^c})} \exp(-\beta(H(\eta) + \text{Move}(\eta \mid \gamma_{\Delta^c}))) \text{Bin}_{\Delta,n}(d\eta),$$

where $\text{Move}(\eta \mid \gamma_{\Delta^c}) = W(\eta \mid \gamma_{\Delta^c}) - nW(0 \mid \gamma_{\Delta^c})$

Adding information on the inside helps to define interaction with the outside

The n -points reduced Campbell measure is defined as

$$\text{Campbell}_P^{(n)}(f) = E \left[\sum_{x_1, \dots, x_n \in \Gamma}^{\neq} f(x_1, \dots, x_n, \Gamma \setminus \{x_1, \dots, x_n\}) \right]$$

where $f : (\mathbb{R}^d)^n \times \text{Config} \rightarrow \mathbb{R}$ positive and measurable.

- If $\Gamma \sim \text{Pois}$ then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = dX_n \text{Pois}(d\gamma)$$

- If $\Gamma \sim \text{Pois}$ then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = dX_n \text{Pois}(d\gamma)$$

- If $\Gamma \sim \text{Bin}_{\Lambda, N}$ and $n \leq N$ then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = \frac{1}{|\Lambda|^n} \frac{N!}{(N-n)!} d_{\Lambda} X_n \text{Bin}_{\Lambda, N-n}(d\gamma)$$

- If $\Gamma \sim \text{Pois}$ then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = dX_n \text{Pois}(d\gamma)$$

- If $\Gamma \sim \text{Bin}_{\Lambda, N}$ and $n \leq N$ then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = \frac{1}{|\Lambda|^n} \frac{N!}{(N-n)!} d_{\Lambda} X_n \text{Bin}_{\Lambda, N-n}(d\gamma)$$

- If Γ satisfies the previous Canonical DLR equations then

$$\text{Campbell}^{(n)}(dX_n d\gamma) = \exp(-\beta(H(X_n) + \text{Move}(X_n, \gamma))) dX_n Q_n^{\beta}(d\gamma)$$

Dereudre, Hardy, Leblé, Maïda

The Sine process satisfies canonical DLR equations and is number-rigid.

Dereudre, Hardy, Leblé, Maïda

The Sine process satisfies canonical DLR equations and is number-rigid.

Ideas of the proof :

Dereudre, Hardy, Leblé, Maïda

The Sine process satisfies canonical DLR equations and is number-rigid.

Ideas of the proof :

- Consequence of the non-number rigidity $\exists n \geq 1 : Q_n^\beta \ll P^\beta$

Dereudre, Hardy, Leblé, Maïda

The Sine process satisfies canonical DLR equations and is number-rigid.

Ideas of the proof :

- Consequence of the non-number rigidity $\exists n \geq 1 : Q_n^\beta \ll P^\beta$
- Use the n-points Campbell measure to obtain a contradiction.

Dereudre, Hardy, Leblé, Maïda

The Sine process satisfies canonical DLR equations and is number-rigid.

Ideas of the proof :

- Consequence of the non-number rigidity $\exists n \geq 1 : Q_n^\beta \ll P^\beta$
- Use the n -points Campbell measure to obtain a contradiction.

We assume $n = 1$, so there exist a function Add such that

$$Q_1^\beta(d\gamma) = e^{-\beta \text{Add}(\gamma)} P^\beta(d\gamma)$$

1-point Campbell measure

$$\text{Campbell}(dx d\gamma) = \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) dx P^\beta(d\gamma)$$

1-point Campbell measure

$$\text{Campbell}(dx d\gamma) = \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) dx P^\beta(d\gamma)$$

2-points Campbell measure

$$\begin{aligned} \text{Campbell}^{(2)}(dx dy d\gamma) = & \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) \\ & \exp(-\beta(\text{Move}(y, \gamma \cup \{x\}) + \text{Add}(\gamma \cup \{x\}))) dx dy P^\beta(d\gamma) \end{aligned}$$

1-point Campbell measure

$$\text{Campbell}(dx d\gamma) = \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) dx P^\beta(d\gamma)$$

2-points Campbell measure

$$\begin{aligned} \text{Campbell}^{(2)}(dx dy d\gamma) &= \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) \\ &\quad \exp(-\beta(\text{Move}(y, \gamma \cup \{x\}) + \text{Add}(\gamma \cup \{x\}))) dx dy P^\beta(d\gamma) \end{aligned}$$

Manipulating Campbell measures shows that

$$\text{Add}(\gamma \cup \{x\}) + \log |x| = \text{Add}(\gamma \cup \{z\}) + \log |z|$$

1-point Campbell measure

$$\text{Campbell}(dx d\gamma) = \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) dx P^\beta(d\gamma)$$

2-points Campbell measure

$$\begin{aligned} \text{Campbell}^{(2)}(dx dy d\gamma) &= \exp(-\beta(\text{Move}(x, \gamma) + \text{Add}(\gamma))) \\ &\quad \exp(-\beta(\text{Move}(y, \gamma \cup \{x\}) + \text{Add}(\gamma \cup \{x\}))) dx dy P^\beta(d\gamma) \end{aligned}$$

Manipulating Campbell measures shows that

$$\text{Add}(\gamma \cup \{x\}) \rightarrow -\infty$$

What we have used


Structure of the campbell measure, stationarity and the fact that $g(x) \rightarrow -\infty$ when $|x| \rightarrow \infty$.


What we have used

Structure of the campbell measure, stationarity and the fact that $g(x) \rightarrow -\infty$ when $|x| \rightarrow \infty$.

Leblé, V. (ongoing work)


2D coulomb gases satisfy canonical DLR equations and is number-rigid.





DLR equations and a control of the cost of exchanging a k point configuration in Δ with a l points configuration in $\Delta + u$ (independently of u) then for any $\Lambda \setminus \Delta$ -local event

$$P(\#\Gamma_{\Delta} = k, \#\Gamma_{\Delta+u} = l, A) \leq CP(\#\Gamma_{\Delta} = l, \#\Gamma_{\Delta+u} = k, A) + \varepsilon$$



DLR equations and a control of the cost of exchanging a k point configuration in Δ with a l points configuration in $\Delta + u$ (independently of u) then for any $\Lambda \setminus \Delta$ -local event

$$P(\#\Gamma_{\Delta} = k, \#\Gamma_{\Delta+u} = l, A) \leq CP(\#\Gamma_{\Delta} = l, \#\Gamma_{\Delta+u} = k, A) + \varepsilon$$

$$P(\#\Gamma_{\Delta} = k \mid \Gamma_{\Delta^c}) > 0 \implies P(\#\Gamma_{\Delta} = l \mid \Gamma_{\Delta^c}) > 0$$

Hyperuniformity can help to show non-number-rigidity.

Hyperuniformity can help to show non-number-rigidity.

Control the cost of exchange configuration in Δ and $\Delta + u$ can be achieved if we show that

$$\lim_{\Lambda \rightarrow \mathbb{R}^d} \sum_{y \in \Gamma_\Lambda} g(x - y) - \int_\Lambda g(x - y) dy$$

exists and is almost surely finite.

Hyperuniformity can help to show non-number-rigidity.

Control the cost of exchange configuration in Δ and $\Delta + u$ can be achieved if we show that

$$\lim_{\Lambda \rightarrow \mathbb{R}^d} \sum_{y \in \Gamma_\Lambda} g(x - y) - \int_\Lambda g(x - y) dy$$

exists and is almost surely finite.

For instance, when $s \in (d - 1, d)$ a sufficient condition is

$$\text{Var}(\#\Gamma_{\Lambda_k}) \leq Ck^{2s-\varepsilon}.$$

Dereudre, V.

For the periodic Riesz gas with $s \in (d - 1, d)$ there is an accumulation point which is non-number-rigid.

Dereudre, V.

For the periodic Riesz gas with $s \in (d - 1, d)$ there is an accumulation point which is non-number-rigid.

An interesting consequence for the one point Campbell measure

$$\text{Campbell}(dx, d\gamma) = e^{-h(x, \gamma)} dx P(d\gamma)$$

Which implies grand canonical DLR equations.

Dereudre, V.

For the periodic Riesz gas with $s \in (d - 1, d)$ there is an accumulation point which is non-number-rigid.

An interesting consequence for the one point Campbell measure

$$\text{Campbell}(dx, d\gamma) = e^{-h(x, \gamma)} dx P(d\gamma)$$

Which implies grand canonical DLR equations.

Existence of a compensator

$$h(x, \Gamma) = \lim_{n \rightarrow +\infty} \sum_{y \in \Gamma_{\Lambda_n}} g(x - y) - \text{Comp}(\Lambda_n, \Gamma_{\Lambda_n^c}, \#\Gamma_{\Lambda_n})$$

Thank you for your attention !