

Hyperuniformity(?) and hyperfluctuation in "curved" spaces

Michael Björklund (Chalmers/GU)

Based on joint works with Tobias Hartnick (Karlsruhe) and Mattias Byléhn (Chalmers/GU)

General framework - Random measures in metric spaces I

Some notation:

Let (X, d) be a metric space, fix a point $x_o \in X$, and assume that the closed ball

$$B_R := \{x \in X : d(x, x_o) \leq R\} \quad \text{is compact for all } R \geq 0.$$

General framework - Random measures in metric spaces I

Some notation:

Let (X, d) be a metric space, fix a point $x_o \in X$, and assume that the closed ball

$$B_R := \{x \in X : d(x, x_o) \leq R\} \quad \text{is compact for all } R \geq 0.$$

Homogeneity: Suppose that $G < \text{Isom}(X, d)$ is a closed transitive subgroup, and thus

$$X = G.x_o \cong G/K, \quad \text{where } K := \text{Stab}_G(x_o) < G \text{ is compact.}$$

General framework - Random measures in metric spaces I

Some notation:

Let (X, d) be a metric space, fix a point $x_o \in X$, and assume that the closed ball

$$B_R := \{x \in X : d(x, x_o) \leq R\} \quad \text{is compact for all } R \geq 0.$$

Homogeneity: Suppose that $G < \text{Isom}(X, d)$ is a closed transitive subgroup, and thus

$$X = G.x_o \cong G/K, \quad \text{where } K := \text{Stab}_G(x_o) < G \text{ is compact.}$$

Let $M^+(X)$ denote the space of positive Radon measures on X , and let μ be a locally square-integrable G -invariant probability measure on $M^+(X)$, i.e.

$$\int_{M^+(X)} p(B_R)^2 d\mu(p) < \infty, \quad \text{for all } R \geq 0.$$

General framework - Random measures in metric spaces II

Volume on X : Fix a left Haar measure m_G on G and define

$$\int_X f \, dm_X = \int_G f(g.x_0) \, dm_G(g), \quad \text{for all } f \in \mathcal{L}_c^\infty(X),$$

so that m_X is a G -invariant Radon measure on X (= Volume). Let m_K denote the Haar probability measure on K .

General framework - Random measures in metric spaces II

Volume on X : Fix a left Haar measure m_G on G and define

$$\int_X f \, dm_X = \int_G f(g.x_0) \, dm_G(g), \quad \text{for all } f \in \mathcal{L}_c^\infty(X),$$

so that m_X is a G -invariant Radon measure on X (= Volume). Let m_K denote the Haar probability measure on K .

Linear statistics: For $f \in \mathcal{L}_c^\infty(X)$, define $Sf(p) = \int_X f \, dp$ for $p \in M^+(X)$.

General framework - Random measures in metric spaces II

Volume on X : Fix a left Haar measure m_G on G and define

$$\int_X f \, dm_X = \int_G f(g.x_0) \, dm_G(g), \quad \text{for all } f \in \mathcal{L}_c^\infty(X),$$

so that m_X is a G -invariant Radon measure on X (= Volume). Let m_K denote the Haar probability measure on K .

Linear statistics: For $f \in \mathcal{L}_c^\infty(X)$, define $Sf(p) = \int_X f \, dp$ for $p \in M^+(X)$.

Note: $Sf \in L^2(\mu)$ for all $f \in \mathcal{L}_c^\infty(X)$ and there exists $\iota_\mu \geq 0$ (intensity of μ) such that

$$\mu(Sf) = \iota_\mu m_X(f), \quad \text{for all } f \in \mathcal{L}_c^\infty(X).$$

Geometric hyperuniformity?

Naive definition: μ is (geometrically) hyperuniform if

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_\chi(B_R)), \quad \text{as } R \rightarrow \infty.$$

Geometric hyperuniformity?

Naive definition: μ is (geometrically) hyperuniform if

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_X(B_R)), \quad \text{as } R \rightarrow \infty.$$

"Problem": Geometrically hyperuniform examples simply do not exist in many geometries (for instance in hyperbolic spaces [B-Bylehn '23]).

Geometric hyperuniformity?

Naive definition: μ is (geometrically) hyperuniform if

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_X(B_R)), \quad \text{as } R \rightarrow \infty.$$

"Problem": Geometrically hyperuniform examples simply do not exist in many geometries (for instance in hyperbolic spaces [B-Bylehn '23]).

Questions:

- ▶ How should spectral hyperuniformity be defined?
- ▶ Do examples exist?
- ▶ What should spectral hyperuniformity correspond to "in the real world"?

Geometric hyperuniformity?

Naive definition: μ is (geometrically) hyperuniform if

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_X(B_R)), \quad \text{as } R \rightarrow \infty.$$

"Problem": Geometrically hyperuniform examples simply do not exist in many geometries (for instance in hyperbolic spaces [B-Bylehn '23]).

Questions:

- ▶ How should spectral hyperuniformity be defined?
- ▶ Do examples exist?
- ▶ What should spectral hyperuniformity correspond to "in the real world"?

Some potential answers below...

Hyperfluctuations

Fact: If (B_R) form a nice ergodic averaging sequence and μ is G -ergodic, then

$$\mathrm{Var}_\mu(S_{\chi_{B_R}}) = o(m_X(B_R)^2), \quad \text{as } R \rightarrow \infty.$$

Hyperfluctuations

Fact: If (B_R) form a nice ergodic averaging sequence and μ is G -ergodic, then

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_X(B_R)^2), \quad \text{as } R \rightarrow \infty.$$

When $X = \mathbb{R}^d$ or a (real/complex) hyperbolic space, then for every $\delta > 0$ there exists a random measure μ_δ (in fact, a point process) such that

$$\mathrm{Var}_{\mu_\delta}(S\chi_{B_R}) \gg m_X(B_R)^{2-\delta}, \quad \text{as } R \rightarrow \infty.$$

Hyperfluctuations

Fact: If (B_R) form a nice ergodic averaging sequence and μ is G -ergodic, then

$$\mathrm{Var}_\mu(S\chi_{B_R}) = o(m_X(B_R)^2), \quad \text{as } R \rightarrow \infty.$$

When $X = \mathbb{R}^d$ or a (real/complex) hyperbolic space, then for every $\delta > 0$ there exists a random measure μ_δ (in fact, a point process) such that

$$\mathrm{Var}_{\mu_\delta}(S\chi_{B_R}) \gg m_X(B_R)^{2-\delta}, \quad \text{as } R \rightarrow \infty.$$

Theorem (B-Bylehn '23) If $X =$ quaterionic hyperbolic space, or a higher rank symmetric space (for instance, the space of positive definite matrices in \mathbb{R}^n for $n \geq 3$), then there exists $\delta_X > 0$ such that for every random measure μ on X , we have

$$\mathrm{Var}_\mu(S\chi_{B_R}) \ll m_X(B_R)^{2-\delta_X}, \quad \text{as } R \rightarrow \infty.$$

What should we mean by spectral hyperuniformity?

In our approach to this question, it is crucial that (G, K) is a so called **Gelfand pair**.
What does that mean?

What should we mean by spectral hyperuniformity?

In our approach to this question, it is crucial that (G, K) is a so called **Gelfand pair**.
What does that mean?

Let $\mathcal{L}_c^\infty(G, K)$ denote the space of bi- K -invariant bounded Borel functions with bounded supports on G . Note that $\mathcal{L}_c^\infty(G, K) * \mathcal{L}_c^\infty(G, K) \subset \mathcal{L}_c^\infty(G, K)$.

What should we mean by spectral hyperuniformity?

In our approach to this question, it is crucial that (G, K) is a so called **Gelfand pair**. What does that mean?

Let $\mathcal{L}_c^\infty(G, K)$ denote the space of bi- K -invariant bounded Borel functions with bounded supports on G . Note that $\mathcal{L}_c^\infty(G, K) * \mathcal{L}_c^\infty(G, K) \subset \mathcal{L}_c^\infty(G, K)$.

We say that (G, K) is a **Gelfand pair** if the algebra $(\mathcal{L}_c^\infty(G, K), *)$ is commutative, i.e. if for every $\varphi \in \mathcal{L}_c^\infty(G, K)$, we have

$$\int_K \varphi(g_1 k g_2) dm_K(k) = \int_K \varphi(g_2 k g_1) dm_K(k), \quad \text{for all } g_1, g_2 \in G.$$

Fact: If (G, K) is a Gelfand pair, then m_G must be bi- G -invariant (unimodular).

Zonal spherical functions

A bi- K -invariant function $\omega : G \rightarrow \mathbb{C}$ is a *zonal spherical function* if it is positive definite and satisfies

$$\int_K \omega(g_1 k g_2) dm_K(k) = \omega(g_1) \omega(g_2), \quad \text{for all } g_1, g_2 \in G.$$

Let Ω denote the space of all zonal functions for (G, K) , equipped with the topology of uniform convergence on compact sets. We can (sloppily) think of these as K -invariant functions on X .

Zonal spherical functions

A bi- K -invariant function $\omega : G \rightarrow \mathbb{C}$ is a *zonal spherical function* if it is positive definite and satisfies

$$\int_K \omega(g_1 k g_2) dm_K(k) = \omega(g_1) \omega(g_2), \quad \text{for all } g_1, g_2 \in G.$$

Let Ω denote the space of all zonal functions for (G, K) , equipped with the topology of uniform convergence on compact sets. We can (sloppily) think of these as K -invariant functions on X .

Given $f \in \mathcal{L}_c^\infty(X)^K$, define the **spherical Fourier transform** $\hat{f} : \Omega \rightarrow \mathbb{C}$ by

$$\hat{f}(\omega) = \int_X f(x) \overline{\omega(x)} dm_X(x), \quad \text{for } \omega \in \Omega.$$

Plancherel and Godement

Fact: There exists a unique positive measure σ (**=Plancherel measure**) on Ω such that

$$\int_X |f(x)|^2 dm_X(x) = \int_{\Omega} |\widehat{f}(\omega)|^2 d\sigma(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

Plancherel and Godement

Fact: There exists a unique positive measure σ (**=Plancherel measure**) on Ω such that

$$\int_X |f(x)|^2 dm_X(x) = \int_{\Omega} |\hat{f}(\omega)|^2 d\sigma(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

Theorem (Godement) There exists a unique positive Radon measure $\hat{\eta}_\mu$ on Ω such that

$$\text{Var}_\mu(Sf) = \int_{\Omega} |\hat{f}(\omega)|^2 d\hat{\eta}_\mu(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

Plancherel and Godement

Fact: There exists a unique positive measure σ (**=Plancherel measure**) on Ω such that

$$\int_X |f(x)|^2 dm_X(x) = \int_{\Omega} |\hat{f}(\omega)|^2 d\sigma(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

Theorem (Godement) There exists a unique positive Radon measure $\hat{\eta}_\mu$ on Ω such that

$$\text{Var}_\mu(Sf) = \int_{\Omega} |\hat{f}(\omega)|^2 d\hat{\eta}_\mu(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

We refer to $\hat{\eta}_\mu$ as the **diffraction measure** of μ .

Plancherel and Godement

Fact: There exists a unique positive measure σ (**=Plancherel measure**) on Ω such that

$$\int_X |f(x)|^2 dm_X(x) = \int_{\Omega} |\hat{f}(\omega)|^2 d\sigma(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

Theorem (Godement) There exists a unique positive Radon measure $\hat{\eta}_\mu$ on Ω such that

$$\text{Var}_\mu(Sf) = \int_{\Omega} |\hat{f}(\omega)|^2 d\hat{\eta}_\mu(\omega), \quad \text{for all } f \in \mathcal{L}_c^\infty(X)^K.$$

We refer to $\hat{\eta}_\mu$ as the **diffraction measure** of μ .

We will see that $\hat{\eta}_\mu$ generalizes the notion of a "spectral density" of a locally square-integrable (translation-invariant) random measure on \mathbb{R}^d .

The Poisson process on X

Let $\mu = \text{Poisson}(m_X)$ (a.k.a. the "Poisson process on X ").

The Poisson process on X

Let $\mu = \text{Poisson}(m_X)$ (a.k.a. the "Poisson process on X ").

Then, for every $f \in \mathcal{L}_c^\infty(X)$, we have

$$\text{Var}_\mu(Sf) = \int_X |f(x)|^2 dm_X(x),$$

and thus $\widehat{\eta}_\mu = \sigma$ (the Plancherel measure).

DPP's on X associated with projections

Fix a Borel set $B \subset \Omega$ with $\sigma(B) = 1$, and define $\Phi(g) = \int_B \omega(g) d\sigma(\omega)$ for $g \in G$, as well as the kernel $\mathcal{K}_\Phi : X \times X \rightarrow \mathbb{C}$ by

$$\mathcal{K}_\Phi(g_1K, g_2K) = \Phi(g_1^{-1}g_2), \quad g_1K, g_2K \in G/K.$$

Since

$$\Phi * \Phi = \Phi * \Phi^* = \Phi \quad \text{and} \quad \Phi(e) = 1,$$

we see that \mathcal{K}_Φ is the kernel of a projection operator on $L^2(m_X)$.

DPP's on X associated with projections

Fix a Borel set $B \subset \Omega$ with $\sigma(B) = 1$, and define $\Phi(g) = \int_B \omega(g) d\sigma(\omega)$ for $g \in G$, as well as the kernel $\mathcal{K}_\Phi : X \times X \rightarrow \mathbb{C}$ by

$$\mathcal{K}_\Phi(g_1 K, g_2 K) = \Phi(g_1^{-1} g_2), \quad g_1 K, g_2 K \in G/K.$$

Since

$$\Phi * \Phi = \Phi * \Phi^* = \Phi \quad \text{and} \quad \Phi(e) = 1,$$

we see that \mathcal{K}_Φ is the kernel of a projection operator on $L^2(m_X)$.

Let μ denote the corresponding determinantal point process ($\iota_\mu = 1$). Then,

$$\widehat{\eta}_\mu = (1 - \kappa_\Phi) \sigma,$$

where $\kappa_\Phi(\omega) = \int_G |\Phi(g)|^2 \omega(g) dm_G(g)$.

Example I: $X = \mathbb{R}^n$, $x_0 = 0$, $G = \mathbb{R}^n$, $K = \{0\}$

Classical Fourier analysis: For $\lambda \in \mathbb{R}^n$, let $\omega_\lambda(x) = e^{2\pi i \langle \lambda, x \rangle}$, for $x \in \mathbb{R}^d$. Then,

$$\Omega = \{\omega_\lambda\}_{\lambda \in \mathbb{R}^n} \cong \mathbb{R}^n \quad \text{and} \quad \sigma = m_{\mathbb{R}^n} (= \text{Lebesgue measure}).$$

Example I: $X = \mathbb{R}^n$, $x_0 = 0$, $G = \mathbb{R}^n$, $K = \{0\}$

Classical Fourier analysis: For $\lambda \in \mathbb{R}^n$, let $\omega_\lambda(x) = e^{2\pi i \langle \lambda, x \rangle}$, for $x \in \mathbb{R}^d$. Then,

$$\Omega = \{\omega_\lambda\}_{\lambda \in \mathbb{R}^n} \cong \mathbb{R}^n \quad \text{and} \quad \sigma = m_{\mathbb{R}^n} (= \text{Lebesgue measure}).$$

Basic lemma of hyperuniformity:

$$\text{Var}_\mu(S\chi_{B_R}) = o(R^n), \quad R \rightarrow \infty \iff \hat{\eta}_\mu([- \epsilon, \epsilon]^n) = o(\epsilon^n), \quad \epsilon \rightarrow 0.$$

Example I: $X = \mathbb{R}^n$, $x_0 = 0$, $G = \mathbb{R}^n$, $K = \{0\}$

Classical Fourier analysis: For $\lambda \in \mathbb{R}^n$, let $\omega_\lambda(x) = e^{2\pi i \langle \lambda, x \rangle}$, for $x \in \mathbb{R}^d$. Then,

$$\Omega = \{\omega_\lambda\}_{\lambda \in \mathbb{R}^n} \cong \mathbb{R}^n \quad \text{and} \quad \sigma = m_{\mathbb{R}^n} (= \text{Lebesgue measure}).$$

Basic lemma of hyperuniformity:

$$\text{Var}_\mu(S\chi_{B_R}) = o(R^n), \quad R \rightarrow \infty \iff \hat{\eta}_\mu([- \epsilon, \epsilon]^n) = o(\epsilon^n), \quad \epsilon \rightarrow 0.$$

DPP (n=1): Take $B = [-1/2, 1/2]$, then $\Phi(x) = \frac{\sin(\pi x)}{\pi x}$ for $x \in \mathbb{R}$, and

$$d\hat{\eta}_\mu(\lambda) = \begin{cases} |\lambda| d\lambda & \text{for } |\lambda| \leq 1 \\ d\lambda & \text{for } |\lambda| \geq 1 \end{cases} \quad (\text{hence hyperuniform}).$$

Example II: $X = \mathbb{R}^n$, $x_0 = 0$, $G = \mathbb{R}^n \rtimes O(n)$, $K = \{0\} \rtimes O(n)$

Radial Fourier analysis: For $\lambda \in [0, \infty)$, let $\omega_\lambda(x) = a_n J_{(n-2)/2}(2\pi\lambda\|x\|)$, for $x \in \mathbb{R}^d$.
Then,

$$\Omega = \{\omega_\lambda\}_{\lambda \in [0, \infty)} \cong [0, \infty) \quad \text{and} \quad d\sigma(\lambda) = b_n \lambda^{n-1} d\lambda.$$

Example II: $X = \mathbb{R}^n$, $x_0 = 0$, $G = \mathbb{R}^n \rtimes O(n)$, $K = \{0\} \rtimes O(n)$

Radial Fourier analysis: For $\lambda \in [0, \infty)$, let $\omega_\lambda(x) = a_n J_{(n-2)/2}(2\pi\lambda\|x\|)$, for $x \in \mathbb{R}^d$.
Then,

$$\Omega = \{\omega_\lambda\}_{\lambda \in [0, \infty)} \cong [0, \infty) \quad \text{and} \quad d\sigma(\lambda) = b_n \lambda^{n-1} d\lambda.$$

Basic lemma of hyperuniformity:

$$\text{Var}_\mu(S\chi_{B_R}) = o(R^n), \quad R \rightarrow \infty \iff \hat{\eta}_\mu([0, \epsilon]) = o(\epsilon^n), \quad \epsilon \rightarrow 0.$$

Example III: $X = \mathbb{H}_{\mathbb{C}}$, $x_o = j$, $G = \mathrm{SL}_2(\mathbb{C})$, $K = \mathrm{SU}(2)$

Let

$$\mathbb{H}_{\mathbb{C}} = \{z + jw : z \in \mathbb{C}, w > 0\},$$

viewed as a subspace of Hamiltonian quaternions.

Fact: Every bi- K -invariant function on G is completely determined by its restriction to the sub-semigroup $a_t = \mathrm{Diag}(e^{t/2}, e^{-t/2})$, for $t \geq 0$.

For $\lambda \in \mathbb{R} \cup i[0, 1]$, let $\omega_{\lambda}(a_t) = \frac{\sin(\lambda t)}{\lambda \sinh(t)}$, for $t \geq 0$. Then,

$$\Omega = \{\omega_{\lambda}\}_{\lambda \in \mathbb{R} \cup i[0, 1]} \cong \mathbb{R} \cup i[0, 1] \quad \text{and} \quad d\sigma(\lambda) = \lambda^2 d\lambda, \quad \lambda \in \mathbb{R}.$$

In particular, the support of σ is a proper subset of Ω and does not contain the "identity" spherical function ($\lambda = i$).

Corollary: If $\mu = \mathrm{Poisson}(m_X)$, then $\sigma([- \epsilon, \epsilon]) = 2\epsilon^3/3$.

Is there a "basic lemma" for this Gelfand pair?

Theorem (B-Bylehn, '23) For every G -invariant locally square-integrable random measure μ on X , we have

$$\overline{\lim}_{R \rightarrow \infty} \frac{\mathrm{Var}_{\mu}(S\chi_{B_R})}{m_X(B_R)} > 0.$$

Is there a "basic lemma" for this Gelfand pair?

Theorem (B-Bylehn, '23) For every G -invariant locally square-integrable random measure μ on X , we have

$$\overline{\lim}_{R \rightarrow \infty} \frac{\text{Var}_\mu(S\chi_{B_R})}{m_X(B_R)} > 0.$$

Recall that $\omega_o(a_t) = \frac{t}{\sinh(t)}$.

Theorem (B-Bylehn, '22) Suppose that μ is a G -invariant random measure on X such that $\hat{\eta}_\mu(i[0, 1]) = 0$. Then,

$$\text{Var}_\mu(S(\omega_o\chi_{B_R})) = o(R^3), \quad R \rightarrow \infty \iff \hat{\eta}_\mu([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

Is there a "basic lemma" for this Gelfand pair?

Theorem (B-Bylehn, '23) For every G -invariant locally square-integrable random measure μ on X , we have

$$\overline{\lim}_{R \rightarrow \infty} \frac{\text{Var}_\mu(S\chi_{B_R})}{m_X(B_R)} > 0.$$

Recall that $\omega_o(a_t) = \frac{t}{\sinh(t)}$.

Theorem (B-Bylehn, '22) Suppose that μ is a G -invariant random measure on X such that $\hat{\eta}_\mu(i[0, 1]) = 0$. Then,

$$\text{Var}_\mu(S(\omega_o\chi_{B_R})) = o(R^3), \quad R \rightarrow \infty \iff \hat{\eta}_\mu([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

The condition $\hat{\eta}_\mu(i[0, 1]) = 0$ is satisfied in "most" situations.

Examples?

Tentative definition: We say that μ is *spectrally hyperuniform* if

$$\widehat{\eta}_{\mu}([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

Examples?

Tentative definition: We say that μ is *spectrally hyperuniform* if

$$\widehat{\eta}_{\mu}([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

Are there any examples?

Examples?

Tentative definition: We say that μ is *spectrally hyperuniform* if

$$\widehat{\eta}_{\mu}([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

Are there any examples? Well, some lattices in G , quasicrystals(?) (too difficult to prove).

How about DPP's?

Examples?

Tentative definition: We say that μ is *spectrally hyperuniform* if

$$\widehat{\eta}_\mu([- \epsilon, \epsilon]) = o(\epsilon^3), \quad \epsilon \rightarrow 0.$$

Are there any examples? Well, some lattices in G , quasicrystals(?) (too difficult to prove).

How about DPP's? No - for every DPP μ on X , there exists $c > 0$ such that

$$d\widehat{\eta}_\mu = \underbrace{(1 - \kappa_\phi)}_{\geq c \text{ on } \text{supp}(\sigma)} d\sigma$$

Very different from \mathbb{R}^n !

The Heisenberg nilmanifold

Heisenberg group: $H = \mathbb{C} \oplus \mathbb{R}$, $(z, s)(w, t) = (z + w, s + t + \frac{1}{2}\Im(z\overline{w}))$.

Rotations: For $k \in U(1)$, let $k.(z, s) = (kz, s)$.

Fact: $G = H \rtimes U(1)$ and $K = \{(0, 0)\} \rtimes U(1)$ form a Gelfand pair (and $G/K \cong H$).

The space of zonal spherical functions for (G, K) can be identified with the *Heisenberg fan*:

$$\Omega = \{(|\lambda|(2j+1), \lambda) : \lambda \neq 0, j \in \mathbb{N}_0\} \sqcup \{(\xi, 0) : \xi \geq 0\} \subset \mathbb{R}^2,$$

and the Plancherel measure is given by

$$\sigma(\psi) = \frac{1}{4\pi^2} \sum_{j=0}^{\infty} \int_{-\infty}^{\infty} \psi(|\lambda|(2j+1), \lambda) |\lambda| d\lambda, \quad \text{for all } \psi \in C_c^\infty(\mathbb{R}^2)$$

More precisely, $\omega_{(j,\lambda)}(z, t) = e^{i\lambda t} e^{-|\lambda||z|^2/4} L_j(|\lambda||z|^2/2)$, where L_j denotes the Laguerre polynomial of degree j .

Ongoing work (with G. Zhang)

Question: Are (isotropic) quasicrystalline point processes in the Heisenberg group "hyperuniform"?

Ongoing work (with G. Zhang)

Question: Are (isotropic) quasicrystalline point processes in the Heisenberg group "hyperuniform"?

Involves delicate estimates of Laguerre polynomials and (yet unproven) Diophantine properties of symplectic lattices. Formulas for the diffraction measures were obtained by B.-Hartnick ('18). A key difficulty is to relate the interactions between the contributions from the line $\{(\xi, 0)\}_{\xi \geq 0}$ and the support of σ .