


Hyperuniformity on the Sphere

Peter Grabner

Institute for Analysis and Number Theory
Graz University of Technology

Hyperuniform structures, rigid point processes
and related topics,
February 21, 2023

Two point distributions



Uniform distribution

Definition

A sequence of point sets $(X_N)_{N \in \mathbb{N}}$ ($X_N \subset \mathbb{S}^d$) is called uniformly distributed, if

$$\lim_{N \rightarrow \infty} \frac{\#(X_N \cap C)}{N} = \sigma_d(C),$$

for all spherical caps C .

Uniform distribution

Definition

A sequence of point sets $(X_N)_{N \in \mathbb{N}}$ ($X_N \subset \mathbb{S}^d$) is called uniformly distributed, if

$$\lim_{N \rightarrow \infty} \frac{\#(X_N \cap C)}{N} = \sigma_d(C),$$

for all spherical caps C .

Uniform distribution

Definition

A sequence of point sets $(X_N)_{N \in \mathbb{N}}$ ($X_N \subset \mathbb{S}^d$) is called uniformly distributed, if

$$\lim_{N \rightarrow \infty} \frac{\#(X_N \cap C)}{N} = \sigma_d(C),$$

for all spherical caps C .

This is equivalent to

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{\mathbf{x} \in X_N} f(\mathbf{x}) = \int_{\mathbb{S}^d} f(\mathbf{x}) d\sigma_d(\mathbf{x})$$

for all continuous (or even Riemann-integrable) functions f .

Uniform distribution

By the density of spherical harmonics in the continuous functions

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$ is equivalent to uniform distribution.

Uniform distribution

By the density of spherical harmonics in the continuous functions

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$ is equivalent to uniform distribution.

We denote by $P_n^{(d)}$ the Legendre-polynomials for \mathbb{S}^d normalised by $P_n^{(d)}(1) = 1$. These are Gegenbauer-polynomials for parameter $\lambda = \frac{d-1}{2}$ up to a scaling factor.

Hyperuniformity on the sphere

Definition (Hyperuniformity)

Let $(X_N)_{N \in \mathbb{N}}$ be a sequence of point sets on the sphere \mathbb{S}^d . The *number variance* of the sequence for caps of opening angle ϕ is given by

$$V(X_N, \phi) = \mathbb{V}_{\mathbf{x}} \# (X_N \cap C(\mathbf{x}, \phi)). \quad (1)$$

A sequence is called

Hyperuniformity on the sphere

Definition (Hyperuniformity)

Let $(X_N)_{N \in \mathbb{N}}$ be a sequence of point sets on the sphere \mathbb{S}^d . The *number variance* of the sequence for caps of opening angle ϕ is given by

$$V(X_N, \phi) = \mathbb{V}_{\mathbf{x}} \# (X_N \cap C(\mathbf{x}, \phi)). \quad (1)$$

A sequence is called

- **hyperuniform for large caps**, if

$$V(X_N, \phi) = o(N) \quad \text{as } N \rightarrow \infty \quad (2)$$

for all $\phi \in (0, \frac{\pi}{2})$;

Hyperuniformity on the sphere (continued)

Definition (continued)

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

① $\lim_{N \rightarrow \infty} \phi_N = 0$

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

- 1 $\lim_{N \rightarrow \infty} \phi_N = 0$
- 2 $\lim_{N \rightarrow \infty} N\sigma(C(\cdot, \phi_N)) = \infty.$

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

- 1 $\lim_{N \rightarrow \infty} \phi_N = 0$
- 2 $\lim_{N \rightarrow \infty} N\sigma(C(\cdot, \phi_N)) = \infty.$

- **hyperuniform for caps at threshold order**, if

$$\limsup_{N \rightarrow \infty} V(X_N, tN^{-\frac{1}{d}}) = \mathcal{O}(t^{d-1}) \quad \text{as } t \rightarrow \infty. \quad (4)$$

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

- 1 $\lim_{N \rightarrow \infty} \phi_N = 0$
- 2 $\lim_{N \rightarrow \infty} N\sigma(C(\cdot, \phi_N)) = \infty.$

- **hyperuniform for caps at threshold order**, if

$$\limsup_{N \rightarrow \infty} V(X_N, tN^{-\frac{1}{d}}) = \mathcal{O}(t^{d-1}) \quad \text{as } t \rightarrow \infty. \quad (4)$$

Hyperuniformity on the sphere (continued)

Definition (continued)

- **hyperuniform for small caps**, if

$$V(X_N, \phi_N) = o(N\sigma(C(\cdot, \phi_N))) \quad \text{as } N \rightarrow \infty \quad (3)$$

and all sequences $(\phi_N)_{N \in \mathbb{N}}$ such that

- 1 $\lim_{N \rightarrow \infty} \phi_N = 0$
- 2 $\lim_{N \rightarrow \infty} N\sigma(C(\cdot, \phi_N)) = \infty.$

- **hyperuniform for caps at threshold order**, if

$$\limsup_{N \rightarrow \infty} V(X_N, tN^{-\frac{1}{d}}) = \mathcal{O}(t^{d-1}) \quad \text{as } t \rightarrow \infty. \quad (4)$$

In all three regimes we require the point distribution to behave **better** than i.i.d. random points.

Large caps

If $(X_N)_{N \in \mathbb{N}}$ is hyperuniform for large caps, then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$. This implies uniform distribution of $(X_N)_{N \in \mathbb{N}}$.

Large caps

If $(X_N)_{N \in \mathbb{N}}$ is hyperuniform for large caps, then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$. This implies uniform distribution of $(X_N)_{N \in \mathbb{N}}$. Furthermore, it is not enough to require the defining relation for hyperuniformity for only one value of ϕ .

If $(X_N)_{N \in \mathbb{N}}$ is hyperuniform for small caps, then

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) < \infty$$

for all $n \geq 1$. This again implies uniform distribution of $(X_N)_{N \in \mathbb{N}}$.

Threshold order

If $(X_N)_{N \in \mathbb{N}}$ is hyperuniform for caps at threshold order, then

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$, which again gives uniform distribution of $(X_N)_{N \in \mathbb{N}}$.

Threshold order

If $(X_N)_{N \in \mathbb{N}}$ is hyperuniform for caps at threshold order, then

$$\lim_{N \rightarrow \infty} \frac{1}{N^2} \sum_{\mathbf{x}, \mathbf{y} \in X_N} P_n^{(d)}(\langle \mathbf{x}, \mathbf{y} \rangle) = 0$$

for all $n \geq 1$, which again gives uniform distribution of $(X_N)_{N \in \mathbb{N}}$. In the cases of small caps and caps of threshold order the conclusion of uniform distribution is not immediately obvious, since the range of caps for testing the distribution is quite restricted.

Deterministic hyperuniform point sets

- t -designs of minimal order

Deterministic hyperuniform point sets

- t -designs of minimal order
- point sets maximising

$$\sum_{\mathbf{x}, \mathbf{y} \in X} \|\mathbf{x} - \mathbf{y}\|$$

Deterministic hyperuniform point sets

- t -designs of minimal order
- point sets maximising

$$\sum_{\mathbf{x}, \mathbf{y} \in X} \|\mathbf{x} - \mathbf{y}\|$$

- sequences of QMC-designs

Deterministic hyperuniform point sets

- t -designs of minimal order
- point sets maximising

$$\sum_{\mathbf{x}, \mathbf{y} \in X} \|\mathbf{x} - \mathbf{y}\|$$

- sequences of QMC-designs
- many candidates like Fibonacci-points or spiral points, but no proofs. . .

Spherical designs

A spherical t -design is a point set $X_N \subset \mathbb{S}^d$, such that

$$\sum_{i,j=1}^N P_n^{(d)}(\langle x_i, x_j \rangle) = 0$$

for $1 \leq n \leq t$. This gives exact integration for all polynomials of degree $\leq t$ on \mathbb{S}^d .

Spherical designs

A spherical t -design is a point set $X_N \subset \mathbb{S}^d$, such that

$$\sum_{i,j=1}^N P_n^{(d)}(\langle x_i, x_j \rangle) = 0$$

for $1 \leq n \leq t$. This gives exact integration for all polynomials of degree $\leq t$ on \mathbb{S}^d .

It was a major breakthrough by Bondarenko, Radchenko, and Viazovska to show that there exist such point sets with $N = \mathcal{O}(t^d)$ points; this is the minimal possible order.

Probabilistic aspects

The original setting of hyperuniformity comes from statistical physics. The points are assumed to be sampled from a point process. The number variance is then the variance with respect to the process.

Probabilistic aspects

The original setting of hyperuniformity comes from statistical physics. The points are assumed to be sampled from a point process. The number variance is then the variance with respect to the process.

In this context the i.i.d. random case is referred to as the “Poissonian point process”. This process is – of course – not hyperuniform.

Determinantal point processes

A point process is determinantal on M with kernel $K : M \times M \rightarrow \mathbb{R}$, if its joint densities are given by

$$\rho_N(x_1, \dots, x_N) = \frac{1}{N!} \det \left(K(x_i, x_j)_{i,j=1}^N \right).$$

Determinantal point processes

A point process is determinantal on M with kernel $K : M \times M \rightarrow \mathbb{R}$, if its joint densities are given by

$$\rho_N(x_1, \dots, x_N) = \frac{1}{N!} \det \left(K(x_i, x_j)_{i,j=1}^N \right).$$

This notion was originally developed in physics, where the joint wave function of N fermionic particles can be expressed as a determinant of the above form.

Determinantal point processes

A point process is determinantal on M with kernel $K : M \times M \rightarrow \mathbb{R}$, if its joint densities are given by

$$\rho_N(x_1, \dots, x_N) = \frac{1}{N!} \det \left(K(x_i, x_j)_{i,j=1}^N \right).$$

This notion was originally developed in physics, where the joint wave function of N fermionic particles can be expressed as a determinant of the above form.

The fact that the determinant vanishes, if $x_i = x_j$ for some $i \neq j$, implies a mutual repulsion of the sample points (“particles”).

Determinantal point processes

The eigenvalues of random matrices, as well as the roots of random polynomials can also be modelled by determinantal point processes.

Determinantal point processes

The eigenvalues of random matrices, as well as the roots of random polynomials can also be modelled by determinantal point processes.

One special case is especially important and easy to understand:

Determinantal point processes

The eigenvalues of random matrices, as well as the roots of random polynomials can also be modelled by determinantal point processes.

One special case is especially important and easy to understand:

Let $H \subset L^2(M)$ be a finite dimensional space and K_H be the orthogonal projection to this space. If $N = \dim H$, then the DPP given by the kernel K_H samples exactly N points.

The spherical ensemble on \mathbb{S}^2

The kernel

$$\tilde{K}^{(N)}(x, y) = \frac{N(1 + x\bar{y})^{N-1}}{4\pi(1 + |x|^2)^{N+1}(1 + |y|^2)^{N+1}}$$

on \mathbb{C}^2 describes the distribution of the eigenvalues of AB^{-1} for two $N \times N$ matrices A, B with i.i.d. complex Gaussian entries.

The spherical ensemble on \mathbb{S}^2

The kernel

$$\tilde{K}^{(N)}(x, y) = \frac{N(1 + x\bar{y})^{N-1}}{4\pi(1 + |x|^2)^{N+1}(1 + |y|^2)^{N+1}}$$

on \mathbb{C}^2 describes the distribution of the eigenvalues of AB^{-1} for two $N \times N$ matrices A, B with i.i.d. complex Gaussian entries. Stereographically projecting the eigenvalues to the sphere \mathbb{S}^2 gives a point process; the spherical ensemble. Its joint densities are given by

$$C_N \prod_{1 \leq i < j \leq N} \|\mathbf{x}_i - \mathbf{x}_j\|^2$$

with a normalising constant C_N .

The spherical ensemble on \mathbb{S}^2

The kernel

$$\tilde{K}^{(N)}(x, y) = \frac{N(1 + x\bar{y})^{N-1}}{4\pi(1 + |x|^2)^{N+1}(1 + |y|^2)^{N+1}}$$


on \mathbb{C}^2 describes the distribution of the eigenvalues of AB^{-1} for two $N \times N$ matrices A, B with i.i.d. complex Gaussian entries. Stereographically projecting the eigenvalues to the sphere \mathbb{S}^2 gives a point process; the spherical ensemble. Its joint densities are given by

$$C_N \prod_{1 \leq i < j \leq N} \|\mathbf{x}_i - \mathbf{x}_j\|^2$$

with a normalising constant C_N .

It has been shown by Alishahi and Zamani that samples of the spherical ensemble are hyperuniform for small caps.

Sample of spherical ensemble



The harmonic ensemble on \mathbb{S}^d

Let H_L be the span of all spherical harmonics of degree $\leq L$ on \mathbb{S}^d . Then the corresponding projection kernel defines a determinantal point process sampling $\dim H \sim L^d$ points. This process was introduced and studied by Beltrán, Marzo, and Ortega-Cerdá.

The harmonic ensemble on \mathbb{S}^d

Let H_L be the span of all spherical harmonics of degree $\leq L$ on \mathbb{S}^d . Then the corresponding projection kernel defines a determinantal point process sampling $\dim H \sim L^d$ points. This process was introduced and studied by Beltrán, Marzo, and Ortega-Cerdá.

They proved *inter alia* that samples of this process are hyperuniform for small caps.

The harmonic ensemble on \mathbb{S}^d

Let H_L be the span of all spherical harmonics of degree $\leq L$ on \mathbb{S}^d . Then the corresponding projection kernel defines a determinantal point process sampling $\dim H \sim L^d$ points. This process was introduced and studied by Beltrán, Marzo, and Ortega-Cerdá.

They proved *inter alia* that samples of this process are hyperuniform for small caps.

In joint work with J. Brauchart, W. Kusner, and J. Ziefle we could prove that samples of this process are also hyperuniform at threshold order (in a slightly weaker sense).

Jittered sampling

Let A_i ($i = 1, \dots, N$) be an area regular partition of \mathbb{S}^d with $\text{diam}(A_i) \leq CN^{-\frac{1}{d}}$ and $\sigma(A_i) = \frac{1}{N}$. Such partitions exist by work of Kuijlaars and Saff and Gigante and Leopardi.

Jittered sampling

Let A_i ($i = 1, \dots, N$) be an area regular partition of \mathbb{S}^d with $\text{diam}(A_i) \leq CN^{-\frac{1}{d}}$ and $\sigma(A_i) = \frac{1}{N}$. Such partitions exist by work of Kuijlaars and Saff and Gigante and Leopardi.

Then define point process by taking N points independently uniformly from the sets A_i (one point per set). This process is the determinantal process given by the projection to the space of functions measurable with respect to the finite σ -algebra generated by $(A_i)_{i=1}^N$.

Jittered sampling

Let A_i ($i = 1, \dots, N$) be an area regular partition of \mathbb{S}^d with $\text{diam}(A_i) \leq CN^{-\frac{1}{d}}$ and $\sigma(A_i) = \frac{1}{N}$. Such partitions exist by work of Kuijlaars and Saff and Gigante and Leopardi.

Then define point process by taking N points independently uniformly from the sets A_i (one point per set). This process is the determinantal process given by the projection to the space of functions measurable with respect to the finite σ -algebra generated by $(A_i)_{i=1}^N$.

Jittered sampling points are hyperuniform in all three regimes.

Poissonian pair correlation

A second local statistic of point distributions has been studied in the context of eigenvalue distributions of Schrödinger operators in the physics context. This has then been extended to zeros of the Riemann zeta function and sequences of the form $(\alpha p(n))$, where p is a polynomial of degree ≥ 2 .

Poissonian pair correlation

A second local statistic of point distributions has been studied in the context of eigenvalue distributions of Schrödinger operators in the physics context. This has then been extended to zeros of the Riemann zeta function and sequences of the form $(\alpha p(n))$, where p is a polynomial of degree ≥ 2 .

Definition

A sequence of points $(x_n)_n$ on the unit circle has **Poissonian pair correlation**, if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \# \{1 \leq i \neq j \leq N \mid N|x_i - x_j| < s\} = 2s$$

for all $s > 0$.

This is a property shared by i.i.d. random point sets.

Poissonian pair correlation continued

It has been proved by Grepstad and Larcher and independently Aistleitner, Lachmann, and Pausinger that PPC implies uniform distribution. The proofs of this expected fact are surprisingly complicated and intricate.

Higher dimensions

In higher dimensions the obvious generalisation is.

Definition

A sequence of point sets $(X_N)_N$ on a compact manifold M of dimension n has Poissonian pair correlation, if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \# \left\{ 1 \leq i \neq j \leq N \mid N^{1/n} d(x_i^{(N)}, x_j^{(N)}) < s \right\} = s^n B_n$$

for all $s > 0$; B_n denotes the volume of the n -dimensional unit ball. Here, d denotes the geodesic distance on M .

PPC implies uniform distribution

Steinerberger studied the case of the torus and gave a proof that PPC implies uniform distribution in this case. Marklof gave a very general proof of this fact for general compact manifolds.

Open questions

- Find relations with other measures of uniformity: discrepancy, error of integration, energy. . .

Open questions

- Find relations with other measures of uniformity: discrepancy, error of integration, energy. . .
- Find more explicit deterministic constructions for hyperuniform point sets for any N .

Open questions

- Find relations with other measures of uniformity: discrepancy, error of integration, energy. . .
- Find more explicit deterministic constructions for hyperuniform point sets for any N .
- Find explicit deterministic constructions for point sets achieving the best possible discrepancy bound (or even a bound better than $N^{-\frac{1}{2}}$)

Thank you for your
attention!