

name: <unnamed> /msu/scratch4/m1cmb07/Connor\_bob/mmb/output/stepwise\_regressions/infl\_per log: rr\_stky\_pr\_other.smcl log type: smcl opened on: 18 Jul 2024, 15:09:53 Interaction effects of stky\_pr\_other and rules on infl\_per\_rr at various horizons note: 1.rule\_g omitted because of collinearity. note: 1.stky\_pr\_other#1.rule\_g omitted because of collinearity.
obtaining LAD starting values ... done M regression (95% efficiency) Number of obs 228 Wald chi2(5) = 75.88 Prob > chi2 = 0.0000 Pseudo R2 = 0.1378 Biweight k 4.685 .25461069 Scale Robust Coefficient std. err. infl\_per\_rr20 t P>|t| [95% conf. interva > 1] stky\_pr\_other 0 (empty) 0 -.0679052 .1119388 -0.61 0.545 -.2885037 .15269 1 > 33 rule tr (empty) 0 0 1 .4312272 .0591256 7.29 0.000 .314708 .54774 > 64 rule\_itr 0 (empty) 0 1 .2690239 .0673714 3.99 0.000 .1362545 .40179 > 32 rule\_g 0 0 (empty) 0 (empty) stky\_pr\_other#rule\_tr 0 (empty) 0 0 0 1 0 (empty) 1 0 0 (empty) -.0183447 .1415603 0.897 1 1 -0.13 -.2973186 . 26062 > 92 stky\_pr\_other#rule\_itr 0 0 0 (empty) (empty) 0 1 0 1 0 0 (empty) 1 1 -.1224913 .1428104 -0.86 0.392 -.4039289 .15894 > 63 stky\_pr\_other#rule\_g 0 0 0 (empty) 0 (empty) 0 1 1 0 0 (empty) 1 1 0 (empty) \_cons 0.000 -.5970891 -.4851886 .0567819 -8.54 -.37328 > 81

Thursday July 18 15:09:53 2024 Page 2

\_cons

> 47

-.5175684

.0579049

-8.94

0.000

-.6316822

- . 40345

\* Interaction effects of stky\_pr\_other and rules on infl\_per\_rr at various horizons note: 1.rule\_g omitted because of collinearity. note: 1.stky\_pr\_other#1.rule\_g omitted because of collinearity. obtaining LAD starting values ... done iterating RLS ..... done iterating RLS ......done fitting empty model ... done computing standard errors ... done M regression (95% efficiency) Number of obs 228 Wald chi2(5) 87.23 Prob > chi2 0.0000 Pseudo R2 = 0.1250 Biweight k = 4.685 Scale .29230746 Robust Coefficient std. err. infl\_per\_rr40 t P>|t| [95% conf. interva > 1] stky\_pr\_other 0 0 (empty) 0.457 1 -.0813723 .109205 -0.75 -.2965833 .13383 > 87 rule\_tr 0 0 (empty) 1 .4721758 .0614037 7.69 0.000 .3511671 . 59318 > 45 rule\_itr 0 (empty) 0 .3042426 .073309 1 4.15 0.000 .159772 .44871 > 32 rule\_g 0 0 (empty) 0 (empty) stky\_pr\_other#rule\_tr 0 0 0 (empty) 0 1 (empty) 0 1 0 0 (empty 1 1 -.0362759 .1451583 -0.25 0.803 -.3223405 .24978 > 87 stky\_pr\_other#rule\_itr 0 0 0 (empty) 0 1 0 (emptý) 1 0 0 (empty 1 1 -.16657 .1615577 -1.03 0.304 - . 484953 .1518 > 13 stky\_pr\_other#rule\_g 0 0 0 (empty) 0 1 0 (empty) 1 0 0 (empty) (empty) 1 1 0

\* Interaction effects of stky\_pr\_other and rules on infl\_per\_rr at various horizons note:  $1.rule_g$  omitted because of collinearity. note: 1.stky\_pr\_other#1.rule\_g omitted because of collinearity. obtaining LAD starting values ... done iterating RLS ..... do iterating RLS  $\ldots\ldots$  done fitting empty model  $\ldots$  done computing standard errors ... done M regression (95% efficiency) Number of obs 228 Wald chi2(5) 79.11 Prob > chi2 0.0000 Pseudo R2 = 0.1178 Biweight k = 4.685 Scale .29055062 Robust Coefficient std. err. infl\_per\_rr60 t P>|t| [95% conf. interva > 1] stky\_pr\_other 0 0 (empty) -.0759242 1 .114779 -0.66 0.509 -.3021199 . 15027 > 16 rule\_tr 0 0 (empty) 1 .4826621 .0672081 7.18 0.000 .3502146 . 61510 > 96 rule\_itr 0 (empty) 0 .3054412 .0791456 1 3.86 0.000 .1494684 .4614 > 14 rule\_g 0 0 (empty) 0 (empty) stky\_pr\_other#rule\_tr 0 0 0 (empty) (empty) 0 1 0 1 0 0 (empty) 1 1 -.0425378 .149296 -0.28 0.776 -.3367566 .25168 > 09 stky\_pr\_other#rule\_itr 0 0 0 (empty) 0 1 0 (emptý) 1 0 0 (empty 1 1 -.1672533 .1774829 -0.94 0.347 -.5170202 .18251 > 35 stky\_pr\_other#rule\_g 0 0 0 (empty) 0 1 0 (empty) 1 0 0 (empty)

(empty)

.063839

-8.28

0.000

-.6542219

-.4026

0

-.5284139

1 1

\_cons

> 06

name: <unnamed>

log: /msu/scratch4/m1cmb07/Connor\_bob/mmb/output/stepwise\_regressions/infl\_per
> \_rr\_stky\_pr\_other.smcl
log type: smcl
closed on: 18 Jul 2024, 15:09:53