

Exposure (Taxonomy)	OR (95% CI)	Effect Size (β)	Standard Error (SE)	P-value	nSNP	Cochran's Q	Q P-value
class Actinobacteria	1.34 (1.08–1.65)	0.2909	0.1072	0.0067	15	30.35	0.0068
order Enterobacteriales	1.76 (1.16–2.67)	0.5661	0.2119	0.0076	7	20.99	0.0018
family Enterobacteriaceae	1.76 (1.16–2.67)	0.5661	0.2119	0.0076	7	20.99	0.0018
order Methanobacteriales	1.13 (1.03–1.25)	0.1262	0.0492	0.0103	10	4.99	0.8355
class Methanobacteria	1.13 (1.03–1.25)	0.1262	0.0492	0.0103	10	4.99	0.8355
family Methanobacteriaceae	1.13 (1.03–1.25)	0.1262	0.0492	0.0103	10	4.99	0.8355
phylum Bacteroidetes	0.77 (0.63–0.94)	-0.2614	0.1027	0.0109	10	4.82	0.8500
family Bifidobacteriaceae	1.38 (1.07–1.78)	0.3251	0.1292	0.0119	13	34.93	4.80e-04
order Bifidobacteriales	1.38 (1.07–1.78)	0.3251	0.1292	0.0119	13	34.93	4.80e-04
genus Ruminococcus2	0.84 (0.73–0.96)	-0.1732	0.0695	0.0127	15	10.29	0.7405
class Coriobacteriia	1.25 (1.04–1.49)	0.2196	0.0916	0.0165	16	18.61	0.2321
order Coriobacteriales	1.25 (1.04–1.49)	0.2196	0.0916	0.0165	16	18.61	0.2321
family Coriobacteriaceae	1.25 (1.04–1.49)	0.2196	0.0916	0.0165	16	18.61	0.2321
family Acidaminococcaceae	0.81 (0.67–0.96)	-0.2167	0.0919	0.0184	7	4.66	0.5875
genus FamilyXIIIAD3011group	0.79 (0.65–0.96)	-0.2363	0.1017	0.0202	13	18.42	0.1036
order Bacteroidales	0.81 (0.67–0.97)	-0.2144	0.0965	0.0262	12	9.01	0.6211
class Bacteroidia	0.81 (0.67–0.97)	-0.2144	0.0965	0.0262	12	9.01	0.6211
genus RuminococcaceaeUCG010	1.26 (1.01–1.58)	0.2339	0.1151	0.0422	6	5.97	0.3087
genus Roseburia	0.80 (0.65–1.00)	-0.2184	0.1106	0.0484	14	20.36	0.0866