

Exploring the causal relationship between gut microbiota, circulating inflammatory proteins, and colorectal neuroendocrine tumors: A Mendelian randomization study

Jie Cui, Tie-Jun Wang, Yue-Chen Zhao, Xuan-Peng Zhou*

* Corresponding author:

Xuan-Peng Zhou

Department of Gastrointestinal and Colorectal Surgery, China-Japan Union Hospital of Jilin University, Changchun, Jilin, China.

E-mail: xpzhou22@mails.jlu.edu.cn.

Journal name: *discover oncology*

Supplementary Figures

Supplementary Fig. 1 Scatter plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors

Supplementary Fig. 2 Funnel plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors

Supplementary Fig. 3 Forest plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors

Supplementary Fig. 4 Leave-one-out sensitivity analysis in the MR for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors


Supplementary Fig. 5 Scatter plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors

Supplementary Fig. 6 Funnel plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors


Supplementary Fig. 7 Forest plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif

chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors


Supplementary Fig. 8 Leave-one-out sensitivity analysis in the MR for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors




Supplementary Fig. 1 Scatter plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors




Supplementary Fig. 2 Funnel plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors




Supplementary Fig. 3 Forest plot of the MR analysis for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors




Supplementary Fig. 4 Leave-one-out sensitivity analysis in the MR for the causal association between five gut microbiota and CR-NETs. **(A)**. family Porphyromonadaceae and CR-NETs. **(B)**. genus Oscillospira and CR-NETs. **(C)**. genus Paraprevotella and CR-NETs. **(D)**. family Porphyromonadaceae and CR-NETs. **(E)**. genus Ruminococcaceae NK4A214 group and CR-NETs. CR-NETs: colorectal neuroendocrine tumors




Supplementary Fig. 5 Scatter plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors



Supplementary Fig. 6 Funnel plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors



Supplementary Fig. 7 Forest plot of the MR analysis for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors



Supplementary Fig. 8 Leave-one-out sensitivity analysis in the MR for the causal association between five inflammatory proteins and CR-NETs. **(A)**. C-C motif chemokine 4 and CR-NETs. **(B)**. C-C motif chemokine 23 and Cr-NETs. **(C)**. TNF-related apoptosis-inducing ligand and CR-NETs. **(D)**. C-X-C motif chemokine 5 and CR-NETs. **(E)**. Monocyte chemoattractant protein 2 and CR-NETs. CR-NETs: colorectal neuroendocrine tumors