

Cayley's Ω -Process And The Reynolds Operator

Bert Lorke

February 26, 2018

Contents

1	Introduction	1
2	Preliminary Work	3
2.1	Concepts From Algebraic Geometry	3
2.2	Concepts From Invariant Theory	4
3	Linearly Reductive Groups, The Reynolds Operator And Hilbert's Finiteness Theorem	6
3.1	The Reynolds Operator And Linearly Reductive Groups	6
3.2	Hilbert's Finiteness Theorem	9
3.3	The Reynolds Operator Of A Linear Algebraic Group	11
4	Cayley's Ω-Process	12
5	Examples (not a section in the final version)	16
6	Further Discussion	20

1 Introduction

A very important concept in mathematics is the idea of an *invariant*: An object which does not change under a certain action. In 1872, Felix Klein came up with a then new method of describing geometries with group theory, called the Klein Erlangen program. Here, the central idea of a geometry is characterized by its associated symmetry group, the group of transformations which leaves certain objects unchanged, for example: angles. The study of these transformations is called conformal geometry.

Let us discuss the following important example in projective geometry: Consider all transformations which map lines to lines, id est, such transformations under which the the property of being a line is invariant. In real projective geometry, the fundamental theorem of projective geometry gives us that these maps are exactly the projective transformations.

Conversely, we can now just consider projective transformations as our given group of transformations. **Invariant theory asks: What invariants exist?** We can loosely notice a kind of duality between geometries viewed as in the Klein Erlangen program and invariant theory. This discipline of mathematics usually only looks at invariants described with so called regular terms, or more concretely formulated: In invariant theory, we try to find invariant polynomial-like functions.

Staying in our example of considering projective transformations as our given group, a well known example for an invariant is the cross ratio. It is a rational polynomial which takes as its input four collinear points. Is this the only invariant? How can we find other invariants? How big is the set (this will be a ring) of all invariants?

Hilbert's finiteness theorem states that for regular actions under certain groups, such that are *linearly reductive*, the invariant ring is finitely generated. If we can find these finite generators, we have a grasp of what all invariants look like. Hilbert's first proof for this theorem was non-constructive. It is claimed¹ that this proof was responsible for Gordan's famous quote "Das ist Theologie und nicht Mathematik". The central idea of this proof is the existence of a Reynolds operator.

One of the most important and most common groups is the general linear group GL_n . It would be great if this group were linearly reductive. But it is! There are multiple ways to see this. In a seminar I held, with the help of the Haar measure, I discussed a way to see that a module complement exists for every representation, making GL_n linearly reductive. One can also show linear reductivity by the Schur-Weyl-duality: The symmetric group is finite, and therefore we can see that rational GL_n representations are semisimple, from which we can again construct module complements.

Here, we will show that GL_n is linear reductive in an even different way. For one, we want to show that a Reynolds Operator exists, which already means that GL_n is linearly reductive. But we want even more than just the existence! What does it help for our motivation to get a grasp of what all (or even just some) invariants look like, if we merely prove the existence of a finite generator set for the invariants? Since this operator projects polynomials to invariant polynomials, if we can find an explicit formula for computing the Reynolds operator applied to a polynomial, we can more easily receive concrete invariants. **This is possible with Cayley's Ω -process!** This is the main focus of my work.

I say "more easily" receive invariants, because if we take a polynomial at random and apply the Reynolds Operator, we might very likely just get a constant polynomial, which is not a very interesting invariant, and we also want to know if there are more invariants. Similar to the first proof of Hilbert's finiteness theorem (by Hilbert himself), we can show that there are certain finitely many polynomials whose images under the Reynolds operator will generate the invariant ring. Although this is not what I will be discussing in detail in my work,

¹I read somewhere that it is not certain

there is in fact an algorithm to compute these certain polynomials. With the help of Cayley's Ω -process, we then get a complete algorithm that gives us the generators of the invariant ring.

2 Preliminary Work

2.1 Concepts From Algebraic Geometry

In the following, K is a field of characteristic 0 and G a linear algebraic group, that is a group whose set is an affine variety, and whose multiplication and inversion are morphisms of affine varieties.

Proposition 2.1

let $V = K^n$ for some $n \in \mathbb{N}$. For a polynomial $p \in K[V] = K[\{X_i\}_{i \in [n]}]$, the set $X_p := \{v \in V \mid p(v) \neq 0\}$ is an affine variety with the coordinate ring $K[X_p] = K[\{X_i\}_{i \in [n]}, p^{-1}]$.

Proof. The set X_p is not an algebraic set itself. The trick is to consider X_p as a subset of $K \times V$. We do this as follows: Consider $\tilde{X}_p := Z(X_0 \cdot p - 1) \subseteq K \times V$. We see that X_p corresponds to \tilde{X}_p by noticing that $\tilde{X}_p = \{(1/p(v), v) \in K \times V \mid v \in X_p\}$, which therefore means we have the one-to-one correspondence $(1/p(v), v) \leftrightarrow v$. By definition \tilde{X}_p is an affine variety, and it is also easy to see that $K[X_p] \cong K[\tilde{X}_p] = K[K \times V]/(X_0 \cdot p - 1)$ via $X_0 \leftrightarrow 1/p$, including evaluations with the above described correspondence of points, which is easy to check. \square

Example 2.2: The General Linear Group GL_n

One of the most important examples is the general linear group GL_n , which will be an essential theme in my work. By the above proposition this group is an affine variety via $p = \det$ with the coordinate ring $K[\{X_{i,j}\}_{i,j \in [n]}, \det]$. This makes GL_n into a *linear algebraic group*, that is a group which is an affine variety whose group operations of the multiplication and inversion are morphisms of affine varieties.

Definition 2.3: Algebraic Cohomomorphism

Let $m: U_1 \times U_2 \rightarrow W$ be a morphism of affine varieties. The strict pullback, which we shall call \hat{m} , is a map of the type $K[W] \rightarrow K[U_1 \times U_2]$. We have $K[U_1 \times U_2] = K[\{X_k\}_{k \in [r]}, \{Y_l\}_{l \in [s]}]$, where $\{X_k\}_{k \in [r]}$ and $\{Y_l\}_{l \in [s]}$ are generators of $K[U_1]$ and $K[U_2]$ respectively. We define the following map:

$$t: K[U_1 \times U_2] \rightarrow K[U_1] \otimes K[U_2]$$

$$\sum_i \lambda_i \prod_j X_j^{d_{i,j}} \prod_k Y_k^{e_{i,k}} \mapsto \sum_i \lambda_i \prod_j X_j^{d_{i,j}} \otimes \prod_k Y_k^{e_{i,k}} \quad (1)$$

This is independent of the choice of generators and independent of the representatives and therefore well-defined. It is even an isomorphism. Now, finally,

we define $m^* := t \circ \hat{m} : K[W] \longrightarrow K[U_1] \otimes K[U_2]$. We call m^* the *algebraic cohomomorphism of m* .

Remark 2.3.1

One might ask why we want to look at these objects $m^*(f)$ instead of $\hat{m}(f)$. Really, these objects are hardly different (the spaces are isomorphic), but it helps to formalize performing operations only on the “left part” or the “right part”, as we will soon see. This is an approach that [DK15] follows, but other literature such as [Stu08] (and probably also Cayley) rather consider $\hat{m}(f)$ written as $\hat{m}(f) = f(XY)$. To give a very simple example: If $f \in K[Z]_G$, we will write $\text{id} \otimes \frac{\partial}{\partial Z_i}(m^*f)$ as in [DK15], whereas [Stu08] would write $\frac{\partial}{\partial Y_i}f(XY)$.

2.2 Concepts From Invariant Theory

Definition 2.4: Regular Action, Finite Rational Representation

Let G be a linear algebraic group, X an affine variety. We call an action $G \times X \longrightarrow X$ a **regular action**, iff μ is a morphism of affine varieties. We say G **acts regularly on X** , and we also call X a **G -variety**.

For a finite-dimensional vector space V , let $\mu : G \times V \longrightarrow V$ be a representation in the classical sense, that is for all $g \in G$ we have $D_\mu(g) := (v \mapsto \mu(g, v)) \in \text{GL}(V)$. We call μ a *finite rational representation* if and only if it is regular.

Definition 2.5: Rational Representation

Let V be a vector space (not necessarily finite dimensional), and $\mu : G \times V \longrightarrow V$ an action. We call μ a **rational representation** iff there exists a linear map $\mu' : V \longrightarrow K[G] \otimes V$ such that $\mu(\sigma, v) = ((\epsilon_\sigma \otimes \text{id}) \circ \mu')(v)$.

Definition 2.6

Let $\mu : G \times X \longrightarrow X$ be a regular action. We define an action $\bar{\mu} : G \times K[X] \longrightarrow K[X]$ via $\bar{\mu}(\sigma, f)(x) := f(\mu(\sigma^{-1}, x))$, or $\sigma.f(x) := f(\sigma^{-1}.x)$, where $\sigma \in G$, $f \in K[X]$ and $x \in X$. This action is obviously regular, but it is also easily shown that it is in fact a rational representation: If $\tilde{\mu}$ is the algebraic map (it is in fact a left action) defined by $\tilde{\mu}(\sigma, x) := \mu(\sigma^{-1}, x)$, ($\sigma \in G$, $x \in X$), then we can define $\bar{\mu}' := \tilde{\mu}^*$ with the desired properties.

Proposition 2.7

If we have $\bar{\mu}'(f) = \sum_i p_i \otimes g_i$, then for $\sigma \in G$ we get $\bar{\mu}'(\sigma.f) = \sum_i p_i \otimes \sigma.g_i$.

Definition 2.8: locally finite

For a vector-space V , we call an action $\mu : G \times V \longrightarrow V$ **locally finite**, iff for every $v \in V$ there exists a G -stable finite-dimensional vector space $U \subseteq V$ such that $v \in U$.

Definition 2.9

Let V be a vector-space and $\mu : G \times V \longrightarrow V$ an action. For $v \in V$ we define $V_v := \text{span } G.v$.

Remark 2.9.1

V_v is always a G -stable subspace of V . For any G -stable subspace $W \subseteq V$ we

have $V_v \subseteq W$. Therefore, an action $\mu: G \times V \rightarrow V$ is locally finite if and only if V_v is finite-dimensional.

Proposition 2.10

An action $\mu: G \times V \rightarrow V$ is a rational representation if and only if the action is locally finite, and every finite-dimensional G -stable subspace W , $\mu|_{G \times W}$ is a finite rational representation.

Proof. See [DK15, A.1.8] and [DK15, 2.2.5(b) \implies (c), 2.2.6]

Assume that μ is a rational representation. Let $v \in V$. We can write $\mu'(v) = \sum_{i=1}^l f_i \otimes v_i$. We then easily see that $V_v \subseteq \text{span}\{v_i\}_{i=1}^l$, showing that the action is locally finite. Now let W be a finite-dimensional G -stable subspace of V with the basis $\{w_i\}_{i=1}^r$. We have $p_{i,j} \in K[G]$ with $\mu'(w_j) = \sum_{i=1}^r p_{i,j} \otimes w_i$. Now let $w = \sum_{j=1}^r \lambda_j w_j \in W$. Then for all $\sigma \in G$ we have

$$\begin{aligned} \mu(\sigma, w) &= ((\epsilon_\sigma \otimes \text{id}) \circ \mu')(w) \\ &= \sum_{j=1}^r \lambda_j \sum_{i=1}^r p_{i,j}(\sigma) \cdot w_i \end{aligned} \tag{2}$$

from which we immediately notice that $D_{\mu|_{G \times W}}(\sigma) \in \text{GL}(W)$. Therefore $\mu|_{G \times W}$ is a finite rational representation.

Now let μ be a locally finite action such that for every finite-dimensional G -stable subspace W , $\mu|_{G \times W}$ is a finite rational representation. Let $v \in V$. By assumption there exists a finite-dimensional G -stable vector space $W \subseteq V$ such that $v \in W$. Choose a basis $\{w_1, \dots, w_r\}$ of W , where $w_1 = v$ (this is done to make some terms easier, but choosing any basis also works). Since $\mu|_{G \times W}$ is a regular representation, we have $D_\mu(\sigma) \in \text{GL}(W)$ for all $\sigma \in G$, and since our action is also regular, for $i \in [r]$ there exist $p_i \in K[G]$ such that for all $\sigma \in G$ we have $\mu(\sigma, v) = D_\mu(\sigma)(v) = \sum_{i=1}^r p_i(\sigma) \cdot w_i$. Now define $\mu'(v) := \sum_{i=1}^r p_i \otimes w_i$. We shall now check that this is well-defined:

PROOF NOT DONE YET

□

Remark 2.10.1

This shows that for a finite-dimensional vector space V , an action is a rational representation if and only if it is a finite rational representation. In other words, we have shown that finite rational representations are exactly the rational representations that are finite dimensional, which justifies the choice of the names of our definitions.

Remark 2.10.2

A finite rational representation $\mu: G \times V \rightarrow V$ is of the following form: Consider $D_\mu: G \rightarrow \text{GL}(V)$. If then $a_{i,j}: G \rightarrow K$ is the function of the (i, j) -entry of D_μ , then $a_{i,j} \in K[G]$.

In fact, it is equivalent to define a representation $\mu: G \times V \rightarrow V$ (V finite dimensional) as rational, iff $D_\mu: G \rightarrow \text{GL}(V)$ is a map of affine varieties.

Definition 2.11: Invariants

Let G act on X regularly.

$$X^G := \{x \in X \mid \forall g \in G : g.x = x\} \quad (3)$$

This defines a linear subspace. The given action induces an action $\bar{\mu}: G \times K[X] \rightarrow K[X]$ as per definition 2.6. The **invariant ring** of the representation is defined as

$$K[X]^G := \{f \in K[X] \mid \forall g \in G : g.f = f\} \quad (4)$$

As the name implies, $K[X]^G$ defines a subalgebra of $K[X]^G$.

The general theme of my work revolves around the question of whether the invariant ring $K[X]^G$ is finitely generated.

Hilbert's finiteness theorem states that if the group G is linearly reductive, $K[V]^G$ is finitely generated. The strict definition of “linearly reductive” is quite tricky, but we will give an alternate equivalent definition shortly.

3 Linearly Reductive Groups, The Reynolds Operator And Hilbert's Finiteness Theorem

3.1 The Reynolds Operator And Linearly Reductive Groups

Definition 3.1: Linearly Reductive Group

Let G be a linear algebraic group. We call G **linearly reductive**, if and only if for any rational representation V and for any $v \in V^G \setminus \{0\}$ there exists an $f \in (V^*)^G$ such that $f(v) \neq 0$.

Proposition 3.2

For a linearly reductive group G and a finite rational representation V , V^G and $(V^*)^G$ are dual to each other with respect to the non-degenerate bilinear form $b: V^* \times V \rightarrow K, (f, v) \mapsto f(v)$.

Proof. We shall first show that $\dim(V^*)^G = \dim V^G$. Let $\{v_1, \dots, v_r\}$ be a basis of V^G , and expand this to a basis $\{v_1, \dots, v_m\}$ of V . We define $\{f_1, \dots, f_r\}$ to be the basis of V^* which is dual to $\{v_1, \dots, v_r\}$, that is we have $f_i(v_j) = \delta_{i,j}$. It should be clear that $\{f_1, \dots, f_r\} \subseteq (V^*)^G$. Now let $f \notin \text{span}\{f_1, \dots, f_r\}$, that is we have $f = \sum_{i=1}^n \lambda_i f_i$, and there exists a $j > r$ such that $\lambda_j \neq 0$. For this j , we have $v_j \notin V^G$, therefore there exists a $\sigma \in G$ such that $\sigma.v_j \neq v_j$. We then get ?? We have now shown that $(V^*)^G = \text{span}\{f_1, \dots, f_r\}$, therefore $\dim(V^*)^G = \dim V^G$. Since G is linearly reductive, we then get via our definition that $b|_{(V^*)^G \times V^G}$ is non-degenerate in the first variable. Since $\dim(V^*)^G = \dim V^G$, we have that $b|_{(V^*)^G \times V^G}$ is non-degenerate in both variables. This exactly means that the spaces $(V^*)^G$ and V^G are dual to each other with respect to b . \square

Definition 3.3: Reynolds Operator

Let X be an affine G -variety. A G -invariant linear projection $R: K[X] \rightarrow K[X]^G$ is called a **Reynolds operator**.

Definition 3.4

Assume that V is a finite rational representation of V such that there exists a unique subrepresentation of V such that $V = V^G \oplus W$. We define $R_V: V \rightarrow V^G$ as the linear projection of V onto V^G along W .

Remark 3.4.1

R_V is a G -invariant projection of V onto V^G : If for $v \in V$ we write $v = u + w$ with $u \in V^G$ and $w \in W$, then for $\sigma \in G$ we have $\sigma.v = \sigma.u + \sigma.w = u + \sigma.w$, and therefore $R_V(\sigma.v) = u = R_V(v)$.

Lemma 3.5

Assume that G is a linear algebraic group with the following property: For every finite rational representation V of G there exists a unique subrepresentation W of V such that $V = V^G \oplus W$, and for this W we have $(W^*)^G = \{0\}$. The following properties hold:

- (a) If V is a subrepresentation of a finite rational representation V' of G , we have $R_{V'}|_V = R_V$.
- (b) If V is a finite rational representation of G and $R'_V: V \rightarrow Y$ is a G -invariant linear map with $V \subseteq Y$ and $R'_V|_{V^G} = \text{id}_{V^G}$, we have $R'_V = R_V$, id est R_V is unique with this property (**Do I need to mention that we should then view $R_V: V \rightarrow V$ instead of $\rightarrow V^G$??**).
- (c) If X is an affine G -variety and $R: K[X] \rightarrow K[X]^G$ is a Reynolds operator, then for every G -stable subspace V of $K[X]$ we have $R|_V = R_V$.
- (d) If X is an affine G -variety, $R: K[X] \rightarrow K[X]^G$ a Reynolds operator and W is any G -stable subspace of $K[X]$, we have $R(W) = W^G$.
- (e) If X is an affine G -variety, the Reynolds operator is unique

Proof.

(a)

Let V be a subrepresentation of a finite rational representation V' of G . We write $V = V^G \oplus W$ and $V' = (V')^G \oplus W'$, where W and W' are each the unique subrepresentations of V and V' respectively with this property as in our assumption. Let $w \in W$. We write $w = u' + w'$ where $u' \in (V')^G$ and $w' \in W'$. We choose a basis $\{u'_i\}_{i \in [r]}$ of $(V')^G$ and $\{w'_j\}_{j \in [s]}$ of W' and write $w = \sum_{i=1}^r \lambda_i u'_i + \sum_{j=1}^s \mu_j w'_j$. For $i \in [r]$, let us consider $\hat{u}'_i \in (V')^*$, the dual basis element of u'_i with respect to the basis $\{u'_i\}_{i \in [r]} \cup \{w'_j\}_{j \in [s]}$ of V' . Because of our assumption we have $(W^*)^G = \{0\}$, so we must have $\hat{u}'_i|_W = 0$, and therefore $\lambda_i = \hat{u}'_i(w) = \hat{u}'_i|_W(w) = 0$. We retrieve $u' = 0$, implying $w = w' \in W'$. We have now shown $W \subseteq W'$. Let $v \in V$. With $V^G \subseteq (V')^G$ and $R_V(v) - v \in W \subseteq W'$, we retrieve $R_{V'}(v) - R_V(v) = R_{V'}(v - R_V(v)) = 0$.

This concludes $R_{V'}|_V = R_V$.

(b)

Let V be a finite rational representation of G , and let $R'_V: V \rightarrow Y$ be a G -invariant linear map where $V \subseteq Y$. Via our assumption, we can find a unique subrepresentation W of V such that $V = V^G \oplus W$. We obviously have $R'_V|_{V^G} = \text{id}_{V^G} = R_V|_{V^G}$. Let $w \in W$. We choose a basis $\{w_i\}_{i \in [r]}$ of $U := \text{span}(W + R'_V(w))$, and we write $R'_V(w) = \sum_{i=1}^r \lambda_i w_i$. Let $\{w'_i\}_{i \in [r]}$ be the basis of U^* dual to the previously mentioned basis of U . For $i \in [r]$, we have $(w'_i \circ R'_V)|_W \in (W^*)^G = \{0\}$ via our assumption, and therefore $\lambda_i = w'_i(R'_V(w)) = (w'_i \circ R'_V)|_W(w) = 0$. This means that $R(w) = 0$. We now have shown $R|_W = 0$. This concludes that $R'_V = R_V$.

(c)

This follows immediately from (b): If X is an affine G -variety and $R: K[X] \rightarrow K[X]^G$ is a Reynolds operator and V is a G -stable subspace of $K[X]$, we have that $R|_V: V \rightarrow K[X]$ is a linear map with $V \subseteq K[X]$ and $R_V|_{V^G} = \text{id}_{V^G}$. Therefore we have $R|_V = R_V$.

(d)

Let X be an affine G -variety, $R: K[X] \rightarrow K[X]^G$ a Reynolds operator and W is any G -stable subspace of $K[X]$. now let $w \in W$. Since W is G -stable we have $V_w \subseteq W$ and with (c) therefore $R(w) = R_{V_w}(w) \in V_w^G \subseteq W^G$. We have therefore shown $R(W) \subseteq W^G$. Also $R|_{W^G} = \text{id}_{W^G}$ since $W^G \subseteq K[X]^G$, concluding $R(W) = W^G$. (e)

This follows immediately from (c): Let X be an affine G -variety and $R_1, R_2: K[X] \rightarrow K[X]^G$ each a Reynolds operator. Now let $f \in K[X]$. Then $R_1(f) = R_{V_f}(f) = R_2(f)$. \square

Remark 3.5.1

Note that in lemma 3.5(d) we just showed uniqueness without mentioning existence. In the following, we see that in fact there always exists a Reynolds operator for groups with the previously described properties.

Theorem 3.6

Let G be a linear algebraic group. The following are equivalent:

- (a) G is linearly reductive
- (b) For every finite rational representation V of G there exists a unique subrepresentation W with $V = V^G \oplus W$. For this subrepresentation W we have $(W^*)^G = \{0\}$.
- (c) For every affine G -variety X there exists a Reynolds operator $R: K[X] \rightarrow K[X]^G$.

Proof.

(a) \implies (b)

Let V be a finite rational representation of G . Consider the subspace $((V^*)^G)^\perp \subseteq V$. It is easily seen that this is a subrepresentation of V . In proposition 3.2, we showed that V^G and $(V^*)^G$ are dual to each other. For this reason, we have

$V = V^G \oplus ((V^*)^G)^\perp$. We have shown the existence, now we shall show uniqueness. Let W be a subrepresentation of V with $V = V^G \oplus W$. Again, it is easily seen that $W^\perp \subseteq V^*$ is a subrepresentation. G must act trivially on $W^\perp \subseteq V^*$: Let $f \in W^\perp$, and let $\sigma \in G$. We have $\sigma.f \in W^\perp$ and therefore $\sigma.f - f \in W^\perp$. Now, let $v \in V$. We write $v = u + w$ for (unique) $u \in V^G$ and $w \in W$ and compute:

$$\begin{aligned} (\sigma.f - f)(v) &= (\sigma.f - f)(u) + (\sigma.f - f)(w) \\ &= f(\sigma^{-1}.u) - f(u) + 0 \\ &= f(u) - f(u) = 0 \end{aligned} \tag{5}$$

Which means that $\sigma.f = f$. Hence G does act trivially on W^\perp . This means that $W^\perp \subseteq (V^*)^G$. For reasons of dimension we get $W^\perp = (V^*)^G$, and therefore also $W = (W^\perp)^\perp = ((V^*)^G)^\perp$, which concludes the claim of uniqueness. Finally, we notice that that W and W^* are isomorphic representations (**How clear is this??**), which also means that $(W^*)^G$ and W^G are isomorphic. Since we have $W^G = \{0\}$, we therefore must also have $(W^*)^G = \{0\}$.

(b) \implies (c)

Let X be an affine G -variety. Let $f \in K[X]$. We define the map $R: K[X] \rightarrow K[X]^G$, $f \mapsto R_{V_f}(f)$. For $f \in K[X]$ we denote by W_f the unique subrepresentation of V_f such that $V_f = V_f^G \oplus W_f$ as in (b). This map is linear: Let $f, g \in K[X]$ and $\lambda \in K$. We notice that $V_f, V_g, V_{\lambda f + g} \subseteq V_f + V_g$, which together with lemma 3.5(a) gives us $R(\lambda f + g) = R_{V_{\lambda f + g}}(\lambda f + g) = R_{V_f + V_g}(\lambda f + g) = \lambda R_{V_f}(f) + R_{V_g}(g) = \lambda R_{V_f}(f) + R_{V_g}(g) = \lambda R(f) + R(g)$. The map R is also a projection onto $K[X]^G$, since for each $f \in K[X]$ we have $V_f^G \subseteq K[X]^G$. R is also G -invariant, since for all $f \in K[X]$ R_{V_f} is G -invariant and for all $\sigma \in G$ we have $V_f = V_{\sigma.f}$. This concludes that R is a Reynolds operator, which shows (c).

(c) \implies (a)

Let V be a finite rational representation of G and let $v \in V^G \setminus \{0\}$. We choose a basis $\{v_i\}_{i \in [r]}$ of V with $v_1 = v$. Let $\hat{v} \in V^*$ be the dual basis vector of v with respect to the afore mentioned basis. Now we define $p_v: K[V^*] \rightarrow K$, $f \mapsto f(\hat{v})$. Consider the isomorphism $\Phi: V \rightarrow (V^*)^*$, $w \mapsto (\phi \mapsto \phi(w))$. We have $(V^*)^* \subseteq K[V^*]$. Since V^* is a finite rational representation and since via our assumption (c) we have a Reynolds operator $R: K[V^*] \rightarrow K[V^*]^G$, we can define $\psi_v := p_v \circ R \circ \Phi: V \rightarrow K$. Since each map is linear, we have $\psi_v \in V^*$. After we notice that since $v \in V^G$ we have $\Phi(v) \in K[V^*]^G$, we can calculate $\psi_v(v) = p_v(\Phi(v)) = \Phi(v)(\hat{v}) = \hat{v}(v) = 1 \neq 0$. This concludes that G is linearly reductive, showing (a). \square

3.2 Hilbert's Finiteness Theorem

Proposition 3.7

See [DK15, p.41 Corollary 2.2.7]

Let G be a linearly reductive group, and let $R: K[X] \rightarrow K[X]^G$ be the Reynolds operator for an affine G -variety X . If $f \in K[X]^G$ and $g \in K[X]$ we have

$R(fg) = fR(g)$, id est the Reynolds operator is a $K[X]^G$ -module homomorphism.

Proof. Let $f \in K[X]^G$ and $g \in K[X]$. We can decompose $V_g = V_g^G \oplus W_g$ uniquely, where W_g is a subrepresentation of V_g . We have $(fW)^G = 0$: Let $h \in (fW)^G$, say $h = fw$ where $w \in W$. \square

Theorem 3.8: Hilbert's Finiteness Theorem

If G is linearly reductive and V is a finite-dimensional rational G -representation, the invariant ring $K[V]^G$ is finitely generated.

Proof. Let $I_{>0}$ denote the ideal generated by all non-constant invariants in $K[V]$. Since $K[V]$ is noetherian, there exist finitely many $\{f_i\}_{i \in [r]} \subseteq K[V]$ such that $(\{f_i\}_{i \in [r]}) = I_{>0}$. These must be non-constant invariants (the zero polynomial will always be omitted). Claim: $K[\{f_i\}_{i \in [r]}] = K[V]^G$. The inclusion " \subseteq " is clear. To show is " \supseteq ". This is equivalent to showing that for all $d \in \mathbb{N}$ we have $K[V]_{\leq d}^G \subseteq K[\{f_i\}_{i \in [r]}]$. We will show our claim via induction over the degree d . For $g \in K[V]_{\leq 1}^G = K$ we are already done since $K \subseteq K[\{f_i\}_{i \in [r]}]$. Now assume that for $d \in \mathbb{N}$ we have $K[V]_{\leq d}^G \subseteq K[\{f_i\}_{i \in [r]}]$. Let $g \in K[V]_{\leq d+1}^G$. By construction, $g \in I_{>0}$, therefore there exist $\{g_i\}_{i \in [r]} \subseteq K[V]$ such that $g = \sum_{i=1}^r g_i f_i$. Since the f_i are of non-constant and $\deg g < d+1$, we must have $\deg g_i < d$. We now make use of the Reynolds Operator:

$$g = R(g) = R\left(\sum_{i=1}^r g_i f_i\right) = \sum_{i=1}^r R(g_i) f_i \quad (6)$$

Since R maps $K[V]_{\leq d}$ to $K[V]_{\leq d}^G$, we have $R(g_i) \in K[V]_{\leq d}^G \subseteq K[\{f_i\}_{i \in [r]}]$ by our induction hypothesis. This finally implies $g \in K[\{f_i\}_{i \in [r]}]$, which concludes our proof: We have $K[V]^G = K[\{f_i\}_{i \in [r]}]$ which means that $K[V]^G$ is finitely generated, which was to show. \square

Lemma 3.9

See [DK15, 2.2.8]

Proof. \square

Lemma 3.10

See [DK15, 2.2.9]

Proof. \square

Theorem 3.11

If G is a linearly reductive group and X is an affine G -variety, $K[X]^G$ is finitely generated.

3.3 The Reynolds Operator Of A Linear Algebraic Group

In theorem 3.6 we have learned about three characterizations of linearly reductive groups, but for a given linear algebraic group, it is still hard to concretely show that it is linearly reductive. We will soon learn about a fourth way to characterize linearly reductive groups, which will motivate the main theme of my work: Cayley's Ω -process.

Definition 3.12

Define the multiplication on $K[G]^*$, denoted by $*$, as follows: For $\alpha, \beta \in K[G]^*$:

$$\alpha * \beta := (\alpha \otimes \beta) \circ m^* \quad (7)$$

More slowly: For $f \in K[G]$ we get $m^*(f) = \sum_i g_i \otimes h_i$ (with $g_i, h_i \in K[G]$), therefore the Kronecker-product gives us

$$(\alpha * \beta)(f) = \sum_i \alpha(g_i) \otimes \beta(h_i) = \sum_i \alpha(g_i) \beta(h_i) \quad (8)$$

As usual, we identify $K \otimes K$ with K canonically.

Example: TODO

Proposition 3.13

The multiplication $*$ makes $K[G]^*$ into an associative algebra with the neutral element $\epsilon := \epsilon_e$ (Note: $\epsilon_\sigma(f) = f(\sigma)$).

Proof. First, a small observation:

$$(m^* \otimes \text{id}) \circ m^* = (\text{id} \otimes m^*) \circ m^* \quad (9)$$

This is true because m (and \otimes) is associative. Then, for $\delta, \gamma, \varphi \in K[G]^*$:

$$\begin{aligned} (\delta * \gamma) * \varphi &= (((\delta \otimes \gamma) \circ m^*) \otimes \varphi) \circ m^* = ((\delta \otimes \gamma) \otimes \varphi) \circ (m^* \otimes \text{id}) \circ m^* \\ &= (\delta \otimes (\gamma \otimes \varphi)) \circ (\text{id} \otimes m^*) \circ m^* = (\delta \otimes ((\gamma \otimes \varphi) \circ m^*)) \circ m^* = \delta * (\gamma * \varphi) \end{aligned} \quad (10)$$

showing the associativity. It should be clear that ϵ is the neutral element. This concludes that $K[G]^*$ is an associative algebra. \square

Now we can formally define $K[G]^*$ -actions.

Definition 3.14

Let G act regularly on a vector-space V via μ , from which we retrieve μ' as described in definition 2.5. $K[G]^*$ then acts on V as follows:

$$\delta \cdot v := ((\delta \otimes \text{id}) \circ \mu')(v) \quad (11)$$

Remark 3.14.1

If we look at definition 2.5, we can see that this newly defined $K[G]^*$ -action is an extension of the given G -action in the following way: The subgroup $\{\epsilon_\sigma \mid \sigma \in G\}$ of $K[G]^*$ is isomorphic to G , and its induced action coincides with the given action: For $\sigma \in G$ and for $v \in V$ we have:

$$\sigma \cdot v = \epsilon_\sigma \cdot v \quad (12)$$

Remark 3.14.2

The subalgebra

$$\{ \delta \in K[G]^* \mid \forall f, g \in K[G] : \delta(fg) = \delta(f)g(e) + f(e)\delta(g) \} \quad (13)$$

is called the **Lie algebra**.

Proposition 3.15

Let G be linearly reductive, and let G act regularly on an affine variety X , which induces a rational G -action on $K[X]$ as described in definition 2.6. Then, the following map

$$R: K[X] \longrightarrow K[X]^G \quad f \mapsto R_G \cdot f \quad (14)$$

defines a Reynolds operator.

Proof. As per our construction from definition 3.14, the linearity of this map should be clear. Let us now check that R does map polynomials to invariant polynomials. For this, let $f \in K[X]$, $\sigma \in G$ and $x \in X$. Write $\mu'(f) = \sum_i p_i \otimes g_i \in K[G] \otimes K[X]$. Now we compute:

$$\begin{aligned} \sigma.(R_G \cdot f)(x) &= (R_G \otimes \text{id})(\mu'(f))(\sigma^{-1}.x) \\ &= \sum_i R_G(p_i) g_i(\sigma^{-1}.x) \\ &= \sum_i R_G(p_i) \cdot (\sigma.g_i)(x) \\ &= \sum_i R_G(\sigma.p_i) \cdot (\sigma.g_i)(x) \\ &= (R_G \otimes \text{id})(\sum_i \sigma.p_i \otimes \sigma.g_i)(x) \\ &= (R_G \otimes \text{id})(\mu'(f))(x) = (R_G \cdot f)(x) \end{aligned} \quad (15)$$

The second-to-last equation can be seen as follows: For all $\tau \in G$ we have

$$\begin{aligned} \sum_i \sigma.p_i(\tau) \sigma.g_i &= \sum_i p_i(\tau \sigma^{-1}) \sigma.g_i \\ &= \tau \sigma^{-1}.(\sigma.f) \\ &= \tau.f \\ &= \sum_i p_i(\tau) g_i \end{aligned} \quad (16)$$

making use of proposition 2.7. We have now shown that $R_G \cdot f \in K[V]^G$. If $f \in K[V]^G$, we have $\mu'(f) = 1 \otimes f$, therefore not done yet pls finish \square

4 Cayley's Ω -Process

We want to express the Reynolds Operator in a concrete way. For the Group GL_n , we can explicitly formulate it with the help of Cayley's Ω -Process. First, how is GL_n an affine variety? Consider $K^{n \times n} \times K$, and its coordinate ring $K[\{Z_{i,j}\}_{i,j \in [n]}, D]$. Now define $I := \left(\left(\sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i \in [n]} Z_{i, \sigma(i)} \right) \cdot D - 1 \right) =$

$(\det(Z) \cdot D - 1)$, where $Z := [Z_{i,j}]_{i,j \in [n]}$. Then $V(I) = \{(z, d) \mid z \in \text{GL}_n, d = \det(z)^{-1}\}$. Equipped with the componentwise multiplication (GL_n and $K \setminus \{0\}$, respectively), this is a linear algebraic group isomorphic to GL_n . The coordinate ring $K[\text{GL}_n]$ is isomorphic to $K\left[\{Z_{i,j}\}_{i,j \in [n]}, \det(Z)^{-1}\right] \subseteq K(\{Z_{i,j}\}_{i,j \in [n]})$, and we will write it as such from now on.

Definition 4.1: Cayley's Ω -Process

We call

$$\Omega := \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i \in [n]} \frac{\partial}{\partial z_{i, \sigma(i)}} \quad (17)$$

Cayley's Ω -Process. It can also be thought of as $\Omega = \det\left(\frac{\partial}{\partial Z}\right)$, where $\frac{\partial}{\partial Z} := \left[\frac{\partial}{\partial z_{i,j}}\right]_{i,j \in [n]}$.

Lemma 4.2

$$\left(\det(Z)^{-1} \cdot \otimes \Omega\right) \circ m^* = m^* \circ \Omega = \left(\Omega \otimes \det(Z)^{-1} \cdot\right) \circ m^* \quad (18)$$

where I write “ $p \cdot$ ” for the operation *multiply with p* for a polynomial $p \in K[\text{GL}_n]$ (but the reader must not worry, this is the only time I will make use of this notation).

Proof. Here, we will follow the same convention as described in chapter 3: The “left” and “right” inputs of m will be represented by $X = [X_{i,j}]_{i,j \in [n]}$ and $Y = [Y_{i,j}]_{i,j \in [n]}$ in the occurring polynomials respectively, and the outputs $m = [m_{i,j}]_{i,j \in [n]}$ are indexed the same as the inputs of the polynomials in $Z_{1,1}, Z_{1,2}, \dots, Z_{n,n}$.

Let $f \in K[\text{GL}_n]$. Then $f \circ m \in K\left[\{X_{i,j}\}_{i,j \in [n]}, \det(X)^{-1}, \{Y_{i,j}\}_{i,j \in [n]}, \det(Y)^{-1}\right]$. For fixed $i, j \in [n]$ we have

$$\begin{aligned} \left(\text{id} \otimes \frac{\partial}{\partial Z_{i,j}}\right) (m^*(f)) &= t \left(\frac{\partial}{\partial Y_{i,j}} (f \circ m) \right) = t \left(\sum_{k,l \in [n]} \left(\left(\frac{\partial}{\partial Z_{k,l}} f \right) \circ m \right) \cdot \frac{\partial}{\partial Y_{i,j}} m_{k,l} \right) \\ &= t \left(\sum_{k=1}^n \left(\left(\frac{\partial}{\partial Z_{k,j}} f \right) \circ m \right) \cdot X_{k,i} \right) = \sum_{k=1}^n (Z_{k,i} \cdot \otimes \text{id}) \left(m^* \left(\frac{\partial}{\partial Z_{k,j}} f \right) \right) \end{aligned} \quad (19)$$

Note the use of t as described in chapter 3 to aid in rephrasing terms. Success-

sively applying this yields

$$\begin{aligned}
(\text{id} \otimes \Omega)(m^*(f)) &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) \left(\text{id} \otimes \prod_{i=1}^n \frac{\partial}{\partial Z_{i,\sigma(i)}} \right) (m^*(f)) \\
&= \sum_{\sigma \in S_n} \text{sgn}(\sigma) \sum_{k \in [n]^n} \left(\prod_{i=1}^n Z_{k(i),i} \cdot \otimes \text{id} \right) \left(m^* \left(\prod_{j=1}^n \frac{\partial}{\partial Z_{k(j),\sigma(j)}} f \right) \right) \\
&= \sum_{k \in [n]^n} \left(\prod_{i=1}^n Z_{k(i),i} \cdot \otimes \text{id} \right) \left(m^* \left(\sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{j=1}^n \frac{\partial}{\partial Z_{k(j),\sigma(j)}} f \right) \right) \\
&= \sum_{k \in S_n} \left(\prod_{i=1}^n Z_{k(i),i} \cdot \otimes \text{id} \right) \left(m^* \left(\sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{j=1}^n \frac{\partial}{\partial Z_{k(j),\sigma(j)}} f \right) \right) \\
&= \sum_{k \in S_n} \left(\prod_{i=1}^n Z_{k(i),i} \cdot \otimes \text{id} \right) (m^*(\text{sgn}(k)\Omega(f))) = (\det(Z) \cdot \otimes \text{id})(m^*(\Omega(f)))
\end{aligned} \tag{20}$$

This immediately shows the first equality, and the second equality is proven analogously. \square

Lemma 4.3

For $p \in \mathbb{N}$, $c_{p,n} := \Omega^p(\det(Z)^p) = \det\left(\frac{\partial}{\partial Z}\right)^p(\det(Z)^p)$ is a nonnegative integer.

Proof. Write $\det(Z)^p = \sum_i a_i m_i(\{Z_{k,l}\}_{k,l \in [n]})$, where $a_i \in \mathbb{Z} \setminus \{0\}$ and m_i are (monic) monomials. Then

$$\Omega^p(\det(Z)^p) = \sum_i a_i m_i \left(\left\{ \frac{\partial}{\partial Z_{k,l}} \right\}_{k,l \in [n]} \right) \left(\sum_j a_j m_j(\{Z_{k,l}\}_{k,l \in [n]}) \right) \tag{21}$$

Notice that $m_i \left(\left\{ \frac{\partial}{\partial Z_{k,l}} \right\}_{k,l \in [n]} \right) (m_j(\{Z_{k,l}\}_{k,l \in [n]}))$ is zero for $i \neq j$ and a strictly positive integer for $i = j$. Therefore in particular

$$c_{p,n} = \sum_i a_i^2 m_i \left(\left\{ \frac{\partial}{\partial Z_{k,l}} \right\}_{k,l \in [n]} \right) (m_i(\{Z_{k,l}\}_{k,l \in [n]})) \in \mathbb{N}_{>0} \tag{22}$$

\square

Now, finally, we have the tools to see the following way of expressing the Reynolds Operator.

Theorem 4.4

For $p \in \mathbb{N}$ and $\tilde{f} \in K \left[\{Z_{i,j}\}_{k,l \in [n]} \right]_{pn}$, define for $f = \frac{\tilde{f}}{\det(Z)^p}$:

$$R(f) := \frac{\Omega^p \tilde{f}}{c_{p,n}} \tag{23}$$

The linear extension of this (mapping anything else in $K[\mathrm{GL}_n]$ to zero), defines the Reynolds Operator R_{GL_n} , which makes GL_n *linearly reductive*.

Proof. First, check that this is well defined: For any such term, expanding the fraction by $\det(Z)^q$ will yield the same result. Also, Ω^p is linear for any $p \in \mathbb{N}$. Now we show that R is GL_n -invariant. First, I will introduce a notation: For $f \in K[\mathrm{GL}_n]$ and $\alpha \in \mathrm{GL}_n$, define $\alpha \cdot f := (x \mapsto f(x\alpha^{-1}))$ (again, I assure the reader that this proof contains the only occurrence of this notation). This is *not* an action, but a right action (normal actions should be called “left actions”). Let $p \in \mathbb{N}$, $\tilde{f} \in K[\mathrm{GL}_n]_{pn}$ and $f := \frac{\tilde{f}}{\det(Z)^p}$. For $\beta, \gamma \in \mathrm{GL}_n$, we notice

$$\begin{aligned} R(\beta \cdot f)(\gamma) &= R\left(\frac{\det(\beta)^p \cdot \beta \cdot \tilde{f}}{\det(Z)^p}\right)(\gamma) = \frac{\det(\beta)^p \cdot \Omega^p(\beta \cdot \tilde{f})(\gamma)}{c_{p,n}} \\ &= \frac{1}{c_{p,n}} \cdot (\epsilon_{\beta^{-1}} \otimes \epsilon_\gamma) \left(((\det(Z)^{-p} \cdot \otimes \Omega^p) \circ m^*) (\tilde{f}) \right) \\ &= \frac{1}{c_{p,n}} \cdot (\epsilon_{\beta^{-1}} \otimes \epsilon_\gamma) \left(((\Omega^p \otimes \det(Z)^{-p} \cdot) \circ m^*) (\tilde{f}) \right) \quad (24) \\ &= \frac{\Omega^p(\gamma^{-1} \cdot \tilde{f})(\beta^{-1}) \cdot \det(\gamma^{-1})^p}{c_{p,n}} = R\left(\frac{\gamma^{-1} \cdot \tilde{f} \cdot \det(\gamma^{-1})^p}{\det(Z)^p}\right)(\beta^{-1}) \\ &= R(\gamma^{-1} \cdot f)(\beta^{-1}) \end{aligned}$$

Since each $\frac{\partial}{\partial Z_{i,j}}$ lowers the degree of a monomial by one or maps it to zero, R maps to K , and therefore for $\delta \in \mathrm{GL}_n$ and $g \in K[\mathrm{GL}_n]$ we have $R(g)(\delta) = R(g) \in K$. We then get for all $\alpha \in \mathrm{GL}_n$

$$\begin{aligned} R(\alpha \cdot f) &= R(\alpha \cdot f)(I_n) = R(I_n^{-1} \cdot f)(\alpha^{-1}) \\ &= R(I_n^{-1} \cdot f) = R(f) \end{aligned} \quad (25)$$

which shows the GL_n -invariance. Finally, the definition immediately gives us that R restricted to K is the identity. As mentioned in lemma 3.5(e), the uniqueness of the Reynolds Operator implies $R = R_{\mathrm{GL}_n}$. \square

Now we will look at the Reynolds Operator R_{SL_n} .

Corollary 4.4.1

With the identification $K[\mathrm{GL}_n] = K\left[\{Z_{k,l}\}_{k,l \in [n]}, \det(Z)^{-1}\right]$, view $K[\mathrm{SL}_n] = K[\mathrm{GL}_n]/I$ where $I = (\det(Z) - 1)$. Now, for $p \in \mathbb{N}$ and $\tilde{f} \in K\left[\{Z_{i,j}\}_{k,l \in [n]}\right]_{pn}$ define for $f = \frac{\tilde{f}}{\det(Z)^p} + I$:

$$R(f) := R_{\mathrm{GL}_n}\left(\frac{\tilde{f}}{\det(Z)^p}\right) + I = \frac{\Omega^p \tilde{f}}{c_{p,n}} + I \quad (26)$$

The linear extension of this (mapping anything else in $K[\mathrm{SL}_n]$ to zero), defines the Reynolds Operator R_{SL_n} , making SL_n *linearly reductive*.

Proof. First, we will show $K[\mathrm{GL}_n]^{\mathrm{SL}_n} = K[\det(Z), \det(Z)^{-1}]$ (action by left multiplication). Let $g \in K[\mathrm{GL}_n]^{\mathrm{SL}_n}$, and let $\alpha \in \mathrm{GL}_n$. Now for $x \in K \setminus \{0\}$, define $M(z) = [z_{i,j}]_{i,j \in [n]} \in \mathrm{GL}_n$ to be the matrix with $z_{1,1} = z$ and $z_{i,j} = \delta_{i,j}$ for $(i,j) \neq (1,1)$. Note that $M(\det(\alpha))^{-1} \alpha \in \mathrm{SL}_n$. Define $h := (\beta \mapsto g(M(\det(\beta)))) \in K[\det(Z), \det(Z)^{-1}]$. Now

$$\begin{aligned} g(\alpha) &= \left(M(\det(\alpha))^{-1} \alpha \right) \cdot g(\alpha) = g(M(\det(\alpha)) \alpha^{-1} \alpha) \\ &= g(M(\det(\alpha))) = h(\alpha) \end{aligned} \quad (27)$$

Therefore $g = h \in K[\det(Z), \det(Z)^{-1}]$. Conversely it is easy to see that $K[\det(Z), \det(Z)^{-1}] \subseteq K[\mathrm{GL}_n]^{\mathrm{SL}_n}$.

Now, define $\hat{R}: K[\mathrm{GL}_n] \rightarrow K[\mathrm{GL}_n]^{\mathrm{SL}_n}$ as follows:

For $p, r \in \mathbb{N}$, $\tilde{f} \in K[\{Z_{k,l \in [n]}\}]_{rn}$, and $f = \frac{\tilde{f}}{\det(Z)^p}$, define

$$\hat{R}(f) := \det(Z)^{r-p} \cdot \frac{\Omega^r \tilde{f}}{c_{r,n}} = \det(Z)^{r-p} \cdot R_{\mathrm{GL}_n} \left(\frac{\tilde{f}}{\det(Z)^r} \right) \quad (28)$$

and as before we define the images of the other elements by linear extension. Well-definedness follows from the same observations as in the proof of the theorem (and $r-p = (r+q) - (p+q)$ for $q \in \mathbb{N}$). This map is the identity on $K[\mathrm{GL}_n]^{\mathrm{SL}_n}$: If $\tilde{f} \in K[\{Z_{k,l \in [n]}\}]_{rn}$ and $\tilde{f} \in K[\mathrm{GL}_n]^{\mathrm{SL}_n}$, then $\tilde{f} = \lambda \det(Z)^r$ with $\lambda \in K$. Therefore we can say w.l.o.g. either $p = 0$ or $r = 0$. Then it is quite clear that then f gets mapped to f . Finally, the SL_n -invariance also follows from the last theorem: Let $\alpha \in \mathrm{SL}_n$. Then

$$\begin{aligned} \hat{R}(\alpha \cdot f) &= \hat{R} \left(\frac{\det(\alpha)^p \cdot \alpha \cdot \tilde{f}}{\det(Z)^p} \right) = \det(Z)^{r-p} \cdot R_{\mathrm{GL}_n} \left(\frac{\det(\alpha)^p \cdot \alpha \cdot \tilde{f}}{\det(Z)^r} \right) \\ &= \det(Z)^{r-p} \cdot R_{\mathrm{GL}_n} \left(\frac{\det(\alpha)^r \cdot \alpha \cdot \tilde{f}}{\det(Z)^r} \right) \\ &= \det(Z)^{r-p} \cdot R_{\mathrm{GL}_n} \left(\alpha \cdot \left(\frac{\tilde{f}}{\det(Z)^r} \right) \right) \\ &= \det(Z)^{r-p} \cdot R_{\mathrm{GL}_n} \left(\frac{\tilde{f}}{\det(Z)^r} \right) = \hat{R}(f) \end{aligned} \quad (29)$$

where we used $\det(\alpha)^p = 1 = \det(\alpha)^r$ and the GL_n -invariance of R_{GL_n} . Thus we have shown that \hat{R} is the Reynolds-Operator for the action of SL_n on GL_n by left-multiplication.

Lastly, this shows our proposed statement $R = R_{\mathrm{SL}_n}$, since $\det(Z) \sim 1$. \square

5 Examples (not a section in the final version)

Let us apply the Reynolds operator with respect to an action on concrete polynomials. Before we look at finite generators of Hilbert's nullcone (which we will

talk about later), we will just look at generators of the polynomial ring.

Example 5.1

Consider the group $G = \text{SL}_2$ and the vector space $V = \{ A \in \mathbb{R}^{2 \times 2} \mid A^T = A \}$. Now we will look at the following action:

$$\begin{aligned} \mu: \quad \text{SL}_2 \times V &\longrightarrow V \\ (S, A) &\longmapsto SAS^T \end{aligned} \quad (30)$$

Now consider the following for $S \in \text{SL}_2$ and $A \in V$:

$$\begin{aligned} S &= \begin{bmatrix} s_{1,1} & s_{1,2} \\ s_{2,1} & s_{2,2} \end{bmatrix} & A &= \begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{bmatrix} \\ S^{-1} &= \begin{bmatrix} s_{2,2} & -s_{1,2} \\ -s_{2,1} & s_{1,1} \end{bmatrix} \end{aligned} \quad (31)$$

We then have

$$\begin{aligned} S^{-1}.A &= S^{-1}A(S^{-1})^T \\ &= \begin{bmatrix} a_{1,1}s_{2,2}^2 - 2a_{1,2}s_{1,2}s_{2,2} + a_{2,2}s_{1,2}^2 & -a_{1,1}s_{2,1}s_{2,2} + a_{1,2}s_{1,1}s_{2,2} + a_{1,2}s_{1,2}s_{2,1} - a_{2,2}s_{1,1}s_{2,1} \\ -a_{1,1}s_{2,1}s_{2,2} + a_{1,2}s_{1,1}s_{2,2} + a_{1,2}s_{1,2}s_{2,1} - a_{2,2}s_{1,1}s_{2,1} & a_{1,1}s_{2,1}^2 - 2a_{1,2}s_{1,1}s_{2,1} + a_{2,2}s_{1,1}^2 \end{bmatrix} \end{aligned} \quad (32)$$

Notice that we also have

$$\begin{aligned} \det \left(\frac{\partial}{\partial S} \right)^n &= \left(\frac{\partial}{\partial S_{1,1}} \frac{\partial}{\partial S_{2,2}} - \frac{\partial}{\partial S_{1,2}} \frac{\partial}{\partial S_{2,1}} \right)^n \\ &= \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{\partial^{n-k}}{\partial S_{1,1}} \frac{\partial^k}{\partial S_{1,2}} \frac{\partial^k}{\partial S_{2,1}} \frac{\partial^{n-k}}{\partial S_{2,2}} \end{aligned} \quad (33)$$

It is quite cumbersome to calculate the Reynolds Operator of general polynomials. We will look at the monomial $A_{1,1}^2$, for which we have

$$\begin{aligned} \mu'(A_{1,1}^2) &= S_{2,2}^4 \otimes A_{1,1}^2 - 4S_{1,2}S_{2,2}^3 \otimes A_{1,1}A_{1,2} + 2S_{1,2}^2S_{2,2}^2 \otimes A_{1,1}A_{2,2} \\ &\quad + 4S_{1,2}^2S_{2,2}^2 \otimes A_{1,2}^2 - 4S_{1,2}^3S_{2,2} \otimes A_{1,2}A_{2,2} + S_{1,2}^4 \otimes A_{2,2}^2 \\ &= \frac{S_{2,2}^4}{\det(S)^2} \otimes A_{1,1}^2 - \frac{4S_{1,2}S_{2,2}^3}{\det(S)^2} \otimes A_{1,1}A_{1,2} \\ &\quad + \frac{2S_{1,2}^2S_{2,2}^2}{\det(S)^2} \otimes A_{1,1}A_{2,2} + \frac{4S_{1,2}^2S_{2,2}^2}{\det(S)^2} \otimes A_{1,2}^2 \\ &\quad - \frac{4S_{1,2}^3S_{2,2}}{\det(S)^2} \otimes A_{1,2}A_{2,2} + \frac{S_{1,2}^4}{\det(S)^2} \otimes A_{2,2}^2 \end{aligned} \quad (34)$$

We can now apply the Reynolds operator in the way we discussed it in proposition 3.15 in combination with Cayley's Ω -process. Since all terms in $K[\text{SL}_2]$ are

already of degree 2, apply the same derivatives to each summand and calculate:

$$\begin{aligned}
& R_G \cdot A_{1,1}^2 \\
&= \left(\frac{\partial}{\partial S_{1,1}} \frac{\partial}{\partial S_{2,2}} - 2 \frac{\partial}{\partial S_{1,1}} \frac{\partial}{\partial S_{1,2}} \frac{\partial}{\partial S_{2,1}} \frac{\partial}{\partial S_{2,2}} + \frac{\partial}{\partial S_{1,2}} \frac{\partial}{\partial S_{2,1}} \right) \cdot A_{1,1}^2 \quad (35) \\
&= 0
\end{aligned}$$

The zero-polynomial is not very interesting, so applying the Reynolds Operator to any polynomial will not always produce interesting results. We will try again for the polynomial $A_{1,2}^2$. We calculate

$$\begin{aligned}
& \mu'(A_{1,2}^2) \\
&= S_{2,1}^2 S_{2,2}^2 \otimes A_{1,1}^2 - 2S_{1,1} S_{2,1} S_{2,2}^2 \otimes A_{1,1} A_{1,2} \\
&\quad - 2S_{1,2} S_{2,1}^2 S_{2,2} \otimes A_{1,2}^2 + 2S_{1,1} S_{1,2} S_{2,1} S_{2,2} \otimes A_{1,1} A_{2,2} \\
&\quad + S_{1,1}^2 S_{2,2}^2 \otimes A_{1,2}^2 + 2S_{1,1} S_{1,2} S_{2,1} S_{2,2} \otimes A_{1,2}^2 \\
&\quad - 2S_{1,1}^2 S_{1,2} S_{2,2} \otimes A_{1,2} A_{2,2} + S_{1,2}^2 S_{2,1}^2 \otimes A_{1,2}^2 \\
&\quad - 2S_{1,1} S_{1,2}^2 S_{2,1} \otimes A_{1,2} A_{2,2} + S_{1,1}^2 S_{1,2}^2 \otimes A_{2,2}^2 \\
&= \frac{S_{2,1}^2 S_{2,2}^2}{\det(S)^2} \otimes A_{1,1}^2 - \frac{2S_{1,1} S_{2,1} S_{2,2}^2}{\det(S)^2} \otimes A_{1,1} A_{1,2} \\
&\quad - \frac{2S_{1,2} S_{2,1}^2 S_{2,2}}{\det(S)^2} \otimes A_{1,2}^2 + \frac{2S_{1,1} S_{1,2} S_{2,1} S_{2,2}}{\det(S)^2} \otimes A_{1,1} A_{2,2} \\
&\quad + \frac{S_{1,1}^2 S_{2,2}^2}{\det(S)^2} \otimes A_{1,2}^2 + \frac{2S_{1,1} S_{1,2} S_{2,1} S_{2,2}}{\det(S)^2} \otimes A_{1,2}^2 \\
&\quad - \frac{2S_{1,1}^2 S_{1,2} S_{2,2}}{\det(S)^2} \otimes A_{1,2} A_{2,2} + \frac{S_{1,2}^2 S_{2,1}^2}{\det(S)^2} \otimes A_{1,2}^2 \\
&\quad - \frac{2S_{1,1} S_{1,2}^2 S_{2,1}}{\det(S)^2} \otimes A_{1,2} A_{2,2} + \frac{S_{1,1}^2 S_{1,2}^2}{\det(S)^2} \otimes A_{2,2}^2
\end{aligned} \quad (36)$$

Again, all $K[\text{SL}_2]$ terms are of degree 2, therefore we can simplify and calculate

$$\begin{aligned}
& R_G \cdot A_{1,2}^2 \\
&= \left(\frac{\partial}{\partial S_{1,1}} \frac{\partial}{\partial S_{2,2}} - 2 \frac{\partial}{\partial S_{1,1}} \frac{\partial}{\partial S_{1,2}} \frac{\partial}{\partial S_{2,1}} \frac{\partial}{\partial S_{2,2}} + \frac{\partial}{\partial S_{1,2}} \frac{\partial}{\partial S_{2,1}} \right) \cdot A_{1,2}^2 \quad (37) \\
&= -\frac{4}{12} A_{1,1} A_{2,2} + \frac{4}{12} A_{1,2}^2 - \frac{4}{12} A_{1,2}^2 + \frac{4}{12} A_{1,2}^2 \\
&= -\frac{1}{3} \det(A)
\end{aligned}$$

This is in line with what we expect: $K[V]^{\text{SL}_n} = K[\det(A)]$.

Example 5.2

Consider the cross ratio. We can view its domain as the affine variety

$$\begin{aligned}
X &:= \{ (a, b, c, d) \in (\mathbb{K}^2)^4 \mid (b_1 c_2 - b_2 c_1)(d_1 a_2 - d_2 a_1) \neq 0 \} \\
&= \{ (a, b, c, d) \in (\mathbb{K}^2)^4 \mid \det(b, c) \det(d, a) \neq 0 \} \quad (38)
\end{aligned}$$

as described in [yaboi]. Now consider the action of GL_2 on X via pointwise application. The *cross ratio* $\mathrm{cr} \in K[X]$ defined as follows

$$\begin{aligned} \mathrm{cr}: \quad X &\longrightarrow K \\ (a, b, c, d) &\longmapsto \frac{\det(a, b) \det(c, d)}{\det(b, c) \det(d, a)} \end{aligned} \quad (39)$$

is an invariant under this action, as is easily seen. Exactly this GL_2 -invariance guarantees a coordinate-independent definition of the cross ratio in the projective space. The invariant ring $K[X]^{\mathrm{GL}_n}$ is finitely generated, though we can't apply the Reynolds operator directly, since X is not a vector space.

Example 5.3

(K needs to be algebraically closed for Zariski-denseness of diagonalizable matrices...)

Consider GL_n viewed as the group of all change-of-coordinates transformations for endomorphisms on K^n , that is the rational representation

$$\begin{aligned} \mu: \quad \mathrm{GL}_n \times K^{n,n} &\longrightarrow K^{n,n} \\ (\sigma, A) &\longmapsto \sigma A \sigma^{-1} \end{aligned} \quad (40)$$

What are its invariants? The invariants are exactly those polynomials that are independent of the choice of the basis. The most well-known invariant is the determinant. From this observation we can find even more: We can follow that the characteristic polynomial of a matrix A , that is $\det(tI_n - A)$, does not change under a change of coordinates. If we write

$$\det(tI_n - A) = \sum_{i=0}^n p_{n,i}(A) t^i \quad (41)$$

this means that every $p_{n,i}$ is an invariant polynomial in $K[K^{n,n}]$! This is how one usually proves that the trace is an invariant polynomial after observing that $p_{n,n-1} = \mathrm{tr}_n$. Are there other invariants than these $p_{n,i}$? No! To see this, we will use a little trick: Consider $D := \{ \delta \in K^{n,n} \mid \delta \text{ diagonalizable} \} \subseteq K^{n,n}$. Since V is Zariski-dense in $K^{n,n}$, we have $K[K^{n,n}] \simeq K[K^{n,n}]_D$ via $p \leftrightarrow p|_D$. For this reason, we will look at the evaluation of an invariant polynomial $p \in K[K^{n,n}]$ only on elements in D , and can deduce what polynomial it is.

Let $p \in K[K^{n,n}]^{\mathrm{GL}_n}$. We define a projection onto the diagonal: $\pi: K^{n,n} \longrightarrow K^n, [A_{i,j}]_{i,j \in [n]} \longmapsto (A_{i,i})_{i \in [n]}$. Consider $\tilde{p} = p([\delta_{i,j} Z_{i,j}]_{i,j \in [n]}) \in K[K^n]$. \tilde{p} is S_n -invariant: If $M_\tau \in \mathrm{GL}_n$ is the matrix corresponding to $\tau \in S_n$, then for all $\tau \in S_n$ and for all $X \in K^n$ we have

$$\begin{aligned} \tau.\tilde{p}(X) &= \tilde{p}(\tau^{-1}.X) \\ &= \tilde{p}(\pi(\mathrm{diag}(\tau^{-1}.X))) \\ &= p(\mathrm{diag}(\tau^{-1}.X)) \\ &= M_\tau.p(\mathrm{diag}(X)) \\ &= p(\mathrm{diag}(X)) = \tilde{p}(X) \end{aligned} \quad (42)$$

From the fundamental theorem of symmetric polynomials we can follow that $\tilde{p} \in \text{span}\{e_{n,i}\}_{i=0}^n$, say $\tilde{p} = \sum_{i=0}^n \lambda_i e_{n,i}$, where $\{e_{n,i}\}_{i=0}^n$ are the elementary symmetric polynomials of dimension n . Now, for a choice (!) of $\sigma_A \in \text{GL}_n$ such that $\sigma_A.A$ is diagonal, we easily see that for $s(A) := \sigma_A.A$ we get $p = p \circ s = \tilde{p} \circ \pi \circ s$, therefore $p = \sum_{i=0}^n \lambda_i e_{n,i} \circ \pi \circ s$. Now we want to show that $e_{n,i} \circ \pi \circ s = p_{n,i}$, which would conclude our claim. For all $A \in D$ we have

$$\begin{aligned} \sum_{i=0}^n (e_{n,i} \circ \pi \circ s)(A) t^i &= \det(t - \sigma_A.A) \\ &= \det(t - A) = \sum_{i=0}^n p_{n,i}(A) t^i \end{aligned} \tag{43}$$

which shows our claim. Note that this is independent of the choice of s , which means that we don't need the axiom of choice (rigorously, as usual, rewrite s as a relation for all possible s instead of a choice of a function...).

6 Further Discussion

fdsafdsa fdsafdsa

References

- [DK15] Harm Derksen and Gregor Kemper. *Computational Invariant Theory*. Springer-Verlag, Berlin Heidelberg, 2015.
- [Stu08] Bernd Sturmfels. *Algorithms in Invariant Theory*. Springer-Verlag, Wien, 2008.