

leroy

Chiara Cimini

Christian Krause

March 30, 2025

# Chapter 1

## Localic Caratheodory Extension

### 1.1 Basics of Locales

**Definition 1** ( $f^*$  and  $f_*$ ). For every continuous Function  $f : X \rightarrow Y$  between topological Spaces, there exists a pair of functors  $(f^*, f_*)$ .

$$f_* = f^{-1} : O(Y) \rightarrow O(X)$$
$$f_* : O(X) \rightarrow O(Y) := A \mapsto \bigcup_{f^*(v) \leq A} v$$

**Lemma 2** ( $f^* \dashv f_*$ ).  $f^*$  is the right adjoint to  $f_*$

*Proof.*

□

### 1.2 Sublocales

**Definition 3** (Nucleus). A nucleus is a map  $e : O(E) \rightarrow O(E)$  with the following three properties:

1.  $e$  is idempotent
2.  $U \leq eU$
3.  $e(U \cap V) = e(U) \cap e(V)$

**Lemma 4** (Nucleus). (*Leroy Lemme 3*) *TODO* Let  $e : O(E) \rightarrow O(E)$  be monotonic. The following are equivalent:

1.  $e$  is a nucleus
2. There is a locale  $X$  and a morphism  $f : X \rightarrow E$  such that  $e = f_* f^*$ .
3. Then there is a locale  $X$  and a embedding  $f : X \rightarrow E$  such that  $e = f_* f^*$ .

*Proof.*

□

**Definition 5** (Nucleus Partial Order). For two nuclei  $e$  and  $f$  on  $O(E)$ , we say that  $e \leq f$  if  $e(U) \leq f(U)$  for all  $U \in O(E)$ . This relation is a partial order.

**Lemma 6** (Nucleus Intersection). *TODO Quelle StoneSpaces S.51 For a set  $S$  of nuclei, the intersection  $\bigcap S$  can be computed by  $\bigcap S(a) = \bigcap \{j(a) | j \in S\}$ . This function satisfies the properties of a nucleus and of an infimum.*

*Proof.* □

**Definition 7** (Sublocal). (Leroy CH 3) A sublocal  $Y \subset X$  is defined by a nucleus  $e_Y : O(X) \rightarrow O(X)$ , such that  $O(Y) = \text{Im}(e_Y) = \{U \in O(X) | e_Y(U) = U\}$ . The corresponding embedding is  $i_X : O(Y) \rightarrow O(X)$ .  $i_X^*(V) = e_X(V)$ ,  $(i_X)_*(U) = U$  And every nucleus  $e$  on  $O(X)$  defines a sublocal  $Y$  of  $X$  by  $O(Y) = \text{Im}(e)$

**Definition 8** (Sublocal Inclusion). (Stimmt das?)(Leroy Ch 3)  $X \subset Y$  if  $e_Y(u) \leq e_X(u)$  for all  $u$ . This means that the Sublocals are a dual order to the nuclei.

### 1.2.1 (1.4) Sublocal Union and Intersection

**Definition 9** (Union of Sublocals). (Leroy CH 1.4) Let  $(X_i)_i$  be a family of sublocals of  $E$  and  $(e_i)_i$  the corresponding nuclei. For all  $V \in O(E)$ , let  $e(V)$  be the union of all  $W \in O(E)$  which are contained in all  $e_i(V)$ .

**Lemma 10** (Union of Sublocals). (Leroy CH 4) Let  $X_i$  be a family of subframes of  $E$  and  $e_i$  be the corresponding nuclei. For every  $V \in O(E)$ , let  $e(V)$  be the union of all  $W \in O(E)$  which are contained in every  $e_i(V)$ . Then

1.  $e$  is the corresponding nucleus of a sublocale  $X$  of  $E$
2. a sublocale  $Z$  of  $E$  contains  $x$  if and only if it contains all  $X_i$ .  $X$  is thus called the union of  $X_i$  denoted by  $\bigcup_i X_i$

*Proof.* The properties of the nucleus (idempotent, increasing, preserving intersection) can be verified by unfolding the definition of  $e(V)$ . □

**Lemma 11** (Sublocal Union equals Nucleus Intersection). For a family of sublocals  $X_i$  of  $E$ , the union  $\bigcup X_i$  is the intersection of the corresponding nuclei. *TODO lean link*

*Proof.* The infimum of the Nuclei is a Supremum of the sublocals, because the Nuclei are a dual order to the sublocals.) This means that it suffices to show that suprema are unique.

TODO Quelle [https://proofwiki.org/wiki/Infimum\\_is\\_Unique](https://proofwiki.org/wiki/Infimum_is_Unique)

Suppose there are two different suprema  $c$  and  $c'$  of a set  $S$ . Because of the definition of a supremum, we that they are both upper bounds of  $S$ . But we also know that the supremum is smaller than any other upper bound, so we get  $c \leq c'$  and  $c' \leq c$ . This means that  $c = c'$ . □

**Definition 12** (Intersection of Sublocals). Let  $(X_i)_i$  be a family of sublocal of  $E$  and  $(e_i)_i$  the corresponding nuclei. For all  $V \in O(E)$ , the intersection  $\bigcap X_i$  is the Union of all Nuclei  $w$  such that  $w \leq x_i$  for all  $x_i \in X_i$

**Lemma 13** (Nucleus Complete Lattice). The Nuclei (and therefore the sublocals) form a complete lattice.

*Proof.* One can prove that the Nuclei are closed under arbitrary intersections by unfolding the definition of the intersection. The supremum is defined as the infimum of the upper Bound. □

**Proposition 14** (Complete Heyting Algebra). *A complete Lattice is a Frame if and only if it is a Heyting Algebra.*

*Proof.* (Source Johnstone:) The Heyting implication is right adjoint to the infimum. This means that the infimum preserves Suprema, since it is a left adjoint. □

**Lemma 15** (Nucleus Heyting Algebra). *The Nuclei form a Heyting Algebra.*

*Proof.* Quelle Johnstone □

**Lemma 16** (Nucleus Frame). *The Nuclei form a frame.*

*Proof.* □

## 1.2.2 (7) Open Sublocals

**Definition 17** ( $e_U$ ). Let  $E$  be a space with  $U, H \in O(E)$ . We denote by  $e_U$  the largest  $W \in O(E)$  such that  $W \cap U \subset H$ . We verify that  $e_U$  is the nucleus of a subspace, which we will temporarily denote by  $[U]$ .

**Lemma 18** ( $e_U$  is a nucleus). *The map  $e_U$  is a nucleus.*

*Proof.* □

**Definition 19** (Open sublocal). For any  $U \in O(E)$ , the sublocal  $[U]$  is called an open sublocal of  $E$ .

**Lemma 20** ((6,7) Open Sublocal Properties). *(Leroy Lemma 6,7)*

1. For all subspaces  $X$  of  $E$  and any  $U \in O(E)$ :

$$X \subset [U] \iff e_X(U) = 1_E$$

2. For all  $U, V \in O(E)$ , we have:

$$[U \cap V] = [U] \cap [V]$$

$$e_{U \cap V} = e_U e_V = e_V e_U$$

$$U \subset V \iff [U] \subset [V]$$

3. For all families  $V_i$  of elements of  $O(E)$ , we have:

$$\cup_i [V_i] = [\cup_i V_i]$$

4.

*Proof.* □

**Definition 21** (Complement). The complement of an open sublocal  $U$  of  $X$  is the sublocal  $X \setminus U$ . (Leroy p. 12) (+ Senf brauchen wir das allgemein??)

**Lemma 22** (Complement Injective). *The complement is injective.*

*Proof.* □

**Definition 23** (Closed Sublocal). A sublocal  $X$  of  $E$  is called closed if  $X = E \setminus U$  for some open sublocal  $U$  of  $E$ .

**Lemma 24** (Intersection of Closed Sublocals). *For any family  $X_i$  of closed sublocals of  $E$ , the intersection  $\bigcap X_i$  is closed (it can be computed by taking the complement of the union of the complements).*

*Proof.* □

**Lemma 25** ((1.8) Properties of Complements). *For any open sublocal  $V$  of  $E$  and any sublocal  $X$  of  $E$ , we have:*

$$V \cup X = E \iff E \setminus V \subset X$$

$$V \cap X = \emptyset \iff X \subset E \setminus V$$

*And thereby:*

$$(E - U = E - V) \implies U = V$$

*Proof.* □

**Lemma 26** ((1.8bis) Properties of Complements Part 2). *For any open sublocal  $V$  of  $E$  and any sublocal  $X$  of  $E$ , we have:*

$$V \cup (E - V) = E \iff V \subset X$$

$$V \cap (E - V) = \emptyset \iff X \subset V$$

*Proof.* □

**Definition 27** (Further Topology).

1.  $IntX$  is the largest open sublocal contained in  $X$
2.  $ExtX$  is the largest open sublocal contained in  $E \setminus X$
3.  $\bar{X}$  is the smallest closed sublocal containing  $X$
4.  $\partial X = \bar{X} \cap (E - IntX)$

**Lemma 28** (Properties of Further Topology).

1.  $\bar{X} = E \setminus Ext(X)$
2.  $\partial X = E \setminus (IntX \cup ExtX)$
3.  $IntX \cup \partial X = \bar{X}$
4.  $ExtX \cup \partial X = E \setminus IntX$

*Proof.* □

### 1.3 Caratheodory Extnesion on Locales

**Definition 29** (Measure on Locales). A measure on a local  $X$  is a map  $\mu : O(X) \rightarrow [0, \infty)$  such that:

1.  $\mu(\emptyset) = 0$
2.  $U \subset V \implies \mu(U) \leq \mu(V)$
3.  $\mu(U \cup V) = \mu(U) + \mu(V) - \mu(V \cap U)$
4. For any increasingly filtered family  $V_i$  of open sublocales of  $X$ , we have:

$$\mu\left(\bigcup_i V_i\right) = \sup_i \mu(V_i)$$

this means: For all  $i$  and  $j$  there exists a  $k$  such that  $V_i \cup V_j \subset V_k$  bzw.  $V_i \subset V_k$  and  $V_j \subset V_k$ .

(Leroy III.1.)

**Definition 30** (Caratheodory). For any measure on a local  $X$ , the caratheodory extension is:

$$\mu(A) = \inf\{\mu(U) \mid A \subset U \in O(X)\}$$

**Lemma 31** (Proptery 0 (Commutes with sup)). *(Leroy lemme 3.1) The caratheodory extension of a measure on a local commutes with unions of increasing families.*

*Proof.*

□

**Lemma 32** (Caratheodory Extensions are monotonic). *The caratheodory extension is monotonic i.e.*

$$A \leq B \implies \mu(A) \leq \mu(B)$$

*Proof.* This is a direct consequence of the definition of the caratheodory extension.

□

**Lemma 33** (Subadditivity). *The Caratheodory extension is subadditive:*

$$\mu(A \cup B) \leq \mu(A) + \mu(b)$$

*Proof.*

□

**Definition 34** (Regular Local). A local is regular, if for all open sublocales  $U$  of  $E$ , the open sublocales  $V$  such that  $V \subset U$  recover  $U$ .

**Definition 35** (Neighborhood).

A neighborhood of a sublocal  $A$  of  $X$  is an open sublocal  $V$  of  $X$  such that  $A \leq V$ .

**Lemma 36** (Regularity of Sublocales). *(Leroy lemme 3.2) In a regular local, any sublocal is regular, meaning that it is the intersection of all open neighborhoods.*

*Proof.*

□

**Lemma 37** (Measure add compl eq top). *(Leroy Lemme 3.3) For any open sublocal  $U$  of a local  $X$ , the caratheodory extension of a measure on  $X$  satisfies*

$$\mu(U) + \mu(X \setminus U) = \mu(X)$$

*Proof.* Siehe Leroy □

**Lemma 38** (Restriction). *The Restriction of a Measure to any open Sublocal is a Measure.*

*Proof.* □

**Lemma 39** (Property 2). *(Leroy Lemm 3.4) For any open sublocal  $U$  and any sublocal  $A$  of a local  $E$ , the caratheodory extension of a measure on  $X$  satisfies*

$$\mu(A) = \mu(A \cap U) + \mu(A \cap (E \setminus U))$$

*Proof.* Siehe Leroy □

**Lemma 40** (Property 3). *(Leroy Lemm 3.5) For a increasing family  $V_\alpha$  of open sublocals of  $E$  and any sublocal  $A$ , we have:*

$$\mu(A \cap (\bigcup V_\alpha)) = \sup_\alpha \mu(A \cap V_\alpha)$$

*Proof.* □

**Lemma 41** (Restriction to a Sublocale). *Let  $A$  be a sublocale of  $E$  with the embedding  $i : A \rightarrow E$ . The restriction of a measure  $\mu$  on  $E$  to  $A$  is a measure on  $A$ :*

$$V \mapsto \mu(i(V)) : \text{Open}(A) \rightarrow \mathbb{R}$$

*Proof.* □

**Proposition 42** (strictly additive). *(Leroy theorem 3.3.1) For any measure on a local  $X$ , the caratheodory extension is strictly additive i.e.  $\mu(A \cup B) = \mu(A) + \mu(B) - \mu(A \cap B)$*

*Proof.* □

**Proposition 43** (reductive). *(Proposition 3.3.1) For any measure on a local  $X$ , the caratheodory extension is reductive i.e. for all  $A \leq X$  the set  $\{A' \subset A, \mu(A') = \mu(A)\}$  has a minimal element*

*Proof.* □

**Lemma 44** (Commutates with inf opens). *(Leroy Lemme 3.6) For any measure on a local  $X$  and a decreasing family  $V_i$  of open sublocals, the caratheodory extension fulfills:  $\mu(\inf V_i) = \inf \mu(V_i)$ .*

*Proof.* □

**Proposition 45** (Commutates with inf). *(Leroy lemme 3.7 et principal) For any measure on a local  $X$ , the caratheodory extension is regular  $\mu(\inf A_i) = \inf \mu(A_i)$ . For decreasing families  $A_i$*

*Proof.* □

**Theorem 46** (Main Theorem (very important)). *For any measure on a local  $X$ , the caratheodory extension is*

1. *strictly additive i.e.  $\mu(A \cup B) = \mu(A) + \mu(B) - \mu(A \cap B)$*
2. *commutes with inf  $\mu(\inf A_i) = \inf \mu(A_i)$*
3. *reductive i.e. for all  $A \leq X$  the set  $\{A' \subset A, \mu(A') = \mu(A)\}$  has a minimal element*

*Proof.* □

## Chapter 2

# Locales correspond to Topology

### 2.1 Leroy Chapter V

**Lemma 47** ((1.10) Intersection of Open and Closed Sublocales). *For any  $U \in O(E)$ , and sublocal  $X$  of  $E$  we have:*

$$e_{U \cap X} = e_U e_X$$

*And for a closed  $F$*

$$e_{X \cap F} = e_X e_F$$

**Lemma 48** (Regular Top to regular local). *Any regular topological space induces a regular local.*

**Lemma 49** (Opens). *(Leroy V.1 Remarque 2) The Open subsets of any good enough topological space correspond precisely to the open sublocales of the corresponding local.*

**Lemma 50** (Subset Sublocal). *(Leroy V.1 Remarque 3) Any subset  $X$  of a good enough topological space  $E$  induces a sublocal  $[X]$  of the corresponding local. This is an order preserving embedding.*

**Definition 51** (Good enough topological space). blackbox to mathlib???

**Lemma 52** (Subset to sublocal Part 1). *(Leroy Proposition 5.1.1)*

*For two subspaces  $X$  and  $Y$  of  $E$  and an open subspaces  $U$  of  $E$ , we have:*

1.  $X \subset Y \implies [X] \subset [Y]$
2.  $X \subset U \iff [X] \subset [U]$
3. *If  $E$  is a good enough topological space, then*

$$X \subset Y \iff [X] \subset [Y]$$

**Lemma 53** (Subset to sublocal Part 2). *(Leroy Proposition 5.1.2, 5.1.3) For an open subspace  $U$  of  $E$  and a subspace  $X$  of  $E$ , we have:*

$$[U \cap X] = [U] \cap [X]$$

$$F = E \setminus U$$

$$[F] = [E]$$

$$[U]$$

$$[X \cap F] = [X] \cap [F]$$



**Lemma 54** (Part 3). *For any subspaces  $X$  of  $E$ , we have:*

1.

$$Ext[X] = [ExtX]$$

2.

$$[\bar{X}] = [\bar{X}]$$

3.

$$[IntX] \subset Int[X]$$

4.

$$\partial[X] \subset [Fr(X)]$$

*For a good enough topological space  $E$ , we have equality in 3 and 4.*

**Proposition 55** (Subset to sublocal preserves structure). *For two subspaces  $X$  and  $Y$  of  $E$  and an open subspaces  $U$  of  $E$ , we have:*

1.  $X \subset Y \implies [X] \subset [Y]$

2.  $X \subset U \iff [X] \subset [U]$

3. *If  $E$  is a good enough topological space, then*

$$X \subset Y \iff [X] \subset [Y]$$

4.

$$[U \cap X] = [U] \cap [X]$$

5. ...

**Theorem 56** (Measure top to loc). *Any measure on a good enough topological space  $X$  induces a measure on the corresponding local. Furthermore, the classical caratheodory extension onto  $\mathcal{P}(X)$  agrees with the restriction of the caratheodory extension of the induced measure on the local.*