

# ME 233 Advanced Control II

## Lecture 2 Introduction to Probability Theory

(ME233 Class Notes pp. PR1-PR3)

# Outline

- Sample Space and Events
- Probability function
- Discrete Random Variables
- Probability mass function, expectation and variance

# Sample Space and Events

Assume:

- We do an experiment many times.
  - Each time we do an experiment we call that a ***trial***
- The outcome of the experiment may be different at each trial.

$\omega_i$  : The  $i^{\text{th}}$  possible outcome of the experiment

# Sample Space and Events

**Sample Space  $\Omega$  :**

The space which contains all possible outcomes of an experiment.

$$\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$$

$\omega_i$  : The  $i^{\text{th}}$  possible outcome of the experiment

Each outcome is an element of  $\Omega$

## Example: Dice

## Experiment:

A situation whose ***outcome*** depends on chance

- throwing a die once



## Sample Space $\Omega$

The set of all possible ***outcomes*** of an experiment

$$\Omega = \left\{ \begin{array}{c} \square \\ \bullet \end{array}, \begin{array}{c} \square \\ \bullet \bullet \end{array}, \begin{array}{c} \square \\ \bullet \bullet \bullet \end{array}, \begin{array}{c} \square \\ \bullet \bullet \bullet \bullet \end{array}, \begin{array}{c} \square \\ \bullet \bullet \bullet \bullet \bullet \end{array}, \begin{array}{c} \square \\ \bullet \bullet \bullet \bullet \bullet \bullet \end{array} \right\}.$$

# Events

**Event**  $S_j$  :

Is a subset of the union of the sample space  $\Omega$   
and the empty set  $\phi$

If a sample space has  $n$  outcomes:

$$\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$$

There are  $2^n$  events:

$$\mathcal{S} = \{S_1, \dots, S_{2^n}\}$$

# Probability - events



Experiment: throwing a die once

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \right\}.$$

**Outcomes:** *elements of the sample space  $S$*

**Events:** *Are subsets of the sample space  $S$*

*An event occurs if any of the outcomes in that event occurs.*

*Empty subsets are **null** or **impossible events***

# Probability - events



Experiment: throwing a die once

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array} \right\}.$$

Some events:

- The event  $E$  of observing an even number of dots:

$$E = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array} \right\}$$

- The event  $O$  of observing an odd number of dots:


$$O = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \bullet \\ \hline \end{array} \right\}.$$



Example: throwing a pair of dice  
(one **red** and one **blue**)

- the sample space has **36** outcomes:

$\Omega =$



- The event  $L$  of obtaining the number **7** is


$$L = \{ (1,6), (2,5), (3,4), (4,3), (5,2), (6,1) \}$$

*$L$  occurs if any of the outcomes in  $L$  occurs.*

# Union, Complement and Intersection


- Union of two events:

$$S_i \cup S_j$$




- Intersection of two events:

$$S_i \cap S_j$$



- Complement of an event:


$$\setminus S_i = S_i^c$$



# Complement


- The **complement** of an event  $A$ , denoted by  $A^c$ , is the set of outcomes that are not in  $A$
- $A^c$  occurring means that  $A$  *does not occur*

$$A^c = \{\omega \mid \omega \in \Omega \text{ and } \omega \notin A\}$$



# Intersection of two events

- The **intersection** of two events ***A*** and ***B***, denoted by  **$A \cap B$** , is the set of outcomes that are in ***A***, **and** ***B***.
- *If the event  $A \cap B$  occurs, then **both** ***A*** and ***B*** occur*



- Events ***A*** and ***B*** are **mutually exclusive** if they cannot both occur at the same time, i.e. if

$$A \cap B = \emptyset$$

# Example of Intersection of two events



14

Experiment: throwing of a dice once

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \right\}.$$


- Events  $E$  and  $O$  are mutually exclusive

$$E = \left\{ \begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \right\} \quad O = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \right\}.$$

$$E \mid O = \blacklozenge$$

# Union of two events

- The **union** of two events  $\mathbf{A}$  and  $\mathbf{B}$ , denoted by  $\mathbf{A} \cup \mathbf{B}$ , is the set of outcomes that are in  $\mathbf{A}$ , or  $\mathbf{B}$ , or both
- *If the event  $\mathbf{A} \cup \mathbf{B}$  occurs, then either  $\mathbf{A}$  or  $\mathbf{B}$  or both occur*



# Probability function

We now consider the probability that a certain event occurs.

Recall: An event occurs if any of the outcomes in that event occurs.

The probability of event  **$A$**  will be denoted by

$$P(A)$$

# Probability

A number between 0 and 1, inclusive, that indicates **how likely an event is to occur.**

- An event with probability of 0 is a **null event.**
- An event with probability of 1 is a **certain event.**
- Probability of event  **$A$**  is denoted as  **$P(A)$** .
- The closer  **$P(A)$**  to 1, the more likely is  **$A$**  to happen.



# Intuitive Notion of Probability

The probability of event **A** is

$$P(A) = \frac{\text{Possible outcomes associated with } A}{\text{Total possible outcomes}}$$

(Assumes each outcome is equally likely)

$$0 \leq P(A) \leq 1$$

## Assigning Probability - Frequentist approach

- An experiment is repeated ***n*** times under essentially identical conditions
- if the event ***A*** occurs ***m*** times and ***n*** is large

$$P(A) \diamond \frac{m}{n}$$

# Dice example

Experiment: throwing a fair die once



$$\Omega = \{\square, \square, \square, \square, \square, \square\} \quad \Omega = \{1, 2, 3, 4, 5, 6\}$$

- $P(\Omega) = 1$
- $P(1) = 1/6, \quad P(3) = 1/6, \quad P(6) = 1/6$
- $P(\text{even number}) = 3/6 = 1/2$
- $P(\text{odd number}) = 3/6 = 1/2$

# Probability Space

The probability space is the triple:

$$(\Omega, \mathcal{S}, P)$$

Where

- $\Omega$  is the sample space
- $\mathcal{S}$  the set of all possible events
- $P : \mathcal{S} \rightarrow [0, 1]$  is the probability function

# Probability function

**Probability function:**  $P : \mathcal{S} \rightarrow [0, 1]$

Satisfies 3 axioms:

1.  $P(S_i) \geq 0, \quad \forall S_i \in \mathcal{S}$


2.  $P(\Omega) = 1$

3.  $P(S_i \cup S_j) = P(S_i) + P(S_j)$  if  $S_i \cap S_j = \emptyset$   
where  $S_i, S_j \in \mathcal{S}$

# Complement

- The **complement** of an event  $A$ , denoted by  $A^c$ , is the set of outcomes that are not in  $A$
- $A^c$  occurring means that  $A$  *does not occur*

$$A^c = \{\omega \mid \omega \in \Omega \text{ and } \omega \notin A\}$$



$$P(A^c) = 1 - P(A)$$

# Independent Events

- Two events are independent if


$$P(A \cap B) = P(A) \times P(B)$$

- Intuitively, two events are independent if the events do not influence each other:
  - Event **A** occurring does not affect the chances of **B** occurring, and vice versa.

# Example of independence

Experiment: throwing a pair of dice (one **red** and one **blue**)

$\Omega =$



*36 possible outcomes*

*The probability of throwing a **red** 1 and a **blue** 5 is*

$$P(\mathbf{1} \cap \mathbf{5}) = 1/36$$

$$= 1/6 \times 1/6 = P(\mathbf{1}) \times P(\mathbf{5})$$



# Law of Union

- Recall: If ***A*** and ***B*** are mutually exclusive

$$P(A \cup B) = P(A) + P(B)$$


- If ***A*** and ***B*** are not mutually exclusive

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

# Law of Union

- If  $A$  and  $B$  are not mutually exclusive


$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$



# Example

Experiment: throwing a pair of dice (one red and one blue)

$\Omega =$



- $P(L)$  = the probability of obtaining a 7

$$L = \{ (1,6), (2,5), (3,4), (4,3), (5,2), (6,1) \}$$

$$P(L) = P(1,6) + P(2,5) + P(3,4) + P(4,3) + P(5,2) + P(6,1)$$

$$P(L) = 6 / 36 = 1 / 6$$

# Joint Probability

Let ***A*** and ***B*** be two events

$$P(A \cap B)$$

is often called the ***joint probability*** of *A* and *B*

$$P(A)$$

$$P(B)$$

are often called the ***marginal probabilities*** of ***A*** and ***B***

# Conditional Probability

Let ***A*** and ***B*** be two events and  $P(B) \neq 0$

The conditional probability of event ***A*** given that event ***B*** has occurred is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

# Bayes' Rule

Let ***A*** and ***B*** be two events

$$\begin{aligned} P(A|B)P(B) &= P(B|A)P(A) \\ &= P(A \cap B) \end{aligned}$$

# Independence

The following are equivalent:

1. ***A*** and ***B*** are *independent*

$$2. P(A \cap B) = P(A) P(B)$$

$$3. P(A|B) = P(A)$$

$$4. P(B|A) = P(B)$$

# Array of Probabilities

Let  $C$  and  $D$  be two chance experiments.

Set of disjoint events associated with  $C$

$$\mathcal{C} = \{C_1, C_2, \dots, C_m\}$$

Set of disjoint events associated with  $D$

$$\mathcal{D} = \{D_1, D_2, \dots, D_n\}$$



# Array of Probabilities


We can construct:

$C \backslash D$	Event $D_1$	Event $D_2$	$\dots$	Event $D_n$	Marginal Probabilities
Event $C_1$	$P(C_1 \cap D_1)$	$P(C_1 \cap D_2)$	$\dots$	$P(C_1 \cap D_n)$	$P(C_1) = \sum_{i=1}^n P(C_1 \cap D_i)$
$\vdots$	$\vdots$	$\vdots$	$\dots$	$\vdots$	$\vdots$
Event $C_m$	$P(C_m \cap D_1)$	$P(C_m \cap D_1)$	$\dots$	$P(C_m \cap D_n)$	$P(C_m) = \sum_{i=1}^n P(C_m \cap D_i)$
Marginal Probabilities	$P(D_1) = \sum_{i=1}^m P(C_i \cap D_1)$	$P(D_2) = \sum_{i=1}^m P(C_i \cap D_2)$	$\dots$	$P(D_n) = \sum_{i=1}^m P(C_i \cap D_n)$	$Sum = 1$

# Example:

There are 4 balls in one jar, 2 balls are red and two balls are black.


- A person can remove a ball from the jar two times, without seeing the balls inside the jar.



# Example:

What is the probability of removing a red ball after having removing a red ball the first time?

To answer this question, lets build the table of probabilities.




# Example:

What is the probability of removing a red ball after having removing a red ball the first time?

To answer this question, lets build the table of probabilities.


Labels:



# Example:

Probability of picking  $R_1$  the first time?


$$P(R_1) = 1/4$$



# Example:

Probability of picking  $R_2$  with only 3 balls left?


$$P(R_2) = 1/3 \quad (\text{second time})$$



# Example:

Probability of picking  $R_1$  the first time and  $R_2$  the second time?

$$P(R_1 \cap R_2) = 1/4 \times 1/3 = 1/12$$



# Example: Array of Probabilities

<div> <div>2 pick</div> <div>1 pick</div> </div>	$R_1$	$R_2$	$B_1$	$B_2$	Marginal Probabilities
$R_1$	0	1/12	1/12	1/12	1/4
$R_2$	1/12	0	1/12	1/12	1/4
$B_1$	1/12	1/12	0	1/12	1/4
$B_2$	1/12	1/12	1/12	0	1/4
Marginal Probabilities	1/4	1/4	1/4	1/4	$Sum = 1$



# Probability of pick red balls consecutively

Probability of event  $A$ : picking a red ball the first time and a red ball the second time?

- Event B: Picking  $R_1$  first and  $R_2$  second
  - Event C: Picking  $R_2$  first and  $R_1$  second
- Mutually exclusive**  
*events*

$$P(A) = P(B \cup C)$$

$$= P(B) + P(C)$$

$$= \frac{1}{12} + \frac{1}{12} = \frac{1}{6}$$

# Example: Array of Probabilities

<div> <div>2 pick</div> <div>1 pick</div> </div>	<i>Red</i>	<i>Black</i>	Marginal Probabilities
<i>Red</i>	<i>1 / 6</i>	<i>1 / 3</i>	<i>1 / 2</i>
<i>Black</i>	<i>1 / 3</i>	<i>1 / 6</i>	<i>1 / 2</i>
Marginal Probabilities	<i>1 / 2</i>	<i>1 / 2</i>	<i>Sum = 1</i>

## Example:

What is the probability of picking a red ball the second time after having picked a red ball the first time?

$$P(Red_2|Red_1) = \frac{P(Red_2 \cap Red_1)}{P(Red_1)}$$

$$P(Red_2|Red_1) = \frac{1/6}{1/2} = \frac{1}{3}$$


# Discrete random variable

Given a sample Space  $\Omega$ , a random variable  $X$  is a function that assigns to each outcome a unique numerical value.

- Example: throwing of a die once



$$\Omega = \{ \square_{\cdot}, \square_{\cdot\cdot}, \square_{\cdot\cdot\cdot}, \square_{\cdot\cdot\cdot\cdot}, \square_{\cdot\cdot\cdot\cdot\cdot}, \square_{\cdot\cdot\cdot\cdot\cdot\cdot} \}.$$



# Discrete random variable

- Example: throwing of a die once



$$\Omega = \left\{ \boxed{\cdot}, \boxed{\cdot \cdot}, \boxed{\cdot \cdot \cdot}, \boxed{\cdot \cdot \cdot \cdot}, \boxed{\cdot \cdot \cdot \cdot \cdot}, \boxed{\cdot \cdot \cdot \cdot \cdot \cdot} \right\}.$$

- In this case, the random variable  $X$  only takes discrete values

$$x_i \in \{1, 2, 3, 4, 5, 6\}$$

- The discrete random variable  $X$  is defined by the **probability mass function**


$$P(x_i) = P(X = x_i)$$

*the probability that,  
after throwing a die,  
 $X$  will be equal to  $x_i$*

# Discrete random variable

- For a **fair die**, the probability mass function of the random variable  $X$  is

$$P(1) = P(2) = P(3) = P(4) = P(5) = P(6) = 1 / 6$$



*the probability mass function satisfies:*

$$\sum_{i=1}^6 P(x_i) = 1$$

# Expected value

- For a discrete random variable  $X$  taking on the  $N$  possible values

$$x_1, x_2, x_3, \dots, x_k, \dots, x_N$$

the **expected value** or **mean** of  $X$  is defined by

$$E[X] = m_x = \hat{x} = \sum_{k=1}^N x_k P(x_k)$$

$$E[X] = x_1 P(x_1) + x_2 P(x_2) + \dots + x_N P(x_N)$$

# Expected value of a function

- For a discrete random variable  $X$  taking on the  $N$  possible values

$$x_1, x_2, x_3, \dots, x_k, \dots, x_N$$

and the real-valued function  $f$

the **expected value** or **mean** of  $Y=f(X)$  is defined by

$$E[Y] = E[f(X)] = \sum_{k=1}^N f(x_k)P(x_k)$$

$$E[Y] = f(x_1)P(x_1) + \dots + f(x_N)P(x_N)$$



# Variance and standard deviation

- For a discrete random variable  $X$  taking on the  $N$  possible values

$x_1, x_2, x_3, \dots, x_k, \dots, x_N$  and a mean  $m_X = \hat{x}$

the **variance** of  $X$  is defined by

$$E[(X - m_X)^2] = \sigma_X^2 = \sum_{k=1}^N (x_k - m_X)^2 P(x_k)$$

where  $\sigma_X$  is the standard deviation of  $X$

# Cumulative Distribution Function

- The cumulative distribution function (CDF) for a discrete random variable  $X$  is

$$F_X(x) = P(X \leq x)$$

Find index  **$k$**  such that



$$x_k \leq x < x_{k+1}$$

$$F_X(x) = \sum_{j=1}^k P(x_j)$$

# Cumulative Distribution Function

- The cumulative distribution function (CDF) for a discrete random variable  $X$  is

$$F_X(x) = \sum_{j=1}^k P(x_j) \quad x_k \leq x < x_{k+1}$$























































































# Sum of two uniform independent random variables

- Let  $X$  and  $Y$  be 2 independent random variables with constant probability mass function
- Let  $Z = X + Y$
- The probability mass function of  $Z$  will not be constant

# Throwing two fair dice

Experiment: throwing a pair of fair dice (red and blue)





















































































$\Omega =$

- the sample space has **36** outcomes:
- each outcome has a **1/36** probability of occurring

# Throwing two fair dice

$\Omega =$


						
	 	 	 	 	 	 
	 	 	 	 	 	 
	 	 	 	 	 	 
	 	 	 	 	 	 
	 	 	 	 	 	 
	 	 	 	 	 	 

- Define the random variable  $Z$  associated with the **event** of observing the total number of dots on both dice after each throw

$Z = k$  when the throw results in the number  $k$

# Throwing two fair dice

$\Omega =$



*number of  
outcomes*


36

$Z$  only takes discrete values

$Z_i \blacklozenge \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$

# Throwing two fair dice

$\Omega =$



*probability of  
each outcome  
 $1/36$*

*we now estimate:*

$$Z=2 \spadesuit P(2) = 1/36$$

$$Z=7 \spadesuit P(7) = 6/36$$

$$Z=3 \spadesuit P(3) = 2/36$$

















































$$Z=12 \spadesuit P(12) = 1/36$$

$$Z=4 \spadesuit P(4) = 3/36$$



# Throwing two fair dice

$\Omega =$

The **probability mass function** is

















































$$P(2) = 1/36 \quad P(5) = 4/36 \quad P(8) = 5/36 \quad P(11) = 2/36$$


$$P(3) = 2/36 \quad P(6) = 5/36 \quad P(9) = 4/36 \quad P(12) = 1/36$$

$$P(4) = 3/36 \quad P(7) = 6/36 \quad P(10) = 3/36$$

# Throwing two fair dice

$\Omega =$



*the probability mass function satisfies:*

$$\sum_{k=2}^{12} P(k) = 1$$