

**Due 01/30/22 11:59 pm**

- Homework 0 consists of both written and coding questions.
- We prefer that you typeset your answers using  $\text{\LaTeX}$  or other word processing software. If you haven't yet learned  $\text{\LaTeX}$ , one of the crown jewels of computer science, now is a good time! Neatly handwritten and scanned solutions will also be accepted for the written questions.
- In all of the questions, **show your work**, not just the final answer.
- **Start early. This is a long assignment. Most of the material is prerequisite material not covered in lecture; you are responsible for finding resources to understand it.**

**Deliverables:**

1. Submit a PDF of your homework to the Gradescope assignment entitled "HW0 Write-Up". **Please start each question on a new page.** If there are graphs, include those graphs in the correct sections. **Do not** put them in an appendix. We need each solution to be self-contained on pages of its own.
  - In your write-up, please state with whom you worked on the homework. This should be on its own page and should be the first page that you submit.
  - In your write-up, please copy the following statement and sign your signature next to it. (Mac Preview and FoxIt PDF Reader, among others, have tools to let you sign a PDF file.) We want to make it *extra* clear so that no one inadvertently cheats. *"I certify that all solutions are entirely in my own words and that I have not looked at another student's solutions. I have given credit to all external sources I consulted."*
  - **Replicate all your code in an appendix.** Begin code for each coding question in a fresh page. Do not put code from multiple questions in the same page. When you upload this PDF on Gradescope, *make sure* that you assign the relevant pages of your code from appendix to correct questions.

## 1 Administrivia (2 points)

1. Please fill out the Welcome Survey if you haven't already. Please write down the email address you used for this survey for reference
2. When are the midterm and final?
3. How many homeworks are there this semester? How many homework drops does each student get?
4. Are you allowed to use online resources to help solve homework problems?
5. What is the penalty for cheating in CS 189?
6. What topic(s) are you most excited to learn about in this course?

## 2 Probability Potpourri (11 points)

1. (2 points) Recall the covariance of two random variables  $X$  and  $Y$  is defined as  $\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$ . For a multivariate random variable  $Z$  (i.e., each index of  $Z$  is a random variable), we define the covariance matrix  $\Sigma$  such that  $\Sigma_{ij} = \text{Cov}(Z_i, Z_j)$ . Concisely,  $\Sigma = \mathbb{E}[(Z - \mu)(Z - \mu)^\top]$ , where  $\mu$  is the mean value of the random column vector  $Z$ . Prove that the covariance matrix is always positive semidefinite (PSD).

*Hint:* Use linearity of expectation.

2. (4 points) The probability that an archer hits her target when it is windy is 0.4; when it is not windy, her probability of hitting the target is 0.7. On any shot, the probability of a gust of wind is 0.3. Find the probability that
  - (i) on a given shot there is a gust of wind and she hits her target.
  - (ii) she hits the target with her first shot.
  - (iii) she hits the target exactly once in two shots.
  - (iv) there was no gust of wind on an occasion when she missed.
3. (2 points) An archery target is made of 3 concentric circles of radii  $1/\sqrt{3}$ , 1 and  $\sqrt{3}$  feet. Arrows striking within the inner circle are awarded 4 points, arrows within the middle ring are awarded 3 points, and arrows within the outer ring are awarded 2 points. Shots outside the target are awarded 0 points.

Consider a random variable  $X$ , the distance of the strike from the center (in feet), and let the probability density function of  $X$  be

$$f(x) = \begin{cases} \frac{2}{\pi(1+x^2)} & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

What is the expected value of the score of a single strike?

4. (3 points) Let  $X \sim \text{Pois}(\lambda)$ ,  $Y \sim \text{Pois}(\mu)$ . given that  $X \perp\!\!\!\perp Y$ , derive an expression for  $\mathbb{P}(X | X + Y = n)$ . What well-known probability distribution is this? What are its parameters?

### 3 Linear Algebra Review (10 points)

1. (3 points) Let  $A \in \mathbb{R}^{n \times n}$  be a symmetric matrix. Prove equivalence between these three different definitions of positive semidefiniteness (PSD).
  - (a) For all  $x \in \mathbb{R}^n$ ,  $x^\top A x \geq 0$ .
  - (b) All the eigenvalues of  $A$  are nonnegative.
  - (c) There exists a matrix  $U \in \mathbb{R}^{n \times n}$  such that  $A = U U^\top$ .

Mathematically, we write positive semidefiniteness as  $A \geq 0$ .

2. (5 points) Now that we're equipped with different definitions of positive semidefiniteness, use them to prove the following properties of PSD matrices.
  - (a) If  $A$  and  $B$  are PSD, then  $2A + 3B$  is PSD.
  - (b) If  $A$  is PSD, all diagonal entries of  $A$  are nonnegative:  $A_{ii} \geq 0, \forall i \in [n]$ .
  - (c) If  $A$  is PSD, the sum of all entries of  $A$  is nonnegative:  $\sum_{j=1}^n \sum_{i=1}^n A_{ij} \geq 0$ .
  - (d) If  $A$  and  $B$  are PSD, then  $\text{Tr}(AB) \geq 0$ , where  $\text{Tr}(M)$  denotes the *trace* of  $M$ .
  - (e) If  $A$  and  $B$  are PSD, then  $\text{Tr}(AB) = 0$  if and only if  $AB = 0$ .
3. (2 points) Let  $A \in \mathbb{R}^{n \times n}$  be a symmetric, PSD matrix. Write  $\|A\|_F$  as a function of the eigenvalues of  $A$ .

*Hint:* Recall that  $\|A\|_F = \sqrt{\text{Tr}(A^\top A)}$ . If you haven't seen this before, you should try to prove it. However, you can accept this as a given fact for this homework assignment.

## 4 Gradients and Norms (10 points)

1. (3 points) Define the  $\ell_p$ -norm as  $\|x\|_p = \left(\sum_{i=1}^n |x_i|^p\right)^{1/p}$ , where  $x \in \mathbb{R}^n$ .  
Prove that  $\frac{1}{\sqrt{n}}\|x\|_2 \leq \|x\|_\infty \leq \|x\|_1 \leq \sqrt{n}\|x\|_2$ . The Cauchy–Schwarz inequality is helpful here.
2. (2 points) Consider a linear decision function  $f(x) = w \cdot x + \alpha$  and the hyperplane decision boundary  $H = \{x : w \cdot x = -\alpha\}$ . Prove that if  $w$  is a unit vector, then the *signed distance* (the  $\ell_2$ -norm distance with an appropriate sign) from  $x$  to the closest point on  $H$  is  $w \cdot x + \alpha$ .
3. (3 points) Consider the function which maps a vector to its maximum entry,  $x \mapsto \max_i x_i$ . While this function is non-smooth, a common trick in machine learning is to use a smooth approximation, *LogSumExp*, defined as follows.

$$\text{LSE} : \mathbb{R}^n \rightarrow \mathbb{R}, \text{LSE}(x) = \ln \left( \sum_{i=1}^n e^{x_i} \right).$$

One of the nice properties of this function is that it is convex, which can be proved by showing its Hessian matrix is positive semidefinite. To that end, compute its gradient and Hessian. You do not need to prove that the Hessian is PSD.

4. (2 points) Let  $X \in \mathbb{R}^{n \times d}$  be a data matrix, consisting of  $n$  samples, each of which has  $d$  features, and let  $y \in \mathbb{R}^n$  be a vector of outcomes. We wish to find the *best linear approximation*, i.e. we want to find the  $\theta$  that minimizes the loss  $L(\theta) = \|y - X\theta\|_2^2$ . Assuming  $X$  has full column rank, compute  $\theta^* = \operatorname{argmin}_\theta L(\theta)$  in terms of  $X$  and  $y$ .

## 5 Isocontours of Normal Distributions (8 points)

Let  $f(\mu, \Sigma)$  be the probability density function of a normally distributed random variable in  $\mathbb{R}^2$ . Write code to plot the isocontours of the following functions, each on its own separate figure. You are free to use Matplotlib, NumPy, and SciPy.

(a) (2 points)  $f(\mu, \Sigma)$ , where  $\mu = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ .

(b) (1 point)  $f(\mu, \Sigma)$ , where  $\mu = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$  and  $\Sigma = \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix}$ .

(c) (2 points)  $f(\mu_1, \Sigma_1) - f(\mu_2, \Sigma_2)$ , where  $\mu_1 = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$ ,  $\mu_2 = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$  and  $\Sigma_1 = \Sigma_2 = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$ .

(d) (2 points)  $f(\mu_1, \Sigma_1) - f(\mu_2, \Sigma_2)$ , where  $\mu_1 = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$ ,  $\mu_2 = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$ ,  $\Sigma_1 = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$  and  $\Sigma_2 = \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix}$ .

(e) (1 point)  $f(\mu_1, \Sigma_1) - f(\mu_2, \Sigma_2)$ , where  $\mu_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mu_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$ ,  $\Sigma_1 = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$  and  $\Sigma_2 = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ .

## 6 Eigenvectors of the Gaussian Covariance Matrix (7 points)

Consider two one-dimensional random variables  $X_1 \sim \mathcal{N}(3, 9)$  and  $X_2 \sim \frac{1}{2}X_1 + \mathcal{N}(4, 4)$ , where  $\mathcal{N}(\mu, \sigma^2)$  is a Gaussian distribution with mean  $\mu$  and variance  $\sigma^2$ . Write a program that draws  $n = 100$  random two-dimensional sample points from  $(X_1, X_2)$  such that the  $i$ th value sampled from  $X_2$  is calculated based on the  $i$ th value sampled from  $X_1$ . In your code, make sure to choose and set a fixed random number seed for whatever random number generator you use, so your simulation is reproducible, and document your choice of random number seed and random number generator in your write-up. For each of the following parts, include the corresponding output of your program.

- (a) (1 point) Compute the mean (in  $\mathbb{R}^2$ ) of the sample.
- (b) (1 point) Compute the  $2 \times 2$  covariance matrix of the sample.
- (c) (2 points) Compute the eigenvectors and eigenvalues of this covariance matrix.
- (d) (1 point) On a two-dimensional grid with a horizontal axis for  $X_1$  with range  $[-15, 15]$  and a vertical axis for  $X_2$  with range  $[-15, 15]$ , plot
  - (i) all  $n = 100$  data points, and
  - (ii) arrows representing both covariance eigenvectors. The eigenvector arrows should originate at the mean and have magnitudes equal to their corresponding eigenvalues.
- (e) (2 points) Let  $U = [v_1 \ v_2]$  be a  $2 \times 2$  matrix whose columns are the eigenvectors of the covariance matrix, where  $v_1$  is the eigenvector with the larger eigenvalue. We use  $U^\top$  as a rotation matrix to rotate each sample point from the  $(X_1, X_2)$  coordinate system to a coordinate system aligned with the eigenvectors. (As  $U^\top = U^{-1}$ , the matrix  $U$  reverses this rotation, moving back from the eigenvector coordinate system to the original coordinate system). *Center* your sample points by subtracting the mean  $\mu$  from each point; then rotate each point by  $U^\top$ , giving  $x_{\text{rotated}} = U^\top(x - \mu)$ . Plot these rotated points on a new two dimensional-grid, again with both axes having range  $[-15, 15]$ .

In your plots, **clearly label the axes and include a title**. Moreover, **make sure the horizontal and vertical axis have the same scale!** The aspect ratio should be one. You are free to use Matplotlib and NumPy.