

Non-linear least squares method for Newton Gravity law using Gauss-Newton method

Berlet Maxim 3384

December 9, 2025

1 Newton's Gravity law (ODE)

Newton's Second law

$$\vec{F} = m\vec{a} \Rightarrow \frac{d^2x}{dt^2} = \frac{\vec{F}}{m}$$

Newton's Gravity law

$$\vec{F} = \frac{GMm}{r^2} \frac{\vec{r}}{r} = \frac{GMm}{r^3} \vec{r} \Rightarrow \frac{d^2x}{dt^2} = \frac{GM}{r^3} \vec{r},$$

where $x = \vec{r}(t) = (x(t), y(t), z(t))$

$$\frac{d^2\vec{r}}{dt^2} = \frac{GM}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} (x \ y \ z)$$

Split vector into scalar equations

$$\begin{cases} \frac{d^2x}{dt^2} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} x \\ \frac{d^2y}{dt^2} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} y \\ \frac{d^2z}{dt^2} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} z \end{cases}$$

Make system first order ODE

$$\begin{cases} \frac{dv_x}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} x \\ \frac{dv_y}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} y \\ \frac{dv_z}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}} z \\ \frac{dx}{dt} = v_x \\ \frac{dy}{dt} = v_y \\ \frac{dz}{dt} = v_z \end{cases}$$

And for solving Cauchy problem add initial values:

$$v(t_0) = (v_{x_0}, v_{y_0}, v_{z_0});$$

$$r(t_0) = (x_0, y_0, z_0).$$

2 Gauss-Newton method

Regression model

$$y_i = g(x(t_i), q) + \varepsilon_i$$

Where:

- y_i - row of observed values in t_i moment,
- $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$ - row of errors in measurements,
- g - nonlinear function that returns row of model values,
- $x(t_i)$ - system state vector in moment t_i
- q - reduction parameters (determines how we measure values)

Dynamical System equation

$$\begin{cases} \frac{dx}{dt} = F(x(t), p) \\ x(t_0) = x_0 \end{cases}$$

Where:

- $x(t)$ - system state in time t
- p - dynamical parameters (e.g. GM, other constants)

Residuals and regression parameters

$$\begin{aligned} r_i &= y_i - g(x(t_i), q) \\ \beta &= (x_0 \ p \ q) \end{aligned}$$

Parameters optimization using least squares

$$\begin{aligned} \sum_{i=1}^N r_i(\hat{\beta})^2 &\rightarrow \min \\ \hat{\beta}^{(s+1)} &= \hat{\beta}^{(s)} - (J^T W J)^{-1} J^T W r(\hat{\beta}^{(s)}) \end{aligned}$$

Where:

- J - Jacobian of residuals ($J_{ij} = \frac{\partial r_i}{\partial \beta_j}$)
- W - weight matrix ($W = \text{diag}(\frac{1}{\sigma_1^2}, \dots, \frac{1}{\sigma_N^2})$)

$$\frac{\partial r_i}{\partial \beta_j} = \left(\frac{\partial r_i}{\partial x_{0j}} \quad \frac{\partial r_i}{\partial p_j} \quad \frac{\partial r_i}{\partial q_j} \right)$$

$$\begin{aligned} \frac{\partial r_i}{\partial q_j} &= - \frac{\partial g(x(t_i), q)}{\partial q_j} \\ \frac{\partial r_i}{\partial p_j} &= - \frac{\partial g(x(t_i), q)}{\partial x} \frac{\partial x}{\partial p_j} \\ \frac{\partial r_i}{\partial x_{0j}} &= - \frac{\partial g(x(t_i), q)}{\partial x} \frac{\partial x}{\partial x_{0j}} \end{aligned}$$

Let's combine last two equations into one matrix $P \equiv (x_0 \ p) \Rightarrow Q(t) = \frac{\partial x}{\partial P}$

Additional Dynamical equation

$$\frac{d}{dt} \frac{\partial x}{\partial P} = \frac{\partial F}{\partial P} + \frac{\partial F}{\partial x} \frac{\partial x}{\partial P} \Rightarrow \frac{d}{dt} Q(t) = \frac{\partial F}{\partial P} + \frac{\partial F}{\partial x} Q(t)$$

$$Q(t_0) = \begin{bmatrix} 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 & \cdots & 0 \end{bmatrix}$$

$\underbrace{\hspace{10em}}$
 x_0

$\underbrace{\hspace{10em}}$
 p

3 Newton law in Gauss-Newton method for Orbit correction

Dynamical system (ODE)

$$\begin{cases} F_1 = \frac{dv_x}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}}x \\ F_2 = \frac{dv_y}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}}y \\ F_3 = \frac{dv_z}{dt} = \frac{GM}{(x^2+y^2+z^2)^{\frac{3}{2}}}z \\ F_4 = \frac{dx}{dt} = v_x \\ F_5 = \frac{dy}{dt} = v_y \\ F_6 = \frac{dz}{dt} = v_z \\ v(t_0) = (v_{x_0}, v_{y_0}, v_{z_0}) \\ r(t_0) = (x_0, y_0, z_0) \end{cases}$$

In this case we have (x, y, z) as observed values therefore our nonlinear function $g(x(t)) = (x(t) \ y(t) \ z(t))$. Lets redefine state vector in function input as $\vec{s}(t) = (x(t) \ y(t) \ z(t) \ v_x(t) \ v_y(t) \ v_z(t))$ from state $\Rightarrow g(s(t)) = (x(t) \ y(t) \ z(t)) = \vec{r}(t)$ We don't have any p or q parameters so differentiation much easier.

$$\frac{\partial r_i}{\partial s_{0_j}} = -\frac{\partial g(s(t_i))}{\partial s} \frac{\partial s}{\partial s_{0_j}} = \begin{bmatrix} \frac{\partial x}{\partial x} & \frac{\partial x}{\partial y} & \frac{\partial x}{\partial z} & \frac{\partial x}{\partial v_x} & \frac{\partial x}{\partial v_y} & \frac{\partial x}{\partial v_z} \\ \frac{\partial y}{\partial x} & \frac{\partial y}{\partial y} & \frac{\partial y}{\partial z} & \frac{\partial y}{\partial v_x} & \frac{\partial y}{\partial v_y} & \frac{\partial y}{\partial v_z} \\ \frac{\partial z}{\partial x} & \frac{\partial z}{\partial y} & \frac{\partial z}{\partial z} & \frac{\partial z}{\partial v_x} & \frac{\partial z}{\partial v_y} & \frac{\partial z}{\partial v_z} \end{bmatrix} \frac{\partial s}{\partial s_{0_j}} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \end{bmatrix} \frac{\partial s}{\partial s_{0_j}}$$

And now to find last component $Q(t) = \frac{\partial s}{\partial s_{0_j}}$ we need to integrate one more equation

$$\begin{aligned} \frac{d}{dt}Q(t) &= \frac{\partial F}{\partial s_0} + \frac{\partial F}{\partial s}Q(t) \\ Q(t_0) &= \underbrace{\begin{bmatrix} 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 & \cdots & 0 \end{bmatrix}}_{s_0} \underbrace{\begin{bmatrix} \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix}}_p = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

In that case first term equals 0 and we have simplified equation

$$\frac{d}{dt}Q(t) = \frac{\partial F}{\partial s}Q(t)$$

Lets find out what's $\frac{\partial F}{\partial s}$ equals to.

$$\begin{aligned}
& \frac{\partial F}{\partial s} = \begin{bmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_1}{\partial y} & \frac{\partial F_1}{\partial z} & \frac{\partial F_1}{\partial v_x} & \frac{\partial F_1}{\partial v_y} & \frac{\partial F_1}{\partial v_z} \\ \frac{\partial F_2}{\partial x} & \frac{\partial F_2}{\partial y} & \frac{\partial F_2}{\partial z} & \frac{\partial F_2}{\partial v_x} & \frac{\partial F_2}{\partial v_y} & \frac{\partial F_2}{\partial v_z} \\ \frac{\partial F_3}{\partial x} & \frac{\partial F_3}{\partial y} & \frac{\partial F_3}{\partial z} & \frac{\partial F_3}{\partial v_x} & \frac{\partial F_3}{\partial v_y} & \frac{\partial F_3}{\partial v_z} \\ \frac{\partial F_4}{\partial x} & \frac{\partial F_4}{\partial y} & \frac{\partial F_4}{\partial z} & \frac{\partial F_4}{\partial v_x} & \frac{\partial F_4}{\partial v_y} & \frac{\partial F_4}{\partial v_z} \\ \frac{\partial F_5}{\partial x} & \frac{\partial F_5}{\partial y} & \frac{\partial F_5}{\partial z} & \frac{\partial F_5}{\partial v_x} & \frac{\partial F_5}{\partial v_y} & \frac{\partial F_5}{\partial v_z} \\ \frac{\partial F_6}{\partial x} & \frac{\partial F_6}{\partial y} & \frac{\partial F_6}{\partial z} & \frac{\partial F_6}{\partial v_x} & \frac{\partial F_6}{\partial v_y} & \frac{\partial F_6}{\partial v_z} \end{bmatrix} = \\
& = \begin{bmatrix} \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-2x^2+y^2+z^2) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3xy) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3xz) & 0 & 0 & 0 \\ \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3yx) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(x^2-2y^2+z^2) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3yz) & 0 & 0 & 0 \\ \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3zx) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(-3zy) & \frac{GM}{(x^2+y^2+z^2)^{\frac{5}{2}}}(x^2+y^2-2z^2) & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \\
& \frac{\partial v_x}{\partial x} = 0 \quad \text{same for } y, z
\end{aligned}$$