Sampling and Estimation - Exercise 2

Jan-Philipp Kolb, Stefan Zins and Matthias Sand
2 Februar 2016

Exercise 2.A

Estimation under a stratified design

- Download the ESS for Sweden and Denmark (round 5)
- Import data to R and combine the two datasets
- Define a survey object (stratified design)
- Estimate the empirical distribution of tv consumption (tvtot) in Sweden and Denmark
- Estimate the joined empirical distribution of tv consumption in Sweden and Denmark

Codebook ESS round 5

Exercise 2.B

• Load the survey package and the api datasets.

library(survey) data(api)

• The dataset apistrat is a sample of schools from apipop stratified by stype. Assuming the selection within the strata was done by SRS, define a survey object (svydesign) and calculate a point and variance estimate for the mean of api00.

Exercise 2.B

• Using stype again as a stratification variable try different allocations for stratified sample. Calculate the allocation of a sample of 60 schools from apipop using equal, proportional and optimal allocation. The proportional allocation should be proportional to the number of schools within the strata and the optimal allocation should be optimal with regard to api99.

Exercise 2.B

- Select a StrSRS from apipop for each allocation.
- Estimate again the mean of api00 from all three samples and compare the results.

Function for stratified samples

Getting the function

source(url)