Noise bends a polynomial fit

Ada Lovelace (Analytical Engine Institute)

Abstract

We measure how noise bends a polynomial fit, and find that its error grows as the square of the noise.

Introduction

State of the art

Polynomial fits are standard[1–3]. Noise is known to bias them[4]. An earlier draft said so. A lost work claimed it first [to be resolved], and a late one agreed[5]. A third said it again.

Results

How does noise change the fit?

The error grows with the noise

The error of the fit follows

$$e(\sigma) = 0.8\,\sigma^2 + 0.02 \tag{1}$$

and Eq. 1 holds at every noise level we ran, as Fig. 1 shows. Table 1 lists the errors. The fit follows Smith's method[1].

Table 1: The error of the fit at each noise level.

noise	rmse
0.1	0.03
0.2	0.05
0.4	0.15

Figure 1: The error of the fit grows with the noise[4].

Discussion

Limits

The noise model is Gaussian, as Lee assumed[4]. A heavier tail would bend the fit further, and the next runs should try one.

Methods

Fitting

We fit by least squares, as Jones did[2].

Acknowledgments

[TO BE GIVEN: acknowledgments]

Funding

[TO BE GIVEN: funding]

Competing interests

The authors declare none.

Data availability

The data are synthetic and the project's runs made them.

Code availability

[TO BE GIVEN: the code repository, which setup has not settled]

References

1. Smith, A. & Li, B. Polynomial fits in practice. *Journal of Fits* https://doi.org/10.5555/smith (2020) doi:10.5555/smith.

2. Jones, C. Least squares for curves. *Numerical Notes* https://doi.org/10.5555/jones (2019) doi:10.5555/jones.
3. Kim, F. A survey of fitting. *Fitting Reviews* https://doi.org/10.5555/kim (2017) doi:10.5555/kim.
4. Lee, D. & Hall, E. Noise and bias in fits. *arXiv* Preprint at https://arxiv.org/abs/1801.00001 (2018).
5. Late, A. A late agreement. *Journal of Tests* https://doi.org/10.5555/late (2024) doi:10.5555/late.