

Real Analysis

Additional Solution to Chapter 1

Xinyuan Niu*

*This file is hereby granted for private, non-commercial and educational purpose only. If you find any flaws, please contact me at bernardniu97@gmail.com

Exercise1 Prove that the Cantor set \mathcal{C} constructed in the text is totally disconnected and perfect.

Solution

Consider the construction of Cantor set. $\forall x, y \in \mathcal{C}$, we can find $k \in \mathbb{N}$ while x and y belong to the same interval in C_k but two different intervals in C_{k+1} . W.L.O.G. note $x, y \in [a_s, b_s]$ which is preserved after k_{th} operation with $x \in [a_s, \frac{2a_s+b_s}{3}]$ and $y \in [\frac{a_s+2b_s}{3}, b_s]$, thus $(\frac{2a_s+b_s}{3}, \frac{a_s+2b_s}{3})$ will be deleted in the next operation. Then it is obvious that $\exists z \in (\frac{2a_s+b_s}{3}, \frac{a_s+2b_s}{3})$ s.t. $x < z < y$ and $z \notin \mathcal{C}$, which means that the Cantor set is totally disconnected.

the other property is also apparently with the two following facts:

1. The length of each interval in C_k is $\frac{1}{3^k}$;
2. The end points of each interval in C_k all belong to \mathcal{C} .

$\forall x \in \mathcal{C}$, we note that $x_n \neq x$ is one of the two end points of the interval in C_k which includes x . Then $\{x_n\}_{n=0}^{\infty} \subset \mathcal{C}$ and $|x - x_k| \leq \frac{1}{3^k}$. So x is not an isolated point, that equally proves the Cantor set is perfect since we've already figured out that \mathcal{C} is closed.

Exercise10 Let $\hat{\mathcal{C}}$ denote a Cantor-like set. Let F_i denote a certain piecewise-linear and continuous function on $[0,1]$ which satisfies:

- a) $F_i = 0$ at the center of all intervals in stage i of the construction of $\hat{\mathcal{C}}$;
- b) $F_i = 1$ in the complement of the intervals in stage i of the construction of $\hat{\mathcal{C}}$;
- c) $F_i(x)$ is a linear function in the rest of $[0,1]$.

Let $f_n = \prod_{i=1}^n F_i$, Prove the following:

- (a) (We see the first part of (a) as kind of a hint) $f_n(x)$ converges to a limit as $n \rightarrow \infty$ which we denote by $f(x)$.
- (b) f is discontinuous at every point of $\hat{\mathcal{C}}$.

Solution

- (a) According to the construction of $F_i(x)$, we can easily notice that $F_i(x) \in [0, 1]$ for $\forall x \in [0, 1]$. Then it can be deduced that $f_i(x) \in [0, 1]$ and $f_{i+1}(x) \leq f_i(x)$ for $\forall x \in [0, 1]$. So $\{f_i(x)\}$ is monotonous and bounded, thus convergent.
- (b) First we point out that $f(x) = 0, \forall x \in \hat{\mathcal{C}}$. Since $\sum_{n=1}^{\infty} 2^{k-1} l_k < 1$, $\lim_{n \rightarrow \infty} l_k = 0$. Then $\forall \epsilon > 0, \exists N \in \mathbb{N}^*$ s.t. after k_{th} operation, the length of each remaining interval, $l_k < \epsilon$. For $\forall x \in \hat{\mathcal{C}}$, denote the interval, left after k_{th} operation, which includes x , I_k . It is obvious that the midpoint of I_k , referred to as x_k , doesn't belong to $\hat{\mathcal{C}}$ with $f(x_k) = 0$ and $|x - x_k| < \epsilon$. Now $\{x_n\}$ converges to x , but $f(x_n) \equiv 0$ with $f(x) = 1$. So $f(x)$ isn't continuous at any point in $\hat{\mathcal{C}}$.

Exercise12 Prove the following:

- (a) An open disc in \mathbb{R}^2 is not the disjoint union of open rectangles.
- (b) An open connected set Ω is the disjoint union of open rectangles if and only if Ω itself is an open rectangle.

Solution

- (a) W.L.O.G. the center of the disc is exactly at the origin. Since all the open rectangles are disjoint, so any point on the boundary of any rectangle must be not in the disc. If there is a point on the boundary of a certain rectangle R_k which is in $\mathbb{R}^2 \setminus (\partial R_k \cup R_k)$, apparently there will be a open set $T \subset R_k$, the distance between whom and the origin is larger than the radii r , which is contradictory to the definition. It indicates that all the bounded points of those rectangles are just on the boundary of the disc, so all of the bounded points share the equal normal, it is obviously incorrect.
- (b) We only need to prove *only if*. Here we prove it with contradiction. Presume there is an open connected set \mathcal{A} matching conditions, however, not being an open rectangle. Then \mathcal{A} is capable of being depicted as the union of no less than two disjointed open rectangles, thus it can be written as two disjointed open sets, this doesn't match the fact that \mathcal{A} is connected.

Exercise14 The **outer Jordan content** $J_*(E)$ of a set of E in \mathbb{R} is defined by

$$J_*(E) = \inf \sum_{j=1}^N |I_j|$$

where the inf is taken over every *finite* covering $E \subset \cup_{j=1}^N I_j$, by intervals I_j .

(a) Prove that $J_*(E) = J_*(\bar{E})$ for every set E (here \bar{E} denotes the closure of E).

(b) Exhibit a countable subset $E \subset [0, 1]$ such that $J_*(E) = 1$ while $m_*(E) = 0$

kaixinjiuhao