

# Graph optimization project

Luca Ferraro (10748116), Bernardo Camajori Tedeschini (10584438),  
William Colombo (10537311)

[https://github.com/LucaFerraro/Graph\\_optimization\\_project](https://github.com/LucaFerraro/Graph_optimization_project)

A.Y. 2020 - 2021

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Problem formulation and continuous relaxation</b>	<b>3</b>
2.1	Variables for ILP formulation . . . . .	3
2.2	ILP formulation . . . . .	3
2.3	Continuous relaxation . . . . .	4
<b>3</b>	<b>Lagrangian relaxation</b>	<b>5</b>
<b>4</b>	<b>Valid inequalities and cutting planes</b>	<b>6</b>
<b>5</b>	<b>Randomized rounding</b>	<b>7</b>
<b>6</b>	<b>Greedy</b>	<b>8</b>
<b>7</b>	<b>Local search - K-opt neighborhood</b>	<b>9</b>
<b>8</b>	<b>Results</b>	<b>10</b>

## List of Figures

<b>1</b>	<b>Problem description</b> . . . . .	<b>2</b>
----------	--------------------------------------	----------

# 1 Introduction

The aim of this project is solving an integer linear programming (ILP) problem by using different strategies.

The problem that we have to solve is the one described in Figure 1.

Consider a directed graph  $G = (N, A)$ , where each node can host a facility. A set of requests for treatment  $K$  is given: each request  $k$  originates in a node  $o_k \in N$  and has a demand for treatment  $d_k$ . The whole demand of  $k$  must be treated by one and only one facility, and a single path must be selected from the origin node  $o_k$  to the facility to which  $k$  is assigned.

Opening a facility in node  $i \in N$  costs  $c_i$  and a facility opened in  $i$  can treat an overall demand  $cap_i$ .

Each arc of the graph has a limited capacity  $uu$ . Arc capacity is the same for all the arcs. For each demand that uses arc  $(i, j)$  a cost  $g_{ij}$  must be paid.

Many requests can originate in the same node but they can be assigned to different facilities and can be routed on different paths even if they are assigned to the same facility.

The problem consists in deciding where to open the facilities, to which facility each request is assigned, and the routing on a single path of each request, with the aim of minimizing the sum of the costs.

Figure 1: Problem description

In the following sections, we describe all the methods that we have implemented to solve this problem.

## 2 Problem formulation and continuous relaxation

In this section, we provide the ILP formulation of the problem and its continuous relaxation.

### 2.1 Variables for ILP formulation

The first thing we have to decide is which variables we need. Since we have to take three decisions, we need 3 variables.

1.  $y_i \in \{0, 1\}$ : this is the location variable, which assumes value 1 if a facility is installed in node  $i \in N$  and 0 otherwise.
2.  $z_{k,i} \in \{0, 1\}$ : this is the assignment variable, and is = 1 if the demand  $k \in K$  is served by node  $i \in N$  and 0 otherwise.
3.  $x_{i,j}^k \in \{0, 1\}$ : this is the routing variable, and it assumes value = 1 if the demand  $k \in K$  uses the arc  $(i,j) \in A$  and 0 otherwise.

### 2.2 ILP formulation

Here, we provide the ILP formulation of the problem.

In the formulation, the flow conservation constraint (the third one) has been written as the difference between the flow entering a node and the one exiting the same node so that at the second member of the constraint we can use directly the variable  $z_{k,i}$  to distinguish between the cases in which the difference has to be 0 or 1. Indeed, if  $z_{k,i}=0$ , then node  $i$  is not serving demand  $k$ , that is the node is a transit node for the demand  $k$  and therefore the difference has to be 0; vide versa, if  $z_{k,i}=0$ , then node  $i$  is serving demand  $k$ , that is the node is the destination node for the demand  $k$  and therefore the difference has to be 1.

$$\begin{aligned}
 & \text{minimize} && \sum_{i \in N} c_i y_i + \sum_{k \in K} \sum_{(i,j) \in A} g_{i,j} x_{i,j}^k \\
 & \text{subject to} && \sum_{k \in K} d_k x_{i,j}^k \leq u u && \forall (i,j) \in A, \\
 & && \sum_{k \in K} d_k z_{k,i} \leq y_i cap_i && \forall i \in N, \\
 & && \sum_{(j,i) \in A} x_{j,i}^k - \sum_{(i,j) \in A} x_{i,j}^k = \begin{cases} -1, & i = o_k \\ z_{k,i}, & \text{otherwise} \end{cases} && \forall i \in N, k \in K, \\
 & && y_i \in \{0, 1\} && \forall i \in N, \\
 & && z_{k,i} \in \{0, 1\} && \forall k \in K, i \in N, \\
 & && x_{i,j}^k \in \{0, 1\} && \forall (i,j) \in A, k \in K
 \end{aligned}$$

### 2.3 Continuous relaxation

The formulation for the CR of the problem is the same as for the ILP. The only difference is in the domain of the variables that now are not binary but continuous and can assume all values in [0,1]:

$$\begin{aligned}
& \text{minimize} && \sum_{i \in N} c_i y_i + \sum_{k \in K} \sum_{(i,j) \in A} g_{i,j} x_{i,j}^k \\
& \text{subject to} && \sum_{k \in K} d_k x_{i,j}^k \leq uu \quad \forall (i,j) \in A, \\
& && \sum_{k \in K} d_k z_{k,i} \leq y_i cap_i \quad \forall i \in N, \\
& && \sum_{(j,i) \in A} x_{j,i}^k - \sum_{(i,j) \in A} x_{i,j}^k = \begin{cases} -1, & i = o_k \\ z_{k,i}, & \text{otherwise} \end{cases} \quad \forall i \in N, k \in K, \\
& && 0 \leq y_i \leq 1 \quad \forall i \in N, \\
& && 0 \leq z_{k,i} \leq 1 \quad \forall k \in K, i \in N, \\
& && 0 \leq x_{i,j}^k \leq 1 \quad \forall (i,j) \in A, k \in K
\end{aligned}$$

### 3 Lagrangian relaxation

We can use the Lagrangian relaxation (LR) since we have agreed that the integrality property does not hold, so we can in theory obtain a lower bound that is better than the one found with the continuous relaxation.

The constraint we have chosen to relax is the node capacity constraint (the second one in formulation in Section 2): since we have one of those constraint for each  $i \in N$ , the constraint is inserted in the objective function through multipliers  $\lambda_i$ .

The new formulation then is:

$$\begin{aligned} & \text{minimize} && \sum_{i \in N} c_i y_i + \sum_{k \in K} \sum_{(i,j) \in A} g_{i,j} x_{i,j}^k - \sum_{i \in N} \lambda_i (y_i cap_i - \sum_{k \in K} d_k z_{k,i}) \\ & \text{subject to} && \sum_{k \in K} d_k x_{i,j}^k \leq uu \quad \forall (i,j) \in A, \\ & && \sum_{(j,i) \in A} x_{j,i}^k - \sum_{(i,j) \in A} x_{i,j}^k = \begin{cases} -1, & i = o_k \\ z_{k,i}, & \text{otherwise} \end{cases} \quad \forall i \in N, k \in K, \\ & && y_i \in \{0, 1\} \quad \forall i \in N, \\ & && z_{k,i} \in \{0, 1\} \quad \forall k \in K, i \in N, \\ & && x_{i,j}^k \in \{0, 1\} \quad \forall (i,j) \in A, k \in K \end{aligned}$$

After many attempts, we haven't found a good bound for the problem: we have tried different starting points for the multipliers  $\lambda_i$  and many different ways of updating the step (both with and without the normalization) of the sub-gradient, but we always find a lower bound that is even worse than the one provided by the continuous relaxation.

For this reason we have tried to change strategy and relax the arc capacity constraint, thus obtaining the following relaxation:

$$\begin{aligned} & \text{minimize} && \sum_{i \in N} c_i y_i + \sum_{k \in K} \sum_{(i,j) \in A} g_{i,j} x_{i,j}^k - \sum_{(i,j) \in A} \lambda_{i,j} (uu - \sum_{k \in K} d_k z_{k,i}) \\ & \text{subject to} && \sum_{k \in K} d_k z_{k,i} \leq y_i cap_i \quad \forall i \in N, \\ & && \sum_{(j,i) \in A} x_{j,i}^k - \sum_{(i,j) \in A} x_{i,j}^k = \begin{cases} -1, & i = o_k \\ z_{k,i}, & \text{otherwise} \end{cases} \quad \forall i \in N, k \in K, \\ & && y_i \in \{0, 1\} \quad \forall i \in N, \\ & && z_{k,i} \in \{0, 1\} \quad \forall k \in K, i \in N, \\ & && x_{i,j}^k \in \{0, 1\} \quad \forall (i,j) \in A, k \in K \end{aligned}$$

This relaxation works much better than the previous one.

## 4 Valid inequalities and cutting planes

The idea we have had for the valid inequalities is to consider each arc capacity constraint as if it was a knapsack problem (so we are considering a knapsack problem for each arc):

- The capacity of each knapsack is  $uu$ , that is the capacity of each arc.
- The items we have to put in the knapsack are represented by the routing variables  $x_{i,j}^k$ : the arc  $(i,j)$  is fixed and we have to choose the requests considering that the weight of the  $k^{th}$  request is  $d_k$

So, what we do is defining a cover, that is a subset  $S \subseteq K$  of elements (flow in this case) such that:

$$\sum_{k \in S} x_{i,j}^k > uu \quad (1)$$

Therefore, we can write the cover inequality as:

$$\sum_{k \in S} x_{i,j}^k \leq |S| - 1 \Rightarrow \sum_{k \in S} (1 - x_{i,j}^k) \geq 1 \quad (2)$$

The separation problem consists then in finding a cover that violates the cover inequality. This is done by solving the following optimization problem for each arc:

$$\begin{aligned} & \text{minimize} && \sum_{k \in K} u_k (1 - x_k^*) \\ & \text{subject to} && \sum_{k \in K} d_k u_k \geq uu + 1, \\ & && u_k \in \{0, 1\} \forall k \in K \end{aligned}$$

Here,  $x_k^*$  is the solution of the current continuous relaxation.  
At each iteration, we try to solve the separation problem for all the arcs, but as soon as we add a cut, we stop and go to the following iteration.  
This is repeated or for a maximum number of iterations or until we don't find any other cut to add.

## 5 Randomized rounding

For applying the randomized rounding approach we consider the continuous relaxation of the problem as described in Section 2 but with the addition of a more explicit consistency constraint relating variables  $z_{k,i}$  and  $y_i$ :

$$z_{k,i} \leq y_i, \forall k \in K, i \in N \quad (3)$$

By adding this constraint, we can apply the randomized rounding procedure only two times, one to the  $z_{k,i}$  variables and then to the  $x_{i,j}^k$  variables. In fact, the solver will automatically fix the value for the variables  $y_i$  in dependance of the values assigned to the  $z_{k,i}$  by the randomized rounding.

## 6 Greedy

The idea of our greedy is looking at one demand at a time and assign try to assign it to a facility that is already opened and that can serve the demand. Of course, at the beginning there is no opened facility, so we have to decide which one to open; similarly, once a facility Our decision is to open the facilities in non-decreasing order of opening cost.

The scheme for the greedy then is:

1. For each demand  $k \in K$ , check if there is a facility that is opened and that can serve the current demand:
  - If yes, compute the shortest path to the first found facility by keeping into account the arc capacity (that is added to the shortest path problem as a constraint) and assign the demand to that facility by routing it on the path found.
  - If no, open as new facility the one whose opening cost is minimum, assign the demand to this facility and route the demand to it on the shortest path (again, keeping into account the arc capacity).

Note that with this algorithm we are able to find a feasible solution in a number of iterations that is at most equal to the number of the demands

## 7 Local search - K-opt neighborhood

We have decided to implement a K-optimum neighborhood.

The initial solution for the algorithm is the one computed by our greedy algorithm.

All what we have to do is writing the ILP formulation of the problem with the addition of the K-opt constraint

$$\sum_{i \in N - Y} y_i + \sum_{i \in Y} (1 - y_i) + \sum_{(i, j, k) \in (A \times D) - X} x_{i, j, k} + \sum_{(i, j, k) \in X} (1 - x_{i, j, k}) + \sum_{(k, i) \in (D \times N) - Z} z_{k, i} + \sum_{(k, i) \in Z} (1 - z_{k, i}) \leq K \quad (4)$$

Note that here, the ensembles X, Y and Z are the ones containing respectively the variables  $x_{i, j, k}$ ,  $y_i$  and  $z_{k, i}$  that are = 1 from the greedy.

For the value of K, we have made many attempts and we have set it equal to the number of variables /50.

## 8 Results

In this section, we report the table with all the results we have found. As it can be seen, there are some results (the ones marked with (\*)) that are not correct: they are about the heuristic and the local search approach for a particular instance and they are providing a value that is lower than the optimum found by the ILP even if they should provide an upper bound of the optimum. This is due to the fact that the ILP we use for finding the shortest path within the greedy doesn't find a feasible solution.

instance	integer optimum	time	continuous relaxation (CR)	gap	(CR)with added inequalities	alternative relaxation	time	randomized rounding	time	greedy	time	LS	time
ABR	777	5.69	722.505	0.070	722.981	777	7.71	777	11.66	1356	0.81	777	45.64
AB1	401	0.35	394.676	0.016	394.958	401	0.39	401	0.64	1091	0.78	401	4.32
ATR	927	8.93	761.74	0.178	761.742	915	100.8	927	6.15	1632	2.13	927	54.92
ATBP	30	1.49	20.6196	0.313	20.62	30	28.27	30	6199.8	30	2.56	30	4.4
NORR	329	5471.7	327.592	0.004	327.592	328	580.89	329	5407	349	44.99	349	8.81
NORROUT	10596	3.9	10556	0.003	10556	10350	162.35	10596	8.05	30510	44.51	10596	33.38
NORB <sub>P</sub>	11	2581.5	9.96	0.094	9.96	10	43.92	14	5304.8	10*	33.89	10*	5.98
NORB <sub>P</sub> 2	4	170.99	3.33	0.167	3.33	4	41.26	55	1946.7	4	38.58	4	91.21
NORROUT1	1022	22.63	705.333	0.31	705.333	1022	40.98	1022	56.12	2129	32.28	1022	195.05
NORROUT2	3014	48.67	1447.33	0.52	1447.33	3014	72.04	3014	107.27	4796	32.27	3014	6221.5