The group G is isomorphic to the group PSL(2,11): C2. Ordinary character table of $G \cong PSL(2,11)$: C2:

	1a	2a	3a	5a	5b	6a	11a	2b	4a	10a	10b	12a	12b
χ_1	1	1	1	1	1	1	1	1	1	1	1	1	1
χ_2	1	1	1	1	1	1	1	-1	-1	-1	-1	-1	-1
<i>χ</i> ₃	10	2	-2	0	0	2	-1	0	0	0	0	0	0
χ_4	10	-2	1	0	0	1	-1	0	2	0	0	-1	-1
χ_5	10	-2	1	0	0	1	-1	0	-2	0	0	1	1
χ_6	10	2	1	0	0	-1	-1	0	0	0	0	$-E(12)^7 + E(12)^{11}$	$E(12)^7 - E(12)^{11}$
χ_7	10	2	1	0	0	-1	-1	0	0	0	0	$E(12)^7 - E(12)^{11}$	$-E(12)^7 + E(12)^{11}$
χ_8	11	-1	-1	1	1	-1	0	1	-1	1	1	-1	-1
χ_9	11	-1	-1	1	1	-1	0	-1	1	-1	-1	1	1
χ_{10}	12	0	0	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	1	2	0	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	0
χ_{11}	12	0	0	$E(5) + E(5)^4$	$E(5)^2 + E(5)^3$	0	1	-2	0	$-E(5) - E(5)^4$	$-E(5)^2 - E(5)^3$	0	0
χ_{12}	12	0	0	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	1	2	0	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	0
χ_{13}	12	0	0	$E(5)^2 + E(5)^3$	$E(5) + E(5)^4$	0	1	-2	0	$-E(5)^2 - E(5)^3$	$-E(5) - E(5)^4$	0	0

Trivial source character table of $G \cong PSL(2,11)$: C2 at p = 5:

Normalisers N_i		N_1								N_2			
p-subgroups of G up to conjugacy in G		P_1								P_2			
Representatives $n_j \in N_i$	1a	2b	2a	3a	4a	6a	12a	12b	11a	1a	2c	2b	2a
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 1 \cdot \chi_{10} + 0 \cdot \chi_{11} + 1 \cdot \chi_{12} + 0 \cdot \chi_{13}$	25	5	1	1	1	1	1	1	3	0	0	0	0
		5	-1	-1	-1	-1	-1	-1	2	0	0	0	0
$ 0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 1 \cdot \chi_{11} + 0 \cdot \chi_{12} + 1 \cdot \chi_{13} $		-5	1	1	-1	1	-1	-1	3	0	0	0	0
		-5	-1	-1	1	-1	1	1	2	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 1 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13}$		0	2	-2	0	2	0	0	-1	0	0	0	0
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 1 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13}$		0	-2	1	2	1	-1	-1	-1	0	0	0	0
$ 0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 1 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} $	10	0	-2	1	-2	_	1	1	-1	0	0	0	0
$ 0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 1 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} $		0	2	1	0		$-E(12)^7 + E(12)^{11}$	$E(12)^7 - E(12)^{11}$	-1	0	0	0	0
$ 0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 1 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} $	10	0	2	1	0	-1	$E(12)^7 - E(12)^{11}$	$-E(12)^7 + E(12)^{11}$	-1	0	0	0	0
$1 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13}$	1	1	1	1	1	1	1	1	1	1	1	1	1
	11	-1	-1	-1	1	-1	1	1	0	1	-1	1	-1
$ 0 \cdot \chi_1 + 1 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 0 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13} $	1	-1	1	1	-1	1	-1	-1	1	1	1	-1	-1
$0 \cdot \chi_1 + 0 \cdot \chi_2 + 0 \cdot \chi_3 + 0 \cdot \chi_4 + 0 \cdot \chi_5 + 0 \cdot \chi_6 + 0 \cdot \chi_7 + 1 \cdot \chi_8 + 0 \cdot \chi_9 + 0 \cdot \chi_{10} + 0 \cdot \chi_{11} + 0 \cdot \chi_{12} + 0 \cdot \chi_{13}$		1	-1	-1	-1	-1	-1	-1	0	1	-1	-1	1

 $P_1 = Group([()]) \cong 1$ $P_2 = Group([(3, 10, 8, 7, 11)(4, 9, 6, 5, 12)]) \cong C5$

 $N_1 = Group([(3,4)(5,7)(6,8)(9,10)(11,12),(1,2,3)(4,5,6)(7,9,10)(8,11,12)]) \cong PSL(2,11) : C2$ $N_2 = Group([(3,10,8,7,11)(4,9,6,5,12),(1,2)(5,6)(7,8)(9,12)(10,11),(1,2)(3,4)(5,8)(6,7)(9,11)(10,12)]) \cong D20$