



© 2000 by The McGraw-Hill Companies, Inc. All rights reserved. This publication is protected by copyright. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without prior written permission from The McGraw-Hill Companies, Inc. This publication may be used for educational or promotional purposes upon written request to The McGraw-Hill Companies, Inc. For more information, contact The McGraw-Hill Companies, Inc., 1221 Avenue of the Americas, New York, NY 10020-1095.

© 2005 Blackwell Publishing Ltd, *Journal of Internal Medicine* 258: 103–110

Another important factor affecting the results of the simulation study is the type of the distribution of the input data. In the present study, the input data were generated from a normal distribution. However, in many practical situations, the input data may follow other distributions, such as the log-normal distribution, the gamma distribution, the beta distribution, etc. Therefore, it is necessary to investigate the effect of the distribution of the input data on the results of the simulation study. This is a topic for future research.

[illegible]

(1) The Commission shall, in accordance with the provisions of the Treaty, ensure that the Commission's work is carried out in a manner which is consistent with the principles of the Commission's work.

[illegible]

© 2004 Blackwell Publishing Ltd, *Journal of Internal Medicine* 255: 103–110

