Midterm presentation

The Quadrocopters

Technische Universität München

26. Mai 2015

Overview

- Motivation
- Model
- Realtime Optimization Approach
- Results

Optimal Control Problem

$$\min_{x,u} J(x,u) \qquad \dot{x} = f(x,u)$$


x : state

u : control

 \rightarrow additional difficulty: realtime approach

Model

Forces and Torques




Newton-Euler Equations

Forces


$$F_{\text{ext}} = F_g + \sum_{i=1}^4 F_i$$

Torques

$$\tau_{\mathsf{ext}} = \sum_{i=1}^{4} \tau_i + (\tau_{\varphi} + \tau_{\theta})$$



Coordinate Systems



$$q = a + ib + jc + kd$$
 $a, b, c, d \in \mathbb{R}$

$$q = a + ib + jc + kd$$
 $a, b, c, d \in \mathbb{R}$

representing rotation $\Leftrightarrow \|q\| = 1$

$$q = a + ib + jc + kd$$
 $a, b, c, d \in \mathbb{R}$

representing rotation $\Leftrightarrow \|q\| = 1$ advantage \to no singularities

$$q = a + ib + jc + kd$$
 $a, b, c, d \in \mathbb{R}$

representing rotation $\Leftrightarrow \|q\|=1$ advantage \to no singularities problem $\to \|q\|=1$ additional constraint

Dynamics

Equations representing dynamics...

$$T(x, u) = M \cdot \begin{pmatrix} \dot{x}_8 \\ \vdots \\ \dot{x}_{13} \end{pmatrix} + \Theta(x)$$

Dynamics

Equations representing dynamics...

$$T(x, u) = M \cdot \begin{pmatrix} \dot{x}_8 \\ \vdots \\ \dot{x}_{13} \end{pmatrix} + \Theta(x)$$

...expressed as system of differential equations:


$$\frac{d}{dt} \begin{pmatrix} x_1 \\ \vdots \\ x_7 \\ x_8 \\ \vdots \\ x_{13} \end{pmatrix} = \begin{pmatrix} \dot{x}_1 \\ \vdots \\ \dot{x}_7 \\ M^{-1}(T(x, u) - \Theta(x)) \end{pmatrix}$$


Prospect


Refinement of the model


- \rightarrow wind
- $\rightarrow \text{ aerodynamical forces}$

Realtime Optimization Approach









Minimization Problem

$$\min_{\substack{s_t, ..., s_N \\ q_t, ..., q_{N-1}}} \sum_{i=t}^{N-1} J_i(s_i, q_i) \quad s.t. \quad \begin{cases} x_t - s_t = 0 \\ h_i(s_i, q_i) - s_{i+1} = 0 \quad \forall i = t, ..., N-1 \end{cases}$$

$$J_i(s_i, q_i)$$
 discretized goal function $x_t - s_t = 0$ expected state = real state $h_i(s_i, q_i)$ solution of the ODE at time i

The Lagrangian

$$L^{t}(y) = \sum_{i=t}^{N-1} J_{i}(s_{i}, q_{i}) + \lambda_{t}^{T}(x_{t} - s_{t}) + \sum_{i=t}^{N-1} \lambda_{i+1}^{T}(h_{i}(s_{i}, q_{i}) - s_{i+1})$$

$$y := (\lambda_t, s_t, q_t, \lambda_{t+1}, s_{t+1}, q_{t+1}, ..., \lambda_N, s_N)$$

We are looking for y^* satisfying the KKT conditions:

$$\Rightarrow \nabla_y L^t(y^*) = 0$$

The SQP Method

How to find y^* ?

$$\begin{aligned} y_{k+1} &= y_k + \alpha_k \Delta y_k \\ \min_{\Delta y_k} &= \frac{1}{2} \Delta y_k^T A_k \Delta y_k + \nabla_{y_k} J(y_k)^T \Delta y_k \end{aligned}$$

The SQP Method

How to find y^* ?

$$y_{k+1} = y_k + \alpha_k \Delta y_k$$

$$\min_{\Delta y_k} = \frac{1}{2} \Delta y_k^T A_k \Delta y_k + \nabla_{y_k} J(y_k)^T \Delta y_k$$

Special case:

$$A_k = H(y_k)$$
 approximated Hessian $\nabla^2_{y_k} L(y_k)$ $\alpha_k = 1$

Newton-Raphson

Find Δy_k with:

$$\nabla_{y_k} L(y_k) + H(y_k) \Delta y_k = 0$$

Newton-Raphson

Find Δy_k with:

$$\nabla_{y_k} L(y_k) + H(y_k) \Delta y_k = 0$$

There is only time for 1 SQP step:

$$y_{t+1}=y_1=y_0+\Delta y_0$$

Riccati Recursion

Approximated Hessian:

What happens in interval $[t_{k-1}, t_k]$?



What happens in interval $[t_{k-1}, t_k]$?



• calculate control u_{k-1} (Riccati Part II)

What happens in interval $[t_{k-1}, t_k]$?



- calculate control u_{k-1} (Riccati Part II)

What happens in interval $[t_{k-1}, t_k]$?



- calculate control u_{k-1} (Riccati Part II)
- \circ calculate y_k (Riccati Part II)

Finite Horizon

$$\min_{\substack{s_t, ..., s_N \\ q_t, ..., q_{N-1}}} \sum_{i=t}^{N-1} J_i(s_i, q_i) \quad s.t. \quad \begin{cases} x_t - s_t = 0 \\ h_i(s_i, q_i) - s_{i+1} = 0 \quad \forall i = t, ..., N-1 \end{cases}$$

How to choose N?

$$N=t_{end} \longrightarrow {\sf problem \ size \ decreasing}$$

Finite Horizon

$$\min_{\substack{s_t, ..., s_N \\ q_t, ..., q_{N-1}}} \sum_{i=t}^{N-1} J_i(s_i, q_i) \quad s.t. \quad \begin{cases} x_t - s_t = 0 \\ h_i(s_i, q_i) - s_{i+1} = 0 \quad \forall i = t, ..., N-1 \end{cases}$$

How to choose N?

$$N=t_{end}
ightarrow {
m problem}$$
 size decreasing $N=t+n
ightarrow {
m problem}$ size constant

Results

Questions?

S. Boyd.
Solving the lqr problem by block elimination.
Lecture notes. 2009.

J. Diebel.

Representing attitude: Euler angles, unit quaternions, and rotation vectors.

10 2006.

M. Diehl, B. H. Georg, J. P. Schlder, R. Findeisen, Z. Nagy, and F. Allgwer.

Real-time optimization and nonlinear model predictive control of processes governed by differential-algebraic equations.

Journal of Process Control, 2002.

M. Diehl, H. G. Bock, and J. P. Schlder.

A real-time iteration scheme for nonlinear optimization in optimal feedback control.

SIAM J. Control Optim., 2005.

- M. M. Diehl.

 Real-Time Optimization for Large Scale Nonlinear Processes.

 PhD thesis, Ruprecht-Karls-Universiteidelberg, 2001.
- L. R. Garcia Carrillo, A. E. Dzul Lopez, R. Lozano, and C. Pegard. Quad Rotorcraft Control. Springer-Verlag London, 2013.
- D. Hartmann, K. Landis, M. Mehrer, S. Moreno, and J. Kim. Quadcopter dynamic modeling and simulation (quad-sim) v1.00. Git, 2014.

URL https://github.com/dch33/Quad-Sim.

E. Reyes-Valeria, R. Enriquez-Caldera, S. Camacho-Lara, and J. Guichard.

Lqr control for a quadrotor using unit quaternions: Modeling and simulation.

IEEE Xplore, 2013.



J. Richter-Gebert and T. Orendt.

Geometriekalkle.

Springer: Berlin, Heidelberg, 2009.