Aguiar, A., Deans, A., Engel, M., Forshage, M., Huber, J., Jennings, J., Johnson, J., Arkady S. Lelej, A., Longino, J., Lohrmann, V., Mikó, I., Michael Ohl, M., Rasmussen, C., Taeger, A. & D. (2013). Order Hymenoptera. In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). Zootaxa 3703: 51-62.

Ahyong, S.T., Lowry, J.K., Alonso, M., Bamber, R.N., Boxshall, G.A., Castro, P., Gerken, S., Karaman, G.S., Goy, J.W., Jones, D.S., Meland, K., Rogers, D.C. & Svavarsson, J. (2011). Subphylum Crustacea Brünnich, 1772. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline c higher-level classification and survey of taxonomic richness. Zootaxa 3148: 165–191.

Allen, Robert T. (2002). A synopsis of the Diplura of North America: Keys to higher taxa, systematics, distributions and descriptions of new taxa (Arthropoda: Insecta), Transactions of the American Entomological Society 128(4): 403-466.

Anderson, O.R. & Cavalier-Smith, T. (2012). Ultrastructure of Diplophrys parva, a new small freshwater species, and a revised analysis of Labrinthulea (Heterokonta). Acta Protozool. 51: 291-304.

Balsamo, M., Guidi, L. & D'hondt, J-L. (2013). Phylum Gastrotricha. In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richn (Addenda 2013). Zootaxa 3703: 79-82.

Bamber, R. (2011). Class Pycnogonida Latreille, 1810. In: Zhang, Z.-Q. (Ed.), Animal biodivers An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 110–1

Beccaloni, G. & Eggleton, P. (2011). Order Blattodea Brunner von Wattenwyl, 1882. In: Zhanş Z.-Q. (Ed.) Animal biodiversity: An outline of

higher-level classification and survey of taxonomic richness. Zootaxa 3148: 199-200.

Beaulieu, F., Dowling, A., Klompen, H., Moraes, G., & Walter, D. (2011). Superorder Parasitiformes Reuter, 1909. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-l classification and survey of taxonomic richness. Zootaxa 3148: 123-128.

Ben-Dov, Y., Miller, D.R. & Gibson, G.A.P. (2012) SCALENET. Accessed at: http://www.sel.barc.usda.gov/scalenet/classif.htm

Betancur-R., R., Wiley, E.O., Miya, M., Lecointre, G., Bailly, N. & Ortí, G. 2013. New and Revised Classification of Bony Fishes Version 2. Retrieved from http://deepfin.org/Classification\_v2.htm

Bieler, R., Carter, J., & Coan, E. 2010. Classification of Bivalve Families. In: Bouchet, P., Ro J-P., Bieler, R., Carter., J., & Coan, E. (2010). Nomenclator of bivalve families and classificatio bivalve families, Malacologia 52(2): 113-133.

Blackburn, D.C. & Wake, D.B. (2011). Class Amphibia Gray, 1825. In: Zhang, Z.-Q. (Ed.), Anibiodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 39–55.

Bock, P. (2014). Belluloporina. In: Bock, P.; Gordon, D. (2014) World List of Bryozoa. Retrieve from: World Register of Marine Species at http://www.marinespecies.org

Bock, P. (Ed.). (2014). Systematic List of Families of Bryozoa. Accessed at: http://bryozoa.net/famsys.html

Bock, P. & Gordon, D. (2013). Phylum Bryozoa Ehrenburg, 1831. In: Zhang, Z.-Q. (Ed.), Anir Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Ad. 2013). Zootaxa 3703: 67-74.

Bouchet, P. (2014a). Crosseolidae Hickman, 2013. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Bouchet, P. (2014b). Globocormidae Espinosa & Ortez, 2010. Retrieved from: World Register a Marine Species at http://www.marinespecies.org

Bouchet, P. (2014c). Eoacmaeoidea Nakano & Ozawa, 2007. Retrieved from: World Register c Marine Species at http://www.marinespecies.org

Bouchet, P. & Gofas, S. (2014a). Caudofoveata. Retrieved from: World Register of Marine Spec at http://www.marinespecies.org

Bouchet, P. & Gofas, S. (2014b). Neomphaloidea McLean, 1981. Retrieved from: World Regists of Marine Species at http://www.marinespecies.org

Bouchet, P., Rocroi, J., Fryda, J., Hausdorf, B., Ponder, W., Valdes, A., & Waren, A. (2005). Classification and nomenclator of gastropod families. Malacologia, 47(1-2): 1-397.

Bourgoin, T. (2014). FLOW - Fulgoromorpha Lists On the Web. Accessesd at: http://www.hemiptera-databases.org/flow

Brands, S. J. (comp.). (2014) The Taxonomicon. Universal Taxonomic Services, Zwaag, The Netherlands. [http://taxonomicon.taxonomy.nl/].

Bray, R., & Cribb, T. (2012). Reorganisation of the superfamily Lepocreadioidea Odhner, 1905 based on an inferred molecular phylogeny. Systematic Parasitology 83 (3):169-177.

Bray, R., Gibson, D., & Jones, A. (Eds.) (2008). Keys to the Trematoda: Volume 3. CAB International, London.

Cavalier-Smith, T. (1997). Amoeboflagellates and mitochondrial cristae in eukaryote evolution: megasystematics of the new protozoan subkingdoms Eozoa and Neozoa. Arch Protistenkd 147, 237–258

Cavalier-Smith, T. (2013). Early evolution of eukaryote feeding modes, cell structural diversity, classification of the protozoan phyla Loukozoa, Sulcozoa, and Choanozoa. European Journal of Protistology 49: 115-178.

Cavalier-Smith, T., & Chao, E. (2004). Protalveolate phylogeny and systematics and the original Sporozoa and dinoflagellates (phylum Myzozoa nom. nov.). European Journal of Protistology 4 185-212.

Cavalier-Smith, T., & Chao, E.E.-Y. (2006). Phylogeny and Megasystematics of Phagotrophic Heterokonts (Kingdom Chromista). Journal of Molecular Evolution 62(4): 388-420.

Cavalier-Smith, T., & Chao, E.E.-Y. (2012). Oxnerella micra sp. n. (Oxnerellidae fam. n.), a Tin Naked Centrohelid, and the Diversity and Evolution of Heliozoa. Protist, 163: 574–601.

Cavalier-Smith, T., Lewis, R., Chao, E., Oates, B., & Bass, D. (2008). Morphology and phyloge of Sainouron acronematica sp. n. and the ultrastructural unity of Cercozoa. Protist, 159(4): 591-6

Cavalier-Smith, T. & Scoble, J.M. (2012). Phylogeny of Heterokonta: Incisomonas marina, a uniciliate gliding opalozoan related to Solenicola (Nanomonadea), and evidence that Actinophyr evolved from raphidophytes. European Journal of Protistology, doi: 10.1016/l.ejop.2012.09.002

Chantangsi, C. & Leander, B. (2010). An SSU rDNA barcoding approach to the diversity of marine interstitial cercozoans, including descriptions of four novel genera and nine novel species International Journal of Systematic and Evolutionary Microbiology 60: 1962-1977.

Choi, E.J., Beatty, D.S., Paul, L.A., Fenical, W. & Jensen, P.R. (2013). Mooreia alkaloidigena gen. nov., sp. nov. and Catalinimonas alkaloidigena gen. nov., sp. nov., alkaloid-producing marii bacteria in the proposed families Mooreiaceae fam. nov. and Catalimonadaceae fam. nov. in the phylum Bacteroidetes. Int. J. Syst. Evol. Microbiol., 63:1219 - 1228.

Christenhusz, M.J.M., Zhang, X.-C. & Schneider, H. (2011). A linear sequence of extant familie and genera of lycophytes and ferns. Phytotaxa 19: 7-54.

Christofferson, M. (2012). Phylogeny of basal descendants of cocoon-forming annelids (Clitell Turk. J. Zool. 36(1): 95-119.

Collins, A.G. (2009). Recent insights into cnidarian phylogeny. Smithsonian Contributions to the Marine Sciences 38: 139–148.

Crandall-Stotler, B., Stotler, R., & Long, D. (2009). Morphology and classification of the Marchantiophyta. In Goffinet, B. & Shaw, A.J. Bryophyte Biology, 2nd Edition. Cambridge University Press, Cambridge UK, pp.1–54.

Condé, B., and J. Pagés. 1991. Diplura. pp. 269-271. In CSIRO (ed.). The Insects of Australia A Textbook for Students and Research Workers. 2nd edition. Melbourne University Press, Carl Australia. 2 volumes. xvii + 1137 + vi pp.

Cribb, T. & Gibson, D. (2014). Digenea. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Crowther, A.L. (2011). Class Anthozoa Ehrenberg, 1834. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 19–23.

Cryan, J. & Urban, J. (2011). Higher-level phylogeny of the insect order Hemiptera: is Auchenorrhyncha really paraphyletic? Systematic Entomology (2011), DOI: 10.1111/j.1365-3113.2011.00611.x

Cutler, E.B. (1994). The Sipuncula: Their Systematics, Biology, and Evolution. Cornell Univers Ithaca, New York, 480 pp.

Daly, M., Brugler, M.R., Cartwright, P., Collins, A.G., Dawson, M.N., Fautin, D.G., France, S.C Catherine, P., McFadden, S., Opresko, D.M., Rodriguez, E., Romano, S.L. & Stake, J.L. (2007). The phylum Cnidaria: A review of phylogenetic patterns and diversity 300 years after Linnaeus. Zootaxa 1668: 127–182.

de Reviers, B., Rousseau, F., & Draisma, S. (2007). Classification of the Phaeophyceae from pato presentand current challenges. In Brodie, J. & Lewis, J. (Eds). Unravelling the Algae: The Pas Present and Future, CRC Press, Florida, pp. 267-284.

Dijkstra, K-D., Bechly, G., Bybee, S., Dow, R., Dumont, H., Fleck, G., Garrison, R., Hämäläinen M., Kalkman, V., Karube, H., May, M., Orr, A., Paulson, D., Rehn, A., Theischinger, G., Truem J., Van Tol, J. Von Ellenrieder, N. & Ware, J. (2013). The classification and diversity of dragor and damselflies (Odonata). In: Zhang, Z.-Q. (Ed.), Animal Biodiversity: An Outline of Higher-Classification and Survey of Taxonomic Richness (Addenda 2013). Zootaxa 3703: 36-45.

Emig, C. (2011). Phylum Phoronida Hatschek, 1888. In: Zhang, Z.-Q. (Ed.), Animal biodiversity outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 230.

Emig, C., Alvarez, F., & Bitner, M. (2013). Brachiopoda World Database. Available at http://www.marinespecies.org/brachiopoda

Emschermann, P. (1972). Loxokalypus socialis gen. et sp. nov. (Kamptozoa, Loxokalypodidae fi nov.), ein neuer Kamptozoentyp aus dem nördlichen Pazifischen Ozean. Ein Vorschlag zur Neufassung der Kamptozoensystematik. Marine Biology 12: 237-254.

Engel, M. & Haas, F. (2007). Family-Group Names for Earwigs (Dermaptera). American Mus Novitates, Number 3567, 20 pp.

Erséus, C. (2005). Phylogeny of Oligochaetous Clitellata. Hydrobiologia 535/536: 357-372.

Erséus, C., Wetzel, M., & Gustavsson, L. (2008). ICZN Rules -- A farewell to Tubificidae. Zootaxa 1744: 66-68.

Eschmeyer, W.N. & Fong, J.D. (2011). Pisces. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 26–38.

Eschmeyer, W.N. & Fong, J.D. (2013). SPECIES BY FAMILY/SUBFAMILY. Accessed at http://research.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp

Euzéby, J. & Whitman, W.B. (2011). Validation List 143, Family I. Rhodothermaceae fam. nov Int. J. Syst. Evol. Microbiol.: 62, 1-4.

Fletcher, M.J. (2009). Key to the Leafhoppers and Treehoppers of Australia and neighbouring areas. Accessed at: http://www1.dpi.nsw.gov.au/keys/leafhop/index.html

Fritz, U. & Havas, P. (2013). Order Testudines: 2013 update. In: Zhang, Z.-Q. (Ed.), Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Ad. 2013). Zootaxa 3703: 12-14.

Garcia-Alvarez, O. & Gofas, S. (2013). Solenogastres. Accessed through World Register of Mar Species at: http://www.marinespecies.org

Garrity, G.M., Lilburn, T.G., Cole, J.R., Harrison, S.H., Euzeby, J., and Tindall, B.J. (2007). Taxonomic Outline of the Bacteria and Archaea, Release 7.7 Michigan State University Board o Trustees. DOI: 10.1601/TOBA7.7

Gibson, D., Jones, A. & Bray, R. (Eds.) (2002). Keys to the Trematoda: Volume 1. CAB International, London.

Gibson, D., Jones, A. & Bray, R. (Eds.) (2005). Keys to the Trematoda: Volume 3. CAB International, London.

Gill, F. & Donsker, D. (Eds). (2014). IOC World Bird List (v 4.1). Accessed at: http://www.worldbirdnames.org [Accessed 23 February 2014].

Gofas, S. (2013a). Monoplacophora. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Gofas, S. (2013b). Polyplacophora. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Gofas, S. (2014a). Patellogastropoda. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Gofas, S. (2014b). Vetigastropoda. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Goffinet, B., Buck, W. & Shaw, A.J. (2008). Morphology, anatomy, and classification of the Bryophyta. In Goffinet, B. & Shaw, A.J. Bryophyte Biology, 2nd Edition. Cambridge University Press, Cambridge UK, pp. 55–138.

Guidetti, R. & Bertolani, R. (2011). Phylum Tardigrada Doyère, 1840. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 96–97.

Guiry, Michael D. (2013a). Chlorophyceae. In: Guiry, M.D. & Guiry, G.M. (2013). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic informative republished from AlgaeBase with permission of M.D. Guiry). Accessed through World Register Marine Species at: http://www.marinespecies.org

Guiry, Michael D. (2013b). Charophyta. In: Guiry, M.D. & Guiry, G.M. (2013). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic informative republished from AlgaeBase with permission of M.D. Guiry). Accessed through World Register Marine Species at: http://www.marinespecies.org

Guiry, Michael D. (2013c). Chromista. In: Guiry, M.D. & Guiry, G.M. (2013). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic informative republished from AlgaeBase with permission of M.D. Guiry). Accessed through World Register Marine Species at: http://www.marinespecies.org

Gupta, R. S., Chander, P., & George, S. (2013). Phylogenetic framework and molecular signatur for the class Chloroflexi and its different clades; proposal for division of the class Chloroflexi clanov. into the suborder Chloroflexineae subord. nov., consisting of the emended family Oscillochloridaceae and the family Chloroflexaceae fam. nov., and the suborder Roseiflexineae subord. nov., containing the family Roseiflexaceae fam. nov. Antonie van Leeuwenhoek 103: 99–119.doi:10.1007/s10482-012-9790-3pmid:22903492.

Harvey, M.S. (2011) Smaller Arachnid Orders Catalogue (SAOCat) database. In: Bisby F.A., Roskov Y.R., Orrell T.M., Nicolson D., Paglinawan L.E., Bailly N., Kirk P.M., Bourgoin T., Baillargeon G. & Ouvrard D. (eds.). Species 2000 & ITIS Catalogue of Life: 2011 Annual Checklist. Digital resource at http://www.catalogueoflife.org/annual-checklist/2011/

Hedlund, B.P. (2012). Validation List 143, Family III. Rubritaleaceae fam. nov. Int. J. Syst. Ev Microbiol.: 62, 1-4.

Hedlund, B.P. & Derrien, M. (2012). Akkermansiaceae fam. nov. VALIDATION LIST no. 143 Int. J. Syst. Evol. Microbiol.: 62, 1-4.

Hess, S., Sausen, N., & Melkonian, M. (2012). Shedding Light on Vampires: The phylogeny vampyrellid amoebae revisited. PLoS ONE 7 (2): e31165.

Henry, T. (2009). Biodiversity of the Heteroptera. In: Footit, R. & Adler, P. (Eds.). Insect Biodiversity: Science and Society. Oxford: Wiley-Blackwell, pp. 223-263.

Hodda, M. (2011). Phylum Nematoda Cobb 1932. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: *I* outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 63–95.

Hooper, J.N.A., Van Soest, R.W.M. & Pisera, A. (2011). Phylum Porifera Grant, 1826. In: Zhan Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonon richness. Zootaxa 3148: 13–18.

Howe, A.T., Bass, D., Vickerman, K., Chao, E.E., & Cavalier-Smith, T. (2009) Phylogeny, taxonomy, and astounding genetic diversity of glissomonadida ord. nov., the dominant gliding zooflagellates in soil (Protozoa: Cercozoa). Protist 160(2): 159-189.

Howe, A.T., Bass, D., Scoble, J.M., Lewis, R., Vickerman, K., Arndt, H. & Cavalier-Smith, T. (2011). Novel cultured protists identify deep-branching environmental DNA clades of Cercozoa new genera Tremula, Micrometopion, Minimassisteria, Nudifila, Peregrinia. Protist 62: 332–37.

Im, W-T., Hu, Z-Y., Kim, K-H., Rhee, S-K., Meng, H., Lee, S-T., & Quan, Z-X. (2012). Description of Fimbriimonas ginsengisoli gen. nov., sp. nov. within the Fimbriimonadia class no the phylum Armatimonadetes. Antonie van Leeuwenhoek: 102 (2): 307-317.

Integrated Taxonomic Information System. (2013). Accessed at: http://www.itis.gov

Integrated Taxonomic Information System. (2014). Accessed at: http://www.itis.gov

Ishitani, Y., Ujiie', Y., de Vargas, C., Not, F., & Takahashi, K. (2012) Phylogenetic relationship and evolutionary patterns of the Order Collodaria (Radiolaria). PLoS ONE 7(5): e35775.

James, S. & Davidson, S. (2012). Molecular phylogeny of earthworms (Annelida: Crassiclitellal based on 28S, 18S, and 16S

Jamieson, B. (1988). On the phylogeny and higher classification of the Oligochaeta. Cladistics 367-401.

Janssens, F. & Christiansen, K.A. (2011). Class Collembola Lubbock, 1870. In: Zhang, Z.-Q. (E Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 192–194.

Karlsbakk, E. (2013). Bivalvulida and Multivalvulida. Accessed through World Register of Marl Species at: http://www.marinespecies.org

Kirk, P. (2013). Updated database version of Kirk, P.M., Cannon, P.F., Minter, D.W. & Stalpers J.A. (2008). Dictionary of Fungi, 10th Edition, CAB International, Oxon, UK.

Konikoff, C. (2013). Graptolithoidea. In: Shenkar, N., Swalla, B.J., & van der Land, J. (2013). Hemichordata World Database. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Kristensen, R. (2002). An introduction to Loricifera, Cycliophora, and Micrognathozoa. Integ. Comp. Biol., 42: 641–651.

Kroh, A. (2014). Echinoidea. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Kury, A. (2013). Order Opiliones Sundevall, 1833. In: Zhang, Z.-Q. (Ed.) Animal Biodiversity Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). Zoc 3703: 27-33.

Lackey, J. (1933). Studies in the life histories of Euglenida III. The morphology of Peranema tichophorum Ehrenberg, with special reference to its kinetic elements and the classification of th Heteronemidæ. Biol. Bull. 65 (2): 238-248.

Lee, J.J., Leedale, G.F. & Bradbury, P. (Eds). (2000). An Illustrated Guide to the Protozoa, 2nd Edition: Organisms Traditionally Referred to as Protozoa, or Newly Discovered Groups. Volume & 2. Society of Protozoologists, Lawrence, KS, 1432 pp.

Lee, K.B., Liu, C.T., Anzai, Y., Kim, H., Aono, T., & Oyaizu, H. (2005). The hierarchical system of the 'Alphaproteobacteria': description of Hyphomonadaceae fam. nov., Xanthobacteraceae far nov. and Erythrobacteraceae fam. nov. Int. J. Syst. Evol. Microbiol., 55, 1907-1919.

Li, W.J., Zhi, X.Y., & Euzéby, J.P. (2008). Proposal of Yaniellaceae fam. nov., Yaniella gen. no and Sinobaca gen. nov. as replacements for the illegitimate prokaryotic names Yaniaceae Li et al. 2005, Yania Li et al. 2004, emend Li et al. 2005, and Sinococcus Li et al. 2006, respectively. Int Syst. Evol. Microbiol. 58: 525-527.

Lynn, D. (2002). The Ciliate Resource Archive. Retrieved from http://www.uoguelph.ca/~ciliates/classification/genera.html

Mah, C. (2014). World Asteroidea database. Retrieved from: World Register of Marine Species http://www.marinespecies.org

Maddison, David R. (1995). Hemiptera. True bugs, cicadas, leafhoppers, aphids, etc.. Version 01 January 1995 (temporary). http://tolweb.org/Hemiptera/8239/1995.01.01 in The Tree of Life We Project, http://tolweb.org/

Marin, B. & Melkonian, M. (2010). Molecular phylogeny and classification of the Mamiellophy class. Nov. (Chlorophyta) based on sequence comparisons of the nuclear- and plastid-encoded rRNA operons. Protist 161: 304-336.

Marshall, B., Bouchet, P. & Gofas, S. (2014). Trochaclididae Thiele, 1928. Retrieved from: W Register of Marine Species at http://www.marinespecies.org

McGavin, G.C. (1993). Bugs of the World. Facts on File, Inc., New York, NY, 192 pp.

Messing, C. (2014). World List of Crinoidea. Retrieved from: World Register of Marine Species http://www.marinespecies.org

Mills, C. (2012). Phylum Ctenophora: List of all valid species names (published by the author, w page established March 1998, last updated 15 May 2012). Available at: http://faculty.washington.edu/cemills/Ctenolist.html.

Minelli, A. (2011). Class Chilopoda, class Symphyla and class Pauropoda. In: Zhang, Z.-Q. (Ed. Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 157–158.

Monk, S. & Richardson, D.J. (2011). Phylum Acanthocephala Kohlreuther, 1771. In: Zhang, Z.-(Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic ric Zootaxa 3148: 234–237.

Mound, L. (2013). Order Thysanoptera Haliday, 1836. In: Zhang, Z.-Q. (Ed.) Animal Biodivers An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). Zootaxa 3703: 49-50.

National Center for Biotechnology Information (2013). NCBI Taxonomy Browser. Accessed at: http://www.ncbi.nlm.nih.gov/

Nicholls, K. (2012). Zoelucasa sablensis n. gen. et n. sp. (Cercozoa, Incertae Sedis), a new scale-covered flagellate from marine sandy shores. Acta Protozool. 51:113-117.

Norenburg, J. & Gibson, R. (2013). World Nemertea Database. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Nunoura, T., Takaki, Y., Kakuta, J., Nishi, S., Sugahara, J., Kazama, H., Chee, G-J., Hattori, M., Kanai, A., Atomi, H., Takai, K., & Takami, H. (2011). Insights into the evolution of Archaea al eukaryotic protein modifier systems revealed by the genome of a novel archaeal group. Nucleic 1 Res. 39: 3204–3223.

Oliveira, I., Read, V., & Mayer, G. (2012). A world checklist of Onychophora (velvet worms), vnotes on nomenclature and status of names. Zookeys 211: 1-70. doi: 10.3897/zookeys.211.3463

Ovrard, D. (2014). Psyl'list - The World Psylloid Database. Accessesd at: http://www.hemiptera-databases.com/psyllist

Pape, T. & Thompson, F. C. (Eds). (2013). Systema Dipterorum, Version 1.5. Accessed at: http://www.diptera.org/

Paulay, G. (2014). Holothuroidea. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Platnick, N. (2013). The World Spider Catalog, Version 14.0. American Museum of Natural History. Accessed at: http://research.amnh.org/iz/spiders/catalog

Prendini, P. (2011a). Order Amblypygi Thorell, 1883. In: Zhang, Z.-Q. (Ed.), Animal biodivers. An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 154.

Prendini, P. (2011b). Order Scorpiones C.L. Koch, 1850. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 115-117.

Prendini, P. (2011c). Order Solifugae Sundevall, 1833. In: Zhang, Z.-Q. (Ed.), Animal biodivers An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 118.

Pyron, R., Burbrink, F., & Wiens, J. (2013). A phylogeny and revised classification of Squamat including 4161 species of lizards and snakes. BMC Evolutionary Biology 13: 93. http://www.biomedcentral.com/1471-2148/13/93

Read, G. & Fauchald, K. (2013). World Polychaeta database. Accessed through World Register Marine Species at: http://www.marinespecies.org

Renzaglia, K., Villareal, J. & Duff, R.J. (2009). New insights into morphology, anatomy, and systematics of hornworts. In Goffinet, B. & Shaw, A.J. (Eds) Bryophyte Biology, 2nd Edition. Cambridge University Press, Cambridge UK, pp.139–172.

Reverter Gil, O. & Hayward, P. (2013). Aethozoontidae. In: Bock, P.; Gordon, D. (2013) World List of Bryozoa. Retrieved from: World Register of Marine Species at http://www.marinespecies

Rogers, C.E., Mattox, K.R., & Stewart, K.D. (1980). The zoospore of Chlorokybus atmophyticus, a charophyte with sarcinoid growth habit. American Journal of Botany 67(5): 774-

Rosling, A., Cox, F., Cruz-Martinez, K., Ihrmark, K., Grelet, G., Lindahl, B., Menkis, A., James (2011). Archaeorhizomycetes: unearthing an ancient class of ubiquitous soil fungi". Science 33: (6044): 876–9.

Rosenberg, G. & Gofas, S. (2014). Sigmurethra. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Ruggiero, M. & Gordon, D. (Eds). (2013). Hierarchical Classification for Managing the Catalog of Life. Species 2000 and ITIS Catalogue of Life Report. 67 pp. (unpublished).

Ruggiero, M. & Gordon, D. (Eds). (2014). Hierarchical Classification for Managing the Catalog of Life. Species 2000 and ITIS Catalogue of Life Report. 57 pp. (in review/press).

Saunders, G.W. & Hommersand, M.H. (2004). Assessing red algal supraordinal diversity and taxonomy in the context of contemporary systematic data. American Journal of Botany 91: 1494-1507.

Schatz, H., Behan-Pelletier, V., O'Connor, B. & Norton, R. (2011). Suborder Oribatida van der Hammen, 1968. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classific and survey of taxonomic richness. Zootaxa 3148: 141-148.

Schuchert, P. (2013). Polypodiidea. Accessed through World Register of Marine Species at: http://www.marinespecies.org

Seenivasan, R., Sausen, N., Medlin, L.K., & Melkonian, M. (2013). Picomonas judraskeda Gen Sp. Nov.: The First Identified Member of the Picozoa Phylum Nov., a Widespread Group of Picoeukaryotes, Formerly Known as 'Picobiliphytes'. PLoS ONE 8(3): e59565. doi:10.1371/journal.pone.0059565

Segers, H. (2011). Phylum Rotifera Cuvier, 1817. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: A outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 231–233.

Shear, W. (2011). Class Diplopoda de Blainville in Gervais, 1844. In: Zhang, Z.-Q. (Ed.), Anima biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 159–164.

Skerman, V.B.D., McGowan, V., & Sneath, P.H.A. (Eds.) (1980). Approved Lists of Bacterial Names. Int. J. Syst. Bacteriol. 30: 225-420.

Smirnov, A.V., Chao, E., Nassonova, E.S., & Cavalier-Smith, T. (2011). A revised classification naked lobose Amoebae (Amoebozoa: Lobosa). Protist: 162 (4): 545-570.

Sørensen, M. (2013). Phylum Kinorhyncha. In: Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Ot of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013). Zootaxa 37(63-66.

Soulier-Perkins, A.. (2014). Cercopoidea Organised On Line. Accessesd at: http://www.hemiptera-databases.org/cool

Steiner G. & Kabat A. R. (2001). Catalogue of supraspecific taxa of Scaphopoda (Mollusca). Zoosystema 23 (3): 433-460.

Stevens, P. F. (2013). Angiosperm Phylogeny Website. Version 12. Accessed at: http://www.mobot.org/MOBOT/research/APweb/

Stöhr, S. (2014). Ophiuroidea. In: Stöhr, S.; O'Hara, T. & Thuy, B. (Eds). (2014). World Ophiuroidea database. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Szeptycki, A. (2007). Catalogue of the World Protura. Wydawnictwa Instytutu Systematyki I Ewolucji Zwierzat, Polskiej Akademii Nauk, Kraków, 210 pp.

Thollesson, M. & Norenburg, J.L. (2003). Ribbon worm relationships: a phylogeny of the phylu Nemertea. Journal of the Royal Society of London, B 270; 407–415.

Thuesen, E. (2009). Phylum Chaetognatha. Retrieved from http://academic.evergreen.edu/t/thuesene/chaetognaths/chaettaxon.htm

Timm, T. (2014). Oligochaeta. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Tree of Life (TOL) Web Project. (1995). Diplura. Version 01 January 1995. Accessed at: http://tolweb.org/Diplura

Tyler, S. (2014). Polycladida. Retrieved from: World Register of Marine Species at http://www.marinespecies.org

Tyler, S. & Schilling, S. (2011). Phylum Xenacoelomorpha Philippe et al., 2011. In: Zhang, Z.-(Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic ric Zootaxa 3148: 24–26.

Tyler, S., Schilling, S., Hooge, M., & Bush, L.F. (Comp.) (2012) Turbellarian taxonomic databate Version 1.7. Accessed at: http://turbellaria.umaine.edu

Vdacný, P., Orsi, W., Bourland, W., Shimano, S., Epstein, S., & Foissner, W. (2011). Morphological and molecular phylogeny of dileptid and tracheliid ciliates: Resolution at the base class Litostomatea (Ciliophora, Rhynchostomatia). European Journal of Protistology 47(4): 295-313.

van der Land, J. (2013). [unassigned] Enteropneusta. Accessed through World Register of Marin Species at: http://www.marinespecies.org

van Nieukerken, E., Kaila, L., Kitching, I., Kristensen, N., Lees, D., Minet, J., Mitter, C., Mutan M., Regier, J., Simonsen, T., Wahlberg, N., Yen, S.-H., Zahiri, R., Adamski, D., Baixeras, J., Bartsch, D., Bengtsson, B., Brown, J., Bucheli, S., Davis, D., De Prins, J., De Prins, W., Epstein, M., Gentili-Poole, P., Gielis, C., Hättenschwiler, P., Hausmann, A., Holloway, J., Kallies, A., Karsholt, O., Kawahara, A., Koster, S., Kozlov, M., Lafontaine, J.D., Lamas, G., Landry, J.-F., Lee, S., Nuss, M., Park, K.-T., Penz, C., Rota, J., Schintlmeister, A., Schmidt, B.C., Sohn, J.-C., Solis, M.A., Tarmann, G., Warren, A., Weller, S., Yakovlev, R., Zolotuhin, V., & Zwick, A. (2011). Order Lepidoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outloof higher-level classification and survey of taxonomic richness. Zootaxa 3148: 212-221.

Walter, D.E. (2001). Endemism and cryptogenesis in 'segmented' mites: A review of Australian Alicorhagiidae, Terpnacaridae, Oehserchestidae and Grandjeanidae (Acari: Sarcoptiformes). Australian Journal of Entomology 40(3): 207-211.

Wilson, D.E. & Reeder, D.M. (2011). Class Mammalia Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.), Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa 3148: 56–60.

World Register of Marine Species (WoRMS). (2013a). Echinodermata. Accessed at: http://www.marinespecies.org

World Register of Marine Species (WoRMS). (2013b). Euglenozoa and Percolozoa. In: Guiry, N & Guiry, G.M. (2013). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M.D. (Accessed at: http://www.marinespecies.org

World Register of Marine Species (WoRMS). (2013c). Foraminifera. In: Hayward, B.W., Cedhagen, T., Kaminski, M., Gross, O. (2013). World Modern Foraminifera Database. Accessed at: http://www.marinespecies.org

World Register of Marine Species (WoRMS). (2014). Platyhelminthes. Accessed at: http://www.marinespecies.org

Yabuki, A., Chao, E., Ishida, K-I., Cavalier-Smith, T. (2012). Microheliella maris (Microhelida ord. n.), an ultrastructurally highly distinctive new axopodial protist species and genus, and the uphylum Heliozoa. Protist 163(3): 356–388.

Yabuki, A., Nakayama, T., Yubuki, N., Hashimoto, T., Ishida, K-I., & Inagaki, Y. (2011). Tsukubamonas globosa n. gen., n. sp., A Novel Excavate Flagellate Possibly Holding a Key for Early Evolution in "Discoba". Journal of Eukaryotic Microbiology 58 (4): 319-331.

Yoon, H.S., Mueller, K.M., Sheath, R.G., Ott, F.D. & Bhattacharya, D. (2006). Defining the maj lineages of red algae (Rhodophyta). Journal of Phycology 42, 482–492.

Zhang, Z-Q., Fan, Q-H., Pesic, V., Smit, H., Bochkov, A., Khaustov, A., Baker, A., Wohltmann, A., Wen, T., Amrine, J., Beron, P., Lin, J., Gabrys, G., & Husband, R. (2011). Order Trombidiformes Reuter, 1909. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher classification and survey of taxonomic richness. Zootaxa 3148: 129-138.