

Challenging Problem1

Shivangi Parashar

Lines and Planes

Abstract—This document contains the solution to find the points on the lines that are closest to each other. Given Lines are skew

Download latex-tikz codes from

https://github.com/shivangi-975/Challenge_1/blob/master/Challenge_1.tex

1 Problem

Find the points on the skew lines that are closest to each other in 3-Dimensions? skew line 1 passing through the point $A(1, 1, 0)$ with directional vector $\mathbf{v}_1(2, -1, 1)$ and skew line 2 passing through the point $B(2, 1, -1)$ with directional vector $\mathbf{v}_2(3, -5, 2)$

$$L_1 : x = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + \lambda_1 \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad (1.0.1)$$

and

$$L_2 : x = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} + \lambda_2 \begin{pmatrix} 3 \\ -5 \\ 2 \end{pmatrix} \quad (1.0.2)$$

2 Solution

Let the closest points be $P(p_1, p_2, p_3)$ on skew line1 and $Q(q_1, q_2, q_3)$ on skew line2, Let \mathbf{p}, \mathbf{q} be two points on the lines L_1, L_2

$$\mathbf{p} = x_1 + \lambda_1 \mathbf{v}_1 \quad (2.0.1)$$

$$\mathbf{q} = x_2 + \lambda_2 \mathbf{v}_2 \quad (2.0.2)$$

$$\mathbf{pq} = x_2 + \lambda_2 \mathbf{v}_2 - x_1 - \lambda_1 \mathbf{v}_1$$

$$\mathbf{pq} = \begin{pmatrix} x_2 & \mathbf{v}_2 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_2 \end{pmatrix} - \begin{pmatrix} x_1 & \mathbf{v}_1 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_1 \end{pmatrix} \quad (2.0.3)$$

points P and Q are closest points, p-q will be perpendicular to both the skew lines, Therefore,

$$\mathbf{v}_1^T(\mathbf{q} - \mathbf{p}) = 0 \quad (2.0.4)$$

$$\mathbf{v}_2^T(\mathbf{q} - \mathbf{p}) = 0 \quad (2.0.5)$$

From 2.0.4 and 2.0.5 we have: The dot product of \mathbf{v}_1 with the line \mathbf{pq} is

$$\mathbf{v}_1^T \begin{pmatrix} x_2 & \mathbf{v}_2 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_2 \end{pmatrix} - \mathbf{v}_1^T \begin{pmatrix} x_1 & \mathbf{v}_1 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_1 \end{pmatrix} = 0 \quad (2.0.6)$$

The dot product of \mathbf{v}_2 with the line \mathbf{pq} is

$$\mathbf{v}_2^T \begin{pmatrix} x_2 & \mathbf{v}_2 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_2 \end{pmatrix} - \mathbf{v}_2^T \begin{pmatrix} x_1 & \mathbf{v}_2 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_1 \end{pmatrix} = 0 \quad (2.0.7)$$

solving 2.0.6 and 2.0.7 we get

$$\begin{pmatrix} \mathbf{v}_1^T x_2 & \mathbf{v}_1^T \mathbf{v}_2 & -\mathbf{v}_1^T x_1 & -\mathbf{v}_1^T \mathbf{v}_1 \end{pmatrix} \begin{pmatrix} 1 \\ \lambda_1 \\ 1 \\ \lambda_2 \end{pmatrix} = 0 \quad (2.0.8)$$

simplifying it further

$$\begin{pmatrix} \mathbf{v}_1^T \mathbf{v}_1 & -\mathbf{v}_1^T \mathbf{v}_2 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} = \begin{pmatrix} \mathbf{v}_1^T(x_2 - x_1) \\ \mathbf{v}_2^T(x_2 - x_1) \end{pmatrix} \quad (2.0.9)$$

Substituting values we have $\lambda_1 = 25/59$ and $\lambda_2 = 7/59$ and coordinates of points would be.

$$P = \begin{pmatrix} 109/59 \\ 34/59 \\ 23/59 \end{pmatrix} \quad (2.0.10)$$