

Intermittent Explosive Disorder: A Predictor of Suicidal Ideation and Behavior in a
Combat Population

A Dissertation

Presented to the faculty of the
California School of Professional Psychology
San Francisco Campus
Alliant International University

In Partial Fulfillment
of the Requirements for the Degree
Doctor of Philosophy in Clinical Psychology

By

Erin K Bailey

December 9th 2013

UMI Number: 3620254

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI 3620254

Published by ProQuest LLC (2014). Copyright in the Dissertation held by the Author.

Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code



ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 - 1346

INTERMITTENT EXPLOSIVE DISORDER AND SUICIDALITY

Intermittent Explosive Disorder: A Predictor of Suicidal Ideation and Behavior in a
Combat Population

This dissertation by Erin K Bailey has been approved
by the committee members signed below who
recommend that it be accepted by the faculty of the
California School of Professional Psychology – San Francisco Campus
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

Dissertation Committee:

Quyen Tiet, PhD (Chair)

Mark Balabanis, PhD

Kristin Samuelson, PhD

Date

PREVIEW

Abstract**Intermittent Explosive Disorder: A Predictor of Suicidal Ideation and Behavior in a Combat Population**

Erin K Bailey

California School of Professional Psychology
San Francisco Campus

Alliant International University

High suicide rates among military personnel and veterans reflect a need to better understand risk factors for suicidal behaviors. Using data from the National Comorbidity Survey-Replication, the present study examined the role of recent (12-month) Intermittent Explosive Disorder (IED) diagnosis on lifetime history of suicidal ideation and behaviors (creating a suicide plan, attempting suicide) in 218 individuals who had previously participated in combat. In addition, we examined the relationship between morally injurious combat experiences and subsequent IED diagnosis. Twelve-month IED diagnosis was significantly associated with all three suicide variables and was a significant predictor of suicidal ideation over and above 12-month PTSD and Major Depressive Disorder. The odds of a combat veteran, who met diagnostic criteria for 12-month IED, reporting past suicidal ideation were 6.20 times greater than those without the diagnosis. For suicide plan and suicide attempts, the odds increase to 5.69 and 7.24 times greater for those with 12-month IED, respectively. Twelve-month IED was not significantly associated with any of the examined combat variables or PTSD. Consistent with published research, morally injurious combat variables significantly predicted 12-month PTSD in this sample. The results of the present study are preliminary and represent an initial examination of IED in a combat population.

Dedication

This doctoral dissertation is dedicated in three parts.

To the men and women who have served in the United States military, I thank you from
the bottom of my heart.

To my family – Scott, Leslie, Clint, and Helen – and my amazing friends, you give my
life meaning.

And lastly, to myself. For every challenge that I have overcome.

PREVIEW

Acknowledgements

I would like to thank my dissertation chair and advisor, Quyen Tiet, PhD for his patience, mentorship, and guidance throughout my graduate years. Your encouragement has been greatly appreciated. I would also like to acknowledge my dissertation committee, Kristin Samuelson, PhD and Mark Balabanis, PhD, for their continued support.

PREVIEW

Table of Contents

	<i>Page</i>
Dedication.....	iv
Acknowledgements.....	v
List of Tables.....	viii
I: Statement of the Problem.....	1
Research Aims.....	4
II: Review of the Literature	6
Posttraumatic Stress Disorder	7
Anger.....	10
Impulsivity.....	16
Intermittent Explosive Disorder.....	17
Suicidality.....	19
Summary.....	24
Purposes of the Current Study.....	26
III: Method.....	29
National Comorbidity Survey – Replication.....	29
Participants	29
Measures	30
Demographic Variables.....	30
Diagnostic Variables.....	31
12-month Major Depressive Disorder.....	31
12-month Posttraumatic Stress Disorder.....	32
12-Month Intermittent Explosive Disorder.....	32

INTERMITTENT EXPLOSIVE DISORDER AND SUICIDALITY

Combat Variables.....	33
General Combat Experience.....	33
Accidental Harm.....	34
Purposeful Harm.....	34
Dependent Variable.....	34
Suicide Variables.....	34
Statistical Analyses	34
IV: Results.....	36
Correlation Analyses - Hypothesis One.....	36
Regression and Mediation Analyses – Hypotheses Two and Three.....	38
Suicidal Ideation.....	38
Creating a Plan for Suicide and Attempting Suicide	38
Post Hoc Analyses.....	41
Additional Mediation Analyses.....	41
Twelve-Month PTSD.....	41
V: Discussion.....	43
Combat and Intermittent Explosive Disorder.....	43
IED as a Predictor of Suicidality.....	45
Summary.....	47
Possible Treatment Implications.....	49
Limitations and Future Research.....	51
References.....	54
Appendix: Additional Tables.....	68

List of Tables

	<i>Page</i>
Table 1: <i>Demographics of Total Sample</i>	31
Table 2: <i>Number of Participants Endorsing Specific Combat Experiences for the Entire Sample and the Three Endorsed Diagnostic Variables Separately (MDD, PTSD, and IED)</i>	33
Table 3: <i>Number of Participants Endorsing Specific Suicidal Symptoms for the Entire Sample and the Three Endorsed Diagnostic Variables Separately (MDD, PTSD, and IED)</i>	33
Table 4: <i>Summary of Variable Associations</i>	37
Table 5: <i>Hierarchical Regression Analysis Predicting Suicidal Ideation</i>	39
Table 6: <i>Hierarchical Regression Analysis Predicting a Plan for Suicide</i>	40
Table 7: <i>Hierarchical Regression Analysis Predicting Suicide Attempt</i>	40
Table 8: <i>Hierarchical Regression Analysis Predicting 12-Month PTSD</i>	42
Table 9: <i>Full Hierarchical Regression Analysis Predicting Suicidal Ideation from Diagnostic Variables</i>	69
Table 10: <i>Full (Men-Only) Hierarchical Regression Analysis Predicting a Plan for Suicide from Diagnostic Variables</i>	70
Table 11: <i>Full Hierarchical Regression Analysis Predicting Suicide Attempt from Diagnostic Variables</i>	71
Table 12: <i>Full Hierarchical Regression Analysis Predicting 12-Month PTSD from All Combat Variables</i>	72
Table 13: <i>Full Hierarchical Regression Analysis Predicting 12-Month IED from All Combat Variables</i>	73
Table 14: <i>Men-Only Hierarchical Logistical Regression Analysis Predicting Suicidal Ideation from Diagnostic Variables</i>	74
Table 15: <i>Men-Only Hierarchical Logistical Regression Analysis Predicting Suicide Attempt from Diagnostic Variables</i>	75

INTERMITTENT EXPLOSIVE DISORDER AND SUICIDALITY

Table 16: <i>Men-Only Hierarchical Regression Analysis Predicting 12-Month PTSD from All Combat Variables</i>	76
---	----

Table 17: <i>Men-Only Hierarchical Regression Analysis Predicting 12-Month IED from All Combat Variables</i>	77
--	----

PREVIEW

CHAPTER I

Statement of the Problem

Overall, suicide is the second leading cause of death in military populations (Center for Disease Control and Prevention, 2009). Between 1993 and 1998, suicide was the second leading cause of death among male veterans and the first among female veterans (Wiebe, Conigliaro, Richmond, & Branas, 2006). Completed suicide among military and veteran personnel has risen dramatically in the last few years. In 2008, the US Army experienced an all-time high with regard to completed suicides (i.e. suicides resulting in the death of an individual; Kuehn, 2009). However, the Associated Press cited July 2012 as the worst month for Army suicides since 2009, nearly doubling from the previous month. It is estimated that the rate is approaching one military suicide per day, with deaths surpassing rates of those killed in action in Afghanistan (Associated Press, 2012).

Examining suicidal ideation and behaviors may aid clinicians in identifying those individuals most at risk for completed suicide. For example, in a study of 185 veterans, Kleepies et al. (2001) found that individuals who had planned self-injury exhibited a higher intent to die – an identified risk factor for completed suicide (Harriss, Hawton, & Zahl, 2005). Additionally, research indicates that the best predictor of completed suicide is a past attempt (Sundararaman, Panangala, & Lister, 2008). Given the rise in military suicide over recent years, psychological research should continue to identify patterns of risk factors for suicidality among veteran populations (Genderson, Schonfeld, Kaplan, & Lyons, 2009).

Although research (Kang & Bullman, 2008) indicates that the suicide rate in the overall military population does not differ from the general public, these results may be misleading. Vulnerable subgroups, such as the Army and Marine Corps have higher rates than the national average (Eaton et al., 2006; Kang & Bullman, 2008) and may be at risk for suicide attempts and completed suicide. These branches may also be more likely to be exposed to combat situations, which may fuel suicidal behaviors.

The impact of psychological factors on suicide risk has been thoroughly documented (Capron, et al., 2012; Brown, Beck, Steer, & Grisham, 2000; Kotler, Iancu, Efroni, & Amir, 2001; Maguen et al., 2011; Mann et al., 1999; Nock et al., 2008; Sokero et al., 2005; Vandivort & Locke, 2010). Overall, veterans who have been exposed to combat are at higher risk for mental health problems, such as PTSD, and depression (Bailey, n.d.; Maguen et al., 2011; Morissette, 2011), which are risk factors for both suicidal ideation and suicidal action (Nock et al., 2008; Sareen, et al., 2007). Moreover, the risk of suicidal behaviors increases when risk factors co-occur (Nock et al. 2008) and studies that examine risk factors for suicidal behaviors will help clinicians to better identify vulnerable subgroups and aid practitioners in developing effective preventative interventions.

Previous research has also indicated that the varying types of combat experiences may lead to differential patterns of psychological stress. For example, Maguen et al. (2009) used existing data from the National Vietnam Veterans Readjustment Survey (NVVRS) to examine the relationships between different types of combat experiences and subsequent PTSD in a sample of 1,200 male Vietnam veterans. The researchers found that, even after controlling for other aspects of combat, killing another person was

significantly predictive of PTSD, as well as disassociation, functional impairment, and violent behavior.

In a follow-up study, Maguen et al. (2011) examined the relationship between suicidal ideation and killing in combat (Maguen et al., 2011). Using a sample of 2,854 returning Operation Iraqi Freedom (OIF) soldiers, they examined suicidal ideation, PTSD, depression, as well as whether a soldier had killed another person in combat. They found that while killing in combat was predictive of subsequent suicidal ideation, PTSD and depression mediated the relationship between killing and suicidal ideation in their sample. These research studies provide a deeper understanding as to the differential relationship between types of combat experiences, psychiatric conditions, and suicidal ideation.

As features of PTSD (Joseph, Dagleish, Thrasher, & Yule, 1997; Taft et al., 2007), anger and impulsivity have both been documented in individuals following traumatic experiences (Dyer et al., 2009, Chemtob et al., 1997). However, anger and impulsivity may be individual risk factors for suicidality independent of PTSD (Chemtob et al. 1994; Kotler et al., 2001), as well as predictive of suicidality when combined into one construct (Mann et al., 1999). Intermittent Explosive Disorder (IED), a psychiatric condition characterized by impulsive acts of explosive anger, has been documented in traumatized populations (Fincham et al, 2009; Nickerson, Aderka, Bryant, & Hofmann, 2012; Silove et al., 2009) and has also been implicated in risk for both suicidal ideation (Bromet et al., 2007) and behaviors (Bromet, et al., 2007; McCloskey, et al., 2008). To our knowledge there are no research studies to date that specifically examining how IED relates to suicidality and different types of combat experiences.

Research Aims

The present study seeks to expand on the aforementioned studies by including suicidal behaviors (such as creating a suicide plan and attempting suicide), in addition to suicidal ideation, as an outcome measure in the relationship between combat experiences and psychiatric conditions. Moreover, the above-mentioned studies provide justification for taking into account both depression and PTSD when examining the relationship between combat experiences and suicidal behaviors. It is possible that, in the current study, depression and PTSD will also mediate the relationship between combat experiences and suicidal behaviors such as creating a plan for suicide or attempting suicide. Depression and PTSD should also be accounted for due to overlapping symptomology with Intermittent Explosive Disorder.

Overall, there is a paucity of research clarifying the role of IED as a potential unique predictor of suicidal ideation and behaviors after taking PTSD, depression, and combat experience into account. In addition, existing research has not examined the relationships between these variables in predicting suicidality together in one model. If IED is found to both relate to violent combat experiences and predict suicidal behaviors after accounting for PTSD and depression, current clinical practices for treatment combat veterans who exhibit suicidal behaviors may need to be modified in order to address the unique experience of IED. While the Department of Defense encourages anger management skills training as an intervention for PTSD recovery (Department of Veterans Affairs, 2007), the results of the present study may support additional anger management and impulse-control interventions for suicide prevention among military veterans. The following literature review describes identified predictors of suicidal

ideation and/or behavior in the literature and outlines a rationale for why IED should be examined in conjunction with existing predictors of suicidality.

PREVIEW

CHAPTER II

Review of Literature

The Reinforcement Sensitivity Theory (RST) developed by J.M. Gray (1972; 1988) identifies two main personality-based pathways of motivational activation in response to stress: the behavioral activation system (BAS) and the behavioral inhibition system (BIS), broadly referred to as *approach* and *avoidance*, respectively (Carver, Sutton, & Scheier, 2000; Elliot & Thrash, 2002). Comparable to the flight-or-flight mechanism (Cannon, 1929), these motivational systems are activated based on certain stressful environmental situations.

Neurological research identifies some degree of anterior cortical (AC) asymmetry in human's emotional processing and motivational systems (Davidson, 1992). Decreased activity in the left AC region has been identified in depressed individuals, while right anterior lesions result in mania-like symptoms (Harmon-Jones & Allen, 1998; Robinson & Downhill, 1995). Electroencephalographic (EEG) studies have provided evidence of resting alpha-band asymmetry (8-13 Hz) for different motivational systems (Harmon-Jones & Allen, 1998, Davidson, 1992) – with some degree of laterality suggestive of separate systems of emotional processing of approach (left frontal activation) and avoidance (right frontal activation) affect. As features of affective disorders (e.g. Posttraumatic Stress Disorder, Intermittent Explosive Disorder), emotions such as shame, anger, and impulsivity can be conceptualized in terms of avoidance and approach systems. Posttraumatic stress disorder has been identified as a risk factor for suicidality among veteran populations (Capron, et al., 2012; Hendin & Haas, 1991; Sareen, et al.,

2007). However, given overlapping symptomology with other emotionally deregulating disorders (i.e. PTSD, MDD) the link between suicidality and IED remains less clear.

Posttraumatic Stress Disorder

Avoidance systems can be operationalized as a process in which an individual avoids aversive circumstances (Carver & Harmon-Jones, 2009). This system is based in anxiety and fear of what might happen should the individual initiate goal-oriented behavior. Posttraumatic stress disorder has been described as an imbalance of motivation systems with the downregulation of approach associated with characteristic emotional numbing and upregulation of avoidance associated with both behavioral avoidance and hyperarousal (Stein & Paulus, 2009). Considering the aversive nature of war experiences, PTSD is often prevalent in military and veteran populations following combat.

Research estimates that 13% of military personnel utilizing health care services meet diagnostic criteria for PTSD (Seal, et al., 2007). One study found that PTSD and major depression cost the Department of Defense (DoD) approximately 204.6 million dollars over two years, including the cost of suicide (Tanielian & Jaycox, 2008). Another study found that combat-induced PTSD costs the US Health Care System approximately \$1.5 to \$2.7 billion dollars over two years (Cesur, Sabia, & Tekin, 2011).

Rates of PTSD have been shown to increase following deployment and are linearly related to the length of deployment, with longer deployment lengths correlated with greater risk for PTSD (Adler, Huffman, Bliese, & Castro, 2005). However, the effect of deployment length on mental health outcomes diminishes when considering number of firefights (exchange of gunfire; Cesur et al., 2011). Therefore, the linear effect of deployment length on risk for PTSD may be mediated by the exchange of gunfire and