

Regional Gray Matter Volume Differences Between Adolescents With ADHD and Typically Developing Controls: Further Evidence for Anterior Cingulate Involvement

Journal of Attention Disorders

1–12

© The Author(s) 2016

Reprints and permissions:

sagepub.com/journalsPermissions.nav

DOI: 10.1177/1087054715619682

jad.sagepub.com



Björn Bonath¹, Jana Tegelbeckers¹, Marko Wilke², Hans-Henning Flechtner¹, and Kerstin Krauel¹

Abstract

Objective: The present study investigated structural brain differences between adolescents with ADHD and matched control participants. **Method:** Voxel-based morphometry (VBM) using the DARTEL approach was performed to assess regional gray matter (GM) volumes. Additionally, individual performance on tests of attention was recorded to correlate ADHD related cognitive impairments with regional gray matter abnormalities. **Results:** We found significantly smaller GM volume in subjects with ADHD compared to their matched controls within the anterior cingulate cortex (ACC), the occipital cortex, bilateral hippocampus/amygdala and in widespread cerebellar regions. Further, reductions of the ACC gray matter volume were found to correlate with scores of selective inattention. **Conclusion:** These findings underline that structural alterations in a widespread cortico-subcortical network seem to underlie the observable attention problems in patients with ADHD. (*J. of Att. Dis.* XXXX; XX(X) XX-XX)

Keywords

adolescent ADHD, ACC, voxel-based morphometry

Introduction

ADHD is one of the most prevalent psychiatric disorders of childhood (Costello, Mustillo, Erkanli, Keeler, & Angold, 2003). Its prevalence has been estimated to be around 4% (Costello et al., 2003); it is characterized by developmentally excessive levels of inattention, impulsivity, and hyperactivity (Polanczyk, de Lima, Horta, Biederman, & Rohde, 2007). In the past, functional magnetic resonance imaging (fMRI) studies have been used to gain insights into the pathophysiology of ADHD and to identify brain regions that are associated with symptoms of attentional deficits, insufficient encoding of reward or motivational goals, and inadequate motor control (Bush, Valera, & Seidman, 2005; Christakou et al., 2013; Cubillo et al., 2014; Edel et al., 2013; Morein-Zamir et al., 2014; Mostofsky et al., 2006; Rubia et al., 2009; Rubia et al., 2006; Schulz et al., 2004; Smith, Taylor, Brammer, Toone, & Rubia, 2006). Among other regions, the anterior cingulate cortex (ACC) has been implicated repeatedly in the lack of attentional control ADHD participants display (Bush, 2011; Kessler, Angstadt,

Welsh, & Sripada, 2014; Konrad, Neufang, Hanisch, Fink, & Herpertz-Dahlmann, 2006).

Previous imaging studies on cognitive control provide evidence that the ACC plays a key role in several processes such as reward based (Bush et al., 2002; Gehring & Willoughby, 2002) and simple decision making (Kennerley, Walton, Behrens, Buckley, & Rushworth, 2006). Most notably, it was found to play an important role in error monitoring (Botvinick, Cohen, & Carter, 2004; Kiehl, Liddle, & Hopfinger, 2000; van Veen, Cohen, Botvinick, Stenger, & Carter, 2001) with either up or down regulation of attention in subsequent performance (Weissman, Giesbrecht, Song, Mangun, & Woldorff, 2003; Weissman, Gopalakrishnan,

¹Otto-von-Guericke University Magdeburg, Germany

²University of Tübingen, Germany

Corresponding Author:

Björn Bonath, Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, Otto-von-Guericke University Magdeburg, Leipziger Str. 44, 39120 Magdeburg, Germany.
Email: bjoern.bonath@ovgu.de

Hazlett, & Woldorff, 2005) or strategic adjustments to prevent subsequent conflict processing (Kerns et al., 2004). Particularly, the ACC is strongly associated with selective attention processes, in particular with stimulus selection due to the appearance of additional distracting events (Weissman et al., 2003; Weissman et al., 2005).

ADHD has also been associated with structural differences in several brain regions, including occipital, cerebellar, temporal, frontal, and subcortical regions (Bush, 2011; Frodl & Skokauskas, 2012; Narr et al., 2009; Pironti et al., 2014; Seidman, Valera, & Makris, 2005; Seidman et al., 2006; Shaw, De Rossi et al., 2014; Shaw et al., 2007; Shaw et al., 2011; Shaw et al., 2009; Shaw et al., 2006; Shaw et al., 2012). Again, one of the most consistently reported brain regions to show ADHD-related reductions in gray matter (GM) volume from childhood to adulthood is the ACC (Frodl & Skokauskas, 2012).

In recent years, voxel-based morphometry (VBM; Ashburner & Friston, 2005) has been used to identify localized differences in GM volume between controls and patients with many different disorders. This method relies on algorithms that partition the human brain into GM and white matter and cerebrospinal fluid (CSF), as well as several other (nuisance) tissue classes. The classification can be based on voxel intensity as well as on voxel location (Ashburner & Friston, 2005). The available volumetric neuroimaging studies in patients with ADHD mostly provide region-of-interest analyses based on a priori hypotheses rather than unbiased whole-brain assessments using automated techniques such as VBM. Recently, a structural magnetic resonance imaging (MRI) study (Semrud-Clikeman, Fine, Bledsoe, & Zhu, 2014) examining the ACC observed a correlation between parent-reported inattention in participants with ADHD and regional reductions in volume of the ACC. So far, however, there is only limited evidence of a direct association between ADHD-related symptoms (such as inattention, impulsivity, or hyperactivity) and reduced ACC GM volume in participants with ADHD.

Based on these previous findings, we hypothesized that adolescents with ADHD would have smaller volumetric measures for the ACC. We further hypothesized that measures of inattention would be related to ACC volume.

Method

Participants and Measures

Thirty-six adolescents (all male, $M_{\text{age}} = 13.7$ years, $SD = 1.5$ years, range = 11-17 years) participated in the current study. Participants with an ADHD diagnosis ($n = 18$) were recruited from the Department of Child and Adolescent Psychiatry at the University of Magdeburg, Germany. Typically developing healthy control participants ($n = 18$) were recruited via advertisements in local newspapers. In-person interviews

with all participants using the Revised Schedule for Affective Disorders and Schizophrenia for School-Age Children (K-SADS-PL; Delmo, et al. 2000) were conducted by three board-certified child psychiatrists (K.K., B.B., and J.T.). The Child Behavior Checklist (CBCL) and the Youth Self-Report (Y-SR) were used as additional measures. Diagnostic criteria for ADHD were fulfilled in all patients ($n = 11$ combined type, $n = 6$ inattentive, $n = 1$ with comorbid oppositional defiant disorder [ODD]) according to *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.; *DSM-IV*; American Psychiatric Association, 1994). Participants with ADHD who met criteria for any comorbid psychiatric disorder other than ODD were excluded from the current study. Healthy control participants were included if they did not report previous or current psychiatric disorder. A clinical meeting to review all participants in this study was performed, and all diagnoses were confirmed during this meeting via clinician consensus (B.B., K.K., H.H.F., J.T.). Most ADHD participants had a medication history with current ($n = 10$) or past ($n = 4$) intake of methylphenidate (MPH). All participants completed the Culture Fair Intelligence Test (CFT 20 R; Weiss, 1997) and the d2-test (Brickenkamp et al., 2002) as a standardized instrument to assess selective attention performance. The total score of the d2-test reflects the accuracy of attentional performance by relating the overall number of processed stimuli to the number of errors of commission and omission. participant's hand preference was assessed using the Edinburgh Handedness Inventory (EHI; Oldfield, 1971). Informed assent/consent was obtained from all participants and their parents in accordance with requirements set forth by the local ethics committee at the University of Magdeburg, Faculty of Medicine, which approved the study. Participants were reimbursed for their participation.

MRI Data Acquisition and Processing

All participants underwent MRI scanning on a whole body Siemens 3 T Trio-scanner (Siemens Magnetom Trio, Erlangen, Germany) using an eight-channel phased-array head coil. A T1-weighted three-dimensional data set was obtained using a magnetization-prepared rapid gradient-echo (3D-MPRAGE) sequence (TR/TE/TI = 2,500 ms/4.77 ms/1,100 ms, flip angle = 7° , field of view = 256 mm, 192 slices, voxel size $1 \times 1 \times 1$ mm³). MRI data were pre-processed and analyzed using SPM8 (<http://www.fil.ion.ucl.ac.uk/>), including the VBM8 toolbox (<http://dbm.neuro.uni-jena.de/vbm>), running on Matlab (The Mathworks, Natick, MA).

Initially, all individual anatomical scans were segmented into tissue classes. Importantly, this was achieved without using information from anatomical priors which may bias the estimation (Wilke, Schmithorst, & Holland, 2003). Using a customized pediatric template (Wilke, Holland, Altay, & Gaser, 2008) created by the TOM8 toolbox (<http://dbm.neuro.uni-jena.de/software/tom/>), individual GM scans

Table 1. Demographic and Clinical Characteristics of Our Sample.

	ADHD (<i>n</i> = 18) <i>M</i> ± <i>SD</i>	Controls (<i>n</i> = 18) <i>M</i> ± <i>SD</i>	<i>p</i>
Age	13.6 (1.7)	14.1 (1.3)	.11
Gray matter volume	784 (49.6)	804 (49.5)	.24
IQ (CFT)	106.8 (11.6)	108.1 (12.9)	.76
Selective attention (d2) ^a	101 (8.9)	112.1 (11.7)	<.005
CBCL			
Attention problems	68.2 (5.7)	53.4 (3.9)	<.001
Delinquent rule-breaking behavior	61.7 (9.9)	51.4 (3.4)	<.01
Aggressive behavior	61.5 (6.2)	53.2 (5.7)	<.001
Internalizing	60.4 (7.1)	50.0 (7.4)	<.001
Externalizing	60.6 (5.6)	49.7 (6.8)	<.001
Y-SR			
Attention problems	60.5 (7.7)	53.2 (3.9)	<.001
Delinquent rule-breaking behavior	53.3 (5.0)	54.3 (4.8)	.51
Aggressive behavior	54.4 (6.8)	53.6 (4.9)	.66
Internalizing	52.1 (11.6)	48.2 (15.2)	.45
Externalizing	51.4 (8.2)	50.8 (7.9)	.84

Note. CFT = Culture Fair Intelligence Test; CBCL = Child Behavior Checklist; Y-SR = Youth Self-Report.

^aStandard values: *M* = 100, *SD* = 15.

were first affine-registered and subsequently deformed using the Diffeomorphic Anatomical Registration Through Exponentiated Lie algebra approach (DARTEL; Ashburner, 2007). GM images were then smoothed (FWHM = 8 mm) and Jacobian scaled, reestablishing the original tissue volume before spatial transformation. Thus, each voxel of the resulting images reflects absolute regional GM volume.

Statistical Analysis

To identify GM volume difference between ADHD and healthy control participants, we conducted a two-sample *t* test with age as a covariate. To improve reliability of the results, statistical maps were thresholded at $p < .001$ (uncorrected on the voxel level) with more than 100 contiguous voxels, corresponding to a $p < .05$, cluster-corrected using the false discovery rate (FDR). Using the MarsBar toolbox (Brett, et al. 2002), we extracted the volume estimates from the resulting ACC cluster and imported these values into SPSS for a correlation analysis with the standardized individual score of selective attention (d2-test) across all participants. Demographic variables were assessed using Student's *t* test for continuous variables and Fisher's exact test for categorical variables. Significance was assumed at $p < .05$.

Results

Demographic Characteristics

As can be seen from Table 1, adolescents with ADHD did not show significant differences relative to control participants in age and IQ. Except for three participants (two ADHD, one control), all participants were right handed

(Fisher's exact test, $p > .05$). Parents of the ADHD group reported more internalizing, $t(34) = 4.3$, $p < .001$, and externalizing, $t(34) = 5.3$, $p < .001$, symptoms as well as more attention problems, $t(34) = 9.0$, $p < .001$, for their children than parents of healthy control participants, whereas ADHD participants (relative to control participants) reported more attention problems, $t(34) = 3.6$, $p < .001$, but did not differ in number of reported internalizing, $t(34) = 0.8$, $p = .4$, and externalizing, $t(34) = 0.2$, $p = .84$, symptoms. Furthermore, Table 1 shows that selective attention scores of the d2-test revealed significantly higher values for healthy control participants than for ADHD patients, $t(34) = 3.0$, $p < .005$.

VBM Results of GM

Figure 1 indicates significant differences in regional GM measures in the cerebellum, the occipital region, the ACC, the hippocampus, and the amygdala. Table 2 lists all clusters showing lower GM volume in participants with ADHD; these were observed within the cerebellum not only with a strong tendency toward the right dorsal/posterior lobe but also with reduced GM measures within the anterior/inferior lobe of the right and left hemisphere as well as the left anterior/posterior lobe (Figure 1). No brain region showed increased GM measures in ADHD participants relative to the control group.

Correlation Analysis for ACC GM Measures and Selective Attention

Correlational analysis between measures of ACC GM volume and standardized measures of the d2-test (Figure 1)




Figure 1. Voxel-based morphometry (VBM); regions with reduced gray matter measures in ADHD.

Note. $p = .05$ corrected for multiple comparisons on cluster level with more than 100 voxels per cluster. A positive correlation was found between measures of gray matter volume of the ACC and selective attention performance using the d2-test (correlation coefficient, $r = .39$, $p < .02$). ACC = anterior cingulate cortex.

revealed a significant positive correlation between ACC GM volume and the test scores of selective attention (Pearson's $r = .39$, $p = .02$; two-tailed) across all participants. However, separate analyses of both groups showed that this positive correlation was mainly driven by the ADHD participants (Pearson's $r = .38$; $p = .058$; one-tailed) and was not present in the healthy controls ($r = -.04$; $p = .44$; one-tailed). These results suggest that low performance in selective attention tends to be associated with lower GM volume within the ACC in participants with ADHD.

Discussion

The present study investigated structural brain differences between adolescents with ADHD and matched control

participants by using an optimized VBM approach. We found significantly lower GM volume in participants with ADHD (relative to their matched controls) in several brain regions, such as the ACC, the occipital cortex, and the hippocampus/amygdala complex, as well as in widespread cerebellar regions. Most interestingly, reductions in GM measures within the ACC seem to be related to attentional deficits in participants with ADHD.

ACC GM-Volume Reduction in ADHD

Previous studies consistently reported that ADHD-related structural abnormalities occur before the age of 10 (Shaw et al., 2007) and can persist into adulthood (Frodl & Skokauskas, 2012; Seidman et al., 2011). This has been

Table 2. Regions With Reduced Gray Matter Measures in ADHD Using Age as Covariate.

Brain regions included within significant cluster	Peak coordinate			Z score	Cluster size
	x	y	z		
Right cerebellum lobules Crus I, Crus II, VI	41	-72	-32	5.1	1,624
Right cerebellum lobules VIII, IX	23	-49	-56	5.3	288
Vermis	-2	-55	-41	5.1	117
Left cerebellum lobule Crus II	-42	-62	-45	4.6	124
Left cerebellum lobules VIII, IX	-36	-54	-53	4.6	294
Left cerebellum lobule VI	-35	-55	-30	4.2	126
Left cerebellum lobule Crus II	-24	-79	-42	4.1	116
Left cerebellum lobule Crus I	-29	-78	-30	4.0	110
Occipital cortex	3	-75	9	4.85	464
Right hippocampus-amygdala complex	33	-21	-15	4.4	279
Left hippocampus-amygdala complex	-27	-13	-23	4.3	261
Left anterior cingulate cortex	-5	38	11	3.4	241

Note. FDR-corrected with 100 as the cluster-level threshold at $p < .05$. FDR = false discovery rate.

suggested to be a consequence of a delay of, rather than a deviation from, normal brain maturation (Shaw et al., 2007). For example, consistent with results in our study, it has been shown that children with ADHD had significant cortical thinning within the ACC (Bledsoe, Semrud-Clikeman, & Pliszka, 2013). The authors reported that the cortical thickness of the right ACC predicts a significant amount of the variance in parent-reported symptoms of ADHD, but failed to show a relationship between this region and scores of a word-color inhibition subtest.

Also, it has been shown that a higher rate of cortical thinning in the cingulate gyrus during cerebral cortical development from childhood into adulthood is associated with higher levels of inattention in adults with ADHD (Shaw et al., 2012).

Recently it has been shown that parent ratings of attention significantly predicted the right volume of the ACC (Semrud-Clikeman et al., 2014). Importantly, our current results expand these previous observations, demonstrating a positive correlation between the GM volume of the ACC and test scores of selective attention. While observed over all participants, the association between reductions in ACC GM volume and scores of selective inattention was mainly driven by the ADHD participants. This is the first study to report a correlation between volumetric aberrations in the ACC and scores of selective attention. Interestingly, GM-volume reduction in the ACC has also been reported in adults with ADHD (Amico, Stauber, Koutsouleris, & Frodl, 2011; Matsuo et al., 2009; Seidman et al., 2011; Seidman et al., 2006) when compared with healthy control participants. Furthermore, Matsuo and colleagues (2009) reported that adult ADHD participants had a tendency of inverse correlation between ADHD-related reductions within the left ACC GM volume and the total score of the Barratt Impulsiveness Scale (BIS), so that higher impulsivity tended to be associated

with smaller ACC GM volumes. This result is also well in line with the findings reported here.

Functional MRI studies have also repeatedly implicated the dorsal as well as the rostral ACC in patients with ADHD in a variety of cognitive tasks, irrespective of age. For example, a functional MRI study (Bush et al., 1999) revealed reduced BOLD responses in the ACC in adult participants with ADHD (relative to healthy controls) during an interference Stroop task, while Tamm, Menon, Ringel, and Reiss (2004) reported a decreased BOLD response of the anterior/mid-cingulate cortex during response inhibition in adolescents with ADHD. Furthermore, Konrad and colleagues (2006) observed less activation in the ACC during attentional alerting in children with ADHD. Importantly, Weissman et al. (2005) reported that the ACC plays a key role in directing attention toward task-relevant stimuli, especially when the task is complicated by distracting stimuli. These findings are therefore well in agreement with ours.

ADHD-related medication treatment might also influence the structural development of children with ADHD. Results from a volumetric MRI study (Pliszka et al., 2006) examining healthy control children, ADHD-treatment-naïve children, and chronically treated children with ADHD reported significantly smaller ACC volume for the ADHD-treatment-naïve group compared with the ADHD-treated and control group.

However, a study by Castellanos and colleagues (2002) reported no differences in GM or total brain volume between participants with and without medication history. Similarly, Makris and colleagues (2010) reported that ADHD-related volumetric deficits in the ACC persist into adulthood and are independent of medical treatment.

In line with this study, Castellanos and colleagues (2008) proposed that ADHD-related lapses of attention might be

associated with a decrease in the functional connectivity between the default-mode network-related structures, such as the ACC and parietal regions. Together, these results suggest a strong association of the ACC with ADHD-related dysfunctions in selective attention, which thus might be responsible for increased distractibility.

Amygdala/Hippocampus GM-Volume Reduction in ADHD

It was suggested that children and adolescents with ADHD suffer from a delay in cortical maturation and gyrification bilaterally in the medial temporal cortex, among other regions (Shaw et al., 2012). However, ADHD-related research focusing on temporal and subcortical brain regions found ambiguous results for the hippocampus, with higher, lower, or no brain volume differences (Amico et al., 2011; Plessen et al., 2006; Posner et al., 2014).

Our results of reduced GM volume within the hippocampus and the amygdala bilaterally, however, are in line with more recent findings. For example, Frodl and colleagues (2010) reported reduced GM volume within the amygdala in adults with ADHD relative to matched controls. The authors also reported that ADHD participants who have been treated with stimulants showed reduced GM volume within the left hippocampus relative to medication-naïve ADHD participants but failed to show significant differences in hippocampal volumes between patients with ADHD and healthy controls in general. Posner et al. (2014) compared medication-naïve children with ADHD with healthy control participants and also observed reduced volumes of the left hippocampus, as well as a reduced functional connectivity between the left hippocampus and the left orbitofrontal cortex (OFC). Furthermore, Das, Cherbuin, Anstey, Abhayaratna, and Easteal (2014) reported that a smaller volume of the left hippocampus was associated with greater hyperactivity symptoms but not inattention. Finally, Brieber and colleagues (2007) reported that children with ADHD showed a decrease of GM volume within the left hippocampus–amygdala complex. The authors proposed that this hippocampal GM reduction might be associated with the presence of the met allele of the brain-derived neurotrophic factor (BDNF) gene (Bueller et al., 2006), a polymorphism which has also been associated with ADHD-related symptoms (Friedel et al., 2005). Thus, Brieber and colleagues speculated that genetic variations in the BDNF gene might contribute to structural brain abnormalities in the hippocampal formation and might be thus the reason for impaired sleep-dependent memory consolidation in ADHD (Prehn-Kristensen et al., 2011). However, the picture is not coherent, with some studies even finding increased GM volume in mesial temporal lobe structures of children and adolescents with ADHD (Plessen et al., 2006). Also, no differences of amygdalar or hippocampal GM volume have been reported in adults with ADHD (Amico et al., 2011;

Castellanos et al., 1996; Filipek et al., 1997; Perlov et al., 2008). One important explanation for these inhomogeneous results across studies may be the number of participants with a history of psychostimulant medication (Frodl & Skokauskas, 2012; Posner et al., 2014). Several studies have shown that MPH can modulate hippocampal physiology and/or functions, including long-term potentiation, learning, and memory (Bethancourt, Camarena, & Britton, 2009; Dommett, Henderson, Westwell, & Greenfield, 2008). Likewise, Verster and colleagues (2010) found that MPH significantly improves declarative memory functioning of adults with ADHD and speculated that MPH enhances the release of dopamine within the ventrolateral prefrontal cortex and the hippocampal formation that is involved in encoding and retrieval (Medin et al., 2013). In contrast to this, Lagace, Yee, Bolanos, and Eisch (2006) reported that juvenile MPH exposure leads to decreased long-term survival of new neurons in the adult hippocampus of rats. A study directly testing this hypothesis in a longitudinal design, however, is lacking.

Other explanations for divergent findings might be differences in age ranges, comorbid disorders, imaging methodologies as well as simply due to the fact that most studies did not focus on this complex when defining regions of interest (ROIs). Furthermore, Das and colleagues (2014) proposed that the hippocampal volume seems more associated with the hyperactivity score instead of the inattention score and might be therefore present only in the hyperactive- or combined, but not in the inattentive subtype of ADHD.

The amygdala is thought to be important in ADHD-related problems with hyperactivity (Frodl et al., 2010), impulsivity, and the regulation of emotion (Brotman et al., 2010; Posner et al., 2011). Several studies already reported reduced GM volume of amygdala in participants with ADHD (Frodl et al., 2010; Geurts, Ridderinkhof, & Scholte, 2013; Lim, Radua, & Rubia, 2014; Valera, Faraone, Murray, & Seidman, 2007), even after controlling for the confounding effect of comorbid ODD and conduct disorder (CD; Sasayama et al., 2010). Importantly, Frodl and colleagues (2010) also reported that more hyperactivity and inattention in participants with ADHD were associated with smaller right amygdala volumes. In addition, a resting-state fMRI study reported abnormal interregional connectivity of the frontal–amygdala–occipital network in young adults with ADHD (Cocchi et al., 2012).

To conclude, ADHD-related volumetric changes within the amygdala and the hippocampus have not been consistently reported in adolescents with ADHD and merit further investigation with optimized brain morphological approaches.

Cerebellar GM-Volume Reduction in ADHD

It has been shown repeatedly that the cerebellar volume is smaller in children and adolescents with ADHD compared

with healthy controls (Bledsoe, Semrud-Clikeman, & Pliszka, 2011; Carmona et al., 2005; Castellanos et al., 1996; Castellanos et al., 2002; Durston et al., 2004; Ivanov, Murrough, Bansal, Hao, & Peterson, 2014; Mackie et al., 2007; Montes et al., 2011; Yang et al., 2008). This ADHD-related volumetric abnormality of the cerebellum can even persist into adulthood (Castellanos et al., 2002; Makris et al., 2015). New findings have also found cerebellar vermis volume to be predictive of behavioral parent ratings of hyperactivity, attention, and restlessness/impulsivity (Bledsoe et al., 2011). Moreover, Mackie and colleagues (2007) examined children with ADHD who were divided into an ADHD group with better clinical outcomes and a group with worse outcomes, compared with matched healthy controls. The authors reported that ADHD participants with a worse clinical outcome show progressive neuroanatomical deficits in GM volumes of the right and left posterior inferior-cerebellar lobes during adolescence relative to both groups. Also, in a sibling study (Durston et al., 2004), the authors reported volumetric reductions in cortical GM and white matter in boys with ADHD as well as in their unaffected siblings relative to healthy controls. This suggests an increased ADHD-related familial risk in general, although GM volume of the cerebellum seems unaffected in siblings without ADHD. Traditionally, cerebellum was associated with motor control, such as real-time fine tuning of movements (Allen & Tsukahara, 1974), but it has increasingly been associated with higher cognitive functions and emotion (Leiner, Leiner, & Dow, 1986; Stoodley & Schmahmann, 2010). It has further been shown that the cerebellum is also involved in multiple cerebro-cerebellar loops via thalamic regions. For example, Middleton and Strick (1994) observed that the cerebellum influences several areas of the prefrontal cortex via thalamic connections (Durston, van Belle, & de Zeeuw, 2011). Therefore, ADHD-related dysfunction might also be associated with an impaired interregional connectivity due to the loss of cerebellar GM and white matter volume (Castellanos & Proal, 2012; Makris et al., 2015; Rubia, Alegria, & Brinson, 2014).

Interestingly, similar to a current study on medication-naïve ADHD adults (Makris et al., 2015), our findings show consistent results of ADHD-related volume reduction within cerebellar regions (Crus I and Lobule VI) that are associated with executive functions (Stoodley & Schmahmann, 2009). It is known that the cerebellum is not only interconnected to motor areas but also to non-motor areas of the cerebral cortex, including frontal, parietal, temporal, and occipital cerebral regions (Strick, Dum, & Fiez, 2009), a cortical network involved in executive functions such as attention and impulsive control, core symptoms of ADHD. These results support the idea that not only cerebral- (Frodl & Skokauskas, 2012; Seidman et al., 2011; Shaw et al., 2007) but also cerebellar cortical developmental dysfunction from childhood into adulthood is associated with ADHD-relevant symptoms.

Occipital Cortex GM-Volume Reduction in ADHD

In addition, in the present study we also found reduced GM volume within occipital cortex for adolescents with ADHD relative to healthy controls. These results are in line with previous studies on ADHD-related volumetric abnormality (Ahrendts et al., 2011; Castellanos et al., 2002; Durston et al., 2004; Makris et al., 2012; Proal et al., 2011; Sasayama et al., 2010; Seidman et al., 2011). The occipital cortex is involved in the well-known attentional network to maintain, suppress, and shift attention (Corbetta & Shulman, 2002; Kanwisher & Wojciulik, 2000), and is thus relevant for ADHD-related aspects. Shaw and colleagues (2007) observed that cortical development in children with ADHD lagged behind that of typically developing children by several years, particularly in prefrontal, middle, and superior temporal, and also in occipital regions. In addition, the occipital cortex of ADHD participants has also been associated with abnormalities of interregional connectivity in distinct networks, such as fronto-occipital and temporo-occipital circuits (Cocchi et al., 2012; Lei et al., 2014; Shaw, Sudre et al., 2014). Together, these findings suggest that occipital brain regions and their relevance for the regulation of attentional processes warrant further studies in participants with ADHD.

Limitations

Our study results differed from results of earlier structural brain studies on participants with ADHD (meta-analysis; Frodl & Skokauskas, 2012) in that we did not observe GM reductions in subcortical regions, such as the striatum, the putamen, and the pallidum. Frodl and Skokauskas (2012) suggested that changes in the basal ganglia seem to diminish over time from child to adulthood; however, ADHD-related adulthood seems to be characterized by decreased GM volume in the ACC. Our sample, with a mean age of 13.7 years, may thus already show features more of the adult, rather than the pediatric, structural phenotype. Furthermore, it is also possible that findings of reduced basal ganglia volume in children and adolescents of earlier studies that have used manual tracing segmentation techniques or ROI-based approaches are biased, and thus overestimated. A further limitation of this study is the exclusive presence of male participants, due to the absence of female patients to which our control group was matched. In general, clinical studies have shown that ADHD predominantly affects males and exhibits a male-to-female sex ratio of 4:1 (Faraone et al., 2015; Polanczyk et al., 2007). Our results may therefore not be generalizable to a female ADHD sample. Also in the present study we did not consider participants' medication history as a study by Castellanos and colleagues (2002) reported no differences in GM or total brain volume between participants with and without medication history.

The small sample size used in this study is another limitation. Future studies with larger sample sizes are needed to confirm relationship between ADHD-related structural brain changes and ADHD-related behavioral outcomes.

Conclusion

Our results provide support for several brain regions being abnormally smaller in participants with ADHD. In particular, we suggest that reduced GM volume in the ACC is directly associated with ADHD-related attentional impairments.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by grants from the SFB 779 TP A03 (K.K.; B.B.).

References

- Ahrendts, J., Rusch, N., Wilke, M., Philipsen, A., Eickhoff, S. B., Glauche, V., . . . van Elst, L. T. (2011). Visual cortex abnormalities in adults with ADHD: A structural MRI study. *World Journal of Biological Psychiatry, 12*, 260-270.
- Allen, G. I., & Tsukahara, N. (1974). Cerebrocerebellar communication systems. *Physiological Reviews, 54*, 957-1006.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Amico, F., Stauber, J., Koutsouleris, N., & Frodl, T. (2011). Anterior cingulate cortex gray matter abnormalities in adults with attention deficit hyperactivity disorder: A voxel-based morphometry study. *Psychiatry Research, 191*, 31-35.
- Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage, 38*, 95-113.
- Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage, 26*, 839-851.
- Bethancourt, J. A., Camarena, Z. Z., & Britton, G. B. (2009). Exposure to oral methylphenidate from adolescence through young adulthood produces transient effects on hippocampal-sensitive memory in rats. *Behavioural Brain Research, 202*, 50-57.
- Bledsoe, J. C., Semrud-Clikeman, M., & Pliszka, S. R. (2011). Neuroanatomical and neuropsychological correlates of the cerebellum in children with attention-deficit/hyperactivity disorder—Combined type. *Journal of the American Academy of Child & Adolescent Psychiatry, 50*, 593-601.
- Bledsoe, J. C., Semrud-Clikeman, M., & Pliszka, S. R. (2013). Anterior cingulate cortex and symptom severity in attention-deficit/hyperactivity disorder. *Journal of Abnormal Psychology, 122*, 558-565.
- Botvinick, M. M., Cohen, J. D., & Carter, C. S. (2004). Conflict monitoring and anterior cingulate cortex: An update. *Trends in Cognitive Sciences, 8*, 539-546.
- Brett, M., Anton, J.-L., Valabregue, R., & Poline, J.-B. (2002). Region of interest analysis using an SPM toolbox. *NeuroImage, 16*.
- Brickenkamp, R., Schmidt-Atzert, L., & Liepmann, D. (2002). *Test d2 concentration test*. Göttingen, Germany: Hogrefe.
- Brieber, S., Neufang, S., Bruning, N., Kamp-Becker, I., Remschmidt, H., Herpertz-Dahlmann, B., . . . Konrad, K. (2007). Structural brain abnormalities in adolescents with autism spectrum disorder and patients with attention deficit/hyperactivity disorder. *Journal of Child Psychology and Psychiatry, 48*, 1251-1258.
- Brotman, M. A., Rich, B. A., Guyer, A. E., Lunsford, J. R., Horsey, S. E., Reising, M. M., . . . Leibenluft, E. (2010). Amygdala activation during emotion processing of neutral faces in children with severe mood dysregulation versus ADHD or bipolar disorder. *American Journal of Psychiatry, 167*, 61-69.
- Bueller, J. A., Aftab, M., Sen, S., Gomez-Hassan, D., Burmeister, M., & Zubieta, J. K. (2006). BDNF Val66Met allele is associated with reduced hippocampal volume in healthy subjects. *Biological Psychiatry, 59*, 812-815.
- Bush, G. (2011). Cingulate, frontal, and parietal cortical dysfunction in attention-deficit/hyperactivity disorder. *Biological Psychiatry, 69*, 1160-1167.
- Bush, G., Frazier, J. A., Rauch, S. L., Seidman, L. J., Whalen, P. J., Jenike, M. A., . . . Biederman, J. (1999). Anterior cingulate cortex dysfunction in attention-deficit/hyperactivity disorder revealed by fMRI and the counting Stroop. *Biological Psychiatry, 45*, 1542-1552.
- Bush, G., Valera, E. M., & Seidman, L. J. (2005). Functional neuroimaging of attention-deficit/hyperactivity disorder: A review and suggested future directions. *Biological Psychiatry, 57*, 1273-1284.
- Bush, G., Vogt, B. A., Holmes, J., Dale, A. M., Greve, D., Jenike, M. A., & Rosen, B. R. (2002). Dorsal anterior cingulate cortex: A role in reward-based decision making. *Proceedings of the National Academy of Sciences, 99*, 523-528.
- Carmona, S., Vilarroya, O., Bielsa, A., Tremols, V., Soliva, J. C., Rovira, M., . . . Bulbena, A. (2005). Global and regional gray matter reductions in ADHD: A voxel-based morphometric study. *Neuroscience Letters, 389*, 88-93.
- Castellanos, F. X., Giedd, J. N., Marsh, W. L., Hamburger, S. D., Vaituzis, A. C., Dickstein, D. P., . . . Rapoport, J. L. (1996). Quantitative brain magnetic resonance imaging in attention-deficit hyperactivity disorder. *Archives of General Psychiatry, 53*, 607-616.
- Castellanos, F. X., Lee, P. P., Sharp, W., Jeffries, N. O., Greenstein, D. K., Clasen, L. S., . . . Rapoport, J. L. (2002). Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit/hyperactivity disorder. *Journal of the American Medical Association, 288*, 1740-1748.
- Castellanos, F. X., Margulies, D. S., Kelly, C., Uddin, L. Q., Ghaffari, M., Kirsch, A., . . . Milham, M. P. (2008). Cingulate-precuneus interactions: A new locus of dysfunction in adult attention-deficit/hyperactivity disorder. *Biological Psychiatry, 63*, 332-337.
- Castellanos, F. X., & Proal, E. (2012). Large-scale brain systems in ADHD: Beyond the prefrontal-striatal model. *Trends in Cognitive Sciences, 16*, 17-26.

- Christakou, A., Murphy, C. M., Chantiluke, K., Cubillo, A. I., Smith, A. B., Giampietro, V., . . . Rubia, K. (2013). Disorder-specific functional abnormalities during sustained attention in youth with attention deficit hyperactivity disorder (ADHD) and with autism. *Molecular Psychiatry*, 18, 236-244.
- Cocchi, L., Bramati, I. E., Zalesky, A., Furukawa, E., Fontenelle, L. F., Moll, J., . . . Mattos, P. (2012). Altered functional brain connectivity in a non-clinical sample of young adults with attention-deficit/hyperactivity disorder. *Journal of Neuroscience*, 32, 17753-17761.
- Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, 3, 201-215.
- Costello, E. J., Mustillo, S., Erkanli, A., Keeler, G., & Angold, A. (2003). Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of General Psychiatry*, 60, 837-844.
- Cubillo, A., Smith, A. B., Barrett, N., Giampietro, V., Brammer, M. J., Simmons, A., & Rubia, K. (2014). Shared and drug-specific effects of atomoxetine and methylphenidate on inhibitory brain dysfunction in medication-naïve ADHD boys. *Cerebral Cortex*, 24, 174-185.
- Das, D., Cherbuin, N., Anstey, K. J., Abhayaratna, W., & Eastale, S. (2014). Regional brain volumes and ADHD symptoms in middle-aged adults: The PATH through life study. *Journal of Attention Disorders*. Advance online publication. doi:10.1177/1087054714523316
- Delmo, C. W. O., Gabriel, M., Bölte, S., Marchio, E., & Poustka, F. (2000). *Schedule for affective disorders and schizophrenia for school-age children* (3rd ed.). Frankfurt, Germany: Klinik für Psychiatrie und Psychotherapie des Kindes- und Jugendalters.
- Domett, E. J., Henderson, E. L., Westwell, M. S., & Greenfield, S. A. (2008). Methylphenidate amplifies long-term plasticity in the hippocampus via noradrenergic mechanisms. *Learning & Memory*, 15, 580-586.
- Durston, S., Hulshoff Pol, H. E., Schnack, H. G., Buitelaar, J. K., Steenhuis, M. P., Minderaa, R. B., . . . van Engeland, H. (2004). Magnetic resonance imaging of boys with attention-deficit/hyperactivity disorder and their unaffected siblings. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43, 332-340.
- Durston, S., van Belle, J., & de Zeeuw, P. (2011). Differentiating frontostriatal and fronto-cerebellar circuits in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 69, 1178-1184.
- Edel, M. A., Enzi, B., Witthaus, H., Tegenthoff, M., Peters, S., Juckel, G., & Lissek, S. (2013). Differential reward processing in subtypes of adult attention deficit hyperactivity disorder. *Journal of Psychiatric Research*, 47, 350-356.
- Faraone, S. V., Asherson, P., Banaschewski, T., Biederman, J., Buitelaar, J. K., Ramos Quiroga, J. A., . . . Franke, B. (2015). Attention deficit/hyperactivity disorder. *Nature Reviews*. Advance online publication. doi:10.1038/nrdp.2015.20
- Filipek, P. A., Semrud-Clikeman, M., Steingard, R. J., Renshaw, P. F., Kennedy, D. N., & Biederman, J. (1997). Volumetric MRI analysis comparing subjects having attention-deficit hyperactivity disorder with normal controls. *Neurology*, 48, 589-601.
- Friedel, S., Horro, F. F., Wermter, A. K., Geller, F., Dempfle, A., Reichwald, K., . . . Hebebrand, J. (2005). Mutation screen of the brain derived neurotrophic factor gene (BDNF): Identification of several genetic variants and association studies in patients with obesity, eating disorders, and attention-deficit/hyperactivity disorder. *American Journal of Medical Genetics, Part B: Neuropsychiatric Genetics*, 132B, 96-99.
- Frodl, T., & Skokauskas, N. (2012). Meta-analysis of structural MRI studies in children and adults with attention deficit hyperactivity disorder indicates treatment effects. *Acta Psychiatrica Scandinavica*, 125, 114-126.
- Frodl, T., Stauber, J., Schaaff, N., Koutsouleris, N., Scheuerecker, J., Ewers, M., . . . Meisenzahl, E. (2010). Amygdala reduction in patients with ADHD compared with major depression and healthy volunteers. *Acta Psychiatrica Scandinavica*, 121, 111-118.
- Gehring, W. J., & Willoughby, A. R. (2002). The medial frontal cortex and the rapid processing of monetary gains and losses. *Science*, 295, 2279-2282.
- Geurts, H. M., Ridderinkhof, K. R., & Scholte, H. S. (2013). The relationship between grey-matter and ASD and ADHD traits in typical adults. *Journal of Autism and Developmental Disorders*, 43, 1630-1641.
- Ivanov, I., Murrough, J. W., Bansal, R., Hao, X., & Peterson, B. S. (2014). Cerebellar morphology and the effects of stimulant medications in youths with attention deficit-hyperactivity disorder. *Neuropsychopharmacology*, 39, 718-726.
- Kanwisher, N., & Wojciulik, E. (2000). Visual attention: Insights from brain imaging. *Nature Reviews Neuroscience*, 1, 91-100.
- Kennerley, S. W., Walton, M. E., Behrens, T. E., Buckley, M. J., & Rushworth, M. F. (2006). Optimal decision making and the anterior cingulate cortex. *Nature Neuroscience*, 9, 940-947.
- Kerns, J. G., Cohen, J. D., MacDonald, A. W., III, Cho, R. Y., Stenger, V. A., & Carter, C. S. (2004). Anterior cingulate conflict monitoring and adjustments in control. *Science*, 303, 1023-1026.
- Kessler, D., Angstadt, M., Welsh, R. C., & Sripada, C. (2014). Modality-spanning deficits in attention-deficit/hyperactivity disorder in functional networks, gray matter, and white matter. *Journal of Neuroscience*, 34, 16555-16566.
- Kiehl, K. A., Liddle, P. F., & Hopfinger, J. B. (2000). Error processing and the rostral anterior cingulate: An event-related fMRI study. *Psychophysiology*, 37, 216-223.
- Konrad, K., Neufang, S., Hanisch, C., Fink, G. R., & Herpertz-Dahlmann, B. (2006). Dysfunctional attentional networks in children with attention deficit/hyperactivity disorder: Evidence from an event-related functional magnetic resonance imaging study. *Biological Psychiatry*, 59, 643-651.
- Lagace, D. C., Yee, J. K., Bolanos, C. A., & Eisch, A. J. (2006). Juvenile administration of methylphenidate attenuates adult hippocampal neurogenesis. *Biological Psychiatry*, 60, 1121-1130.
- Lei, D., Ma, J., Du, X., Shen, G., Jin, X., & Gong, Q. (2014). Microstructural abnormalities in the combined and inattentive subtypes of attention deficit hyperactivity disorder: A diffusion tensor imaging study. *Scientific Reports*, 4, Article 6875.
- Leiner, H. C., Leiner, A. L., & Dow, R. S. (1986). Does the cerebellum contribute to mental skills? *Behavioral Neuroscience*, 100, 443-454.
- Lim, L., Radua, J., & Rubia, K. (2014). Gray matter abnormalities in childhood maltreatment: A voxel-wise meta-analysis. *American Journal of Psychiatry*, 171, 854-863.

- Mackie, S., Shaw, P., Lenroot, R., Pierson, R., Greenstein, D. K., Nugent, T. F., III, . . . Rapoport, J. L. (2007). Cerebellar development and clinical outcome in attention deficit hyperactivity disorder. *American Journal of Psychiatry*, 164, 647-655.
- Makris, N., Liang, L., Biederman, J., Valera, E. M., Brown, A. B., Petty, C., . . . Seidman, L. J. (2015). Toward Defining the Neural Substrates of ADHD: A Controlled Structural MRI Study in Medication-Naive Adults. *Journal of Attention Disorders*, 19, 944-953.
- Makris, N., Seidman, L. J., Brown, A., Valera, E. M., Kaiser, J. R., Petty, C. R., . . . Biederman, J. (2012). Further understanding of the comorbidity between Attention-Deficit/Hyperactivity Disorder and bipolar disorder in adults: An MRI study of cortical thickness. *Psychiatry Research*, 202, 1-11.
- Makris, N., Seidman, L. J., Valera, E. M., Biederman, J., Monuteaux, M. C., Kennedy, D. N., . . . Faraone, S. V. (2010). Anterior cingulate volumetric alterations in treatment-naive adults with ADHD: A pilot study. *Journal of Attention Disorders*, 13, 407-413.
- Matsuo, K., Nicoletti, M., Nemoto, K., Hatch, J. P., Peluso, M. A., Nery, F. G., & Soares, J. C. (2009). A voxel-based morphometry study of frontal gray matter correlates of impulsivity. *Human Brain Mapping*, 30, 1188-1195.
- Medin, T., Rinholm, J. E., Owe, S. G., Sagvolden, T., Gjedde, A., Storm-Mathisen, J., & Bergersen, L. H. (2013). Low dopamine D5 receptor density in hippocampus in an animal model of attention-deficit/hyperactivity disorder (ADHD). *Neuroscience*, 242, 11-20.
- Middleton, F. A., & Strick, P. L. (1994). Anatomical evidence for cerebellar and basal ganglia involvement in higher cognitive function. *Science*, 266, 458-461.
- Montes, L. G., Ricardo-Garcell, J., De la Torre, L. B., Alcantara, H. P., Garcia, R. B., Acosta, D. A., & Bouzas, A. F. (2011). Cerebellar gray matter density in females with ADHD combined type: A cross-sectional voxel-based morphometry study. *Journal of Attention Disorders*, 15, 368-381.
- Morein-Zamir, S., Dodds, C., van Hartevelt, T. J., Schwarzkopf, W., Sahakian, B., Muller, U., & Robbins, T. (2014). Hypoactivation in right inferior frontal cortex is specifically associated with motor response inhibition in adult ADHD. *Human Brain Mapping*, 35, 5141-5152.
- Mostofsky, S. H., Rimrodt, S. L., Schafer, J. G., Boyce, A., Goldberg, M. C., Pekar, J. J., & Denckla, M. B. (2006). Atypical motor and sensory cortex activation in attention-deficit/hyperactivity disorder: A functional magnetic resonance imaging study of simple sequential finger tapping. *Biological Psychiatry*, 59, 48-56.
- Narr, K. L., Woods, R. P., Lin, J., Kim, J., Phillips, O. R., Del'Homme, M., . . . Levitt, J. G. (2009). Widespread cortical thinning is a robust anatomical marker for attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48, 1014-1022.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, 9, 97-113.
- Perlov, E., Philipsen, A., Tebartz van Elst, L., Ebert, D., Henning, J., Maier, S., . . . Hesslinger, B. (2008). Hippocampus and amygdala morphology in adults with attention-deficit hyperactivity disorder. *Journal of Psychiatry & Neuroscience*, 33, 509-515.
- Pironti, V. A., Lai, M. C., Muller, U., Dodds, C. M., Suckling, J., Bullmore, E. T., & Sahakian, B. J. (2014). Neuroanatomical abnormalities and cognitive impairments are shared by adults with attention-deficit/hyperactivity disorder and their unaffected first-degree relatives. *Biological Psychiatry*, 76, 639-647.
- Plessen, K. J., Bansal, R., Zhu, H., Whiteman, R., Amat, J., Quackenbush, G. A., . . . Peterson, B. S. (2006). Hippocampus and amygdala morphology in attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*, 63, 795-807.
- Pliszka, S. R., Glahn, D. C., Semrud-Clikeman, M., Franklin, C., Perez, R., Xiong, J., & Liotti, M. (2006). Neuroimaging of inhibitory control areas in children with attention deficit hyperactivity disorder who were treatment naive or in long-term treatment. *American Journal of Psychiatry*, 163, 1052-1060.
- Polanczyk, G., de Lima, M. S., Horta, B. L., Biederman, J., & Rohde, L. A. (2007). The worldwide prevalence of ADHD: A systematic review and meta-regression analysis. *American Journal of Psychiatry*, 164, 942-948.
- Posner, J., Nagel, B. J., Maia, T. V., Mechling, A., Oh, M., Wang, Z., & Peterson, B. S. (2011). Abnormal amygdalar activation and connectivity in adolescents with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50, 828-837.e823.
- Posner, J., Siciliano, F., Wang, Z., Liu, J., Sonuga-Barke, E., & Greenhill, L. (2014). A multimodal MRI study of the hippocampus in medication-naive children with ADHD: What connects ADHD and depression? *Psychiatry Research*, 224, 112-118.
- Prehn-Kristensen, A., Goder, R., Fischer, J., Wilhelm, I., Seeck-Hirschner, M., Aldenhoff, J., & Baving, L. (2011). Reduced sleep-associated consolidation of declarative memory in attention-deficit/hyperactivity disorder. *Sleep Medicine*, 12, 672-679.
- Proal, E., Reiss, P. T., Klein, R. G., Mannuzza, S., Gotimer, K., Ramos-Olazagasti, M. A., . . . Castellanos, F. X. (2011). Brain gray matter deficits at 33-year follow-up in adults with attention-deficit/hyperactivity disorder established in childhood. *Archives of General Psychiatry*, 68, 1122-1134.
- Rubia, K., Alegria, A. A., & Brinson, H. (2014). Brain abnormalities in attention-deficit hyperactivity disorder: A review. *Revista de Neurologia*, 58, S3-S18.
- Rubia, K., Halari, R., Cubillo, A., Mohammad, A. M., Brammer, M., & Taylor, E. (2009). Methylphenidate normalises activation and functional connectivity deficits in attention and motivation networks in medication-naive children with ADHD during a rewarded continuous performance task. *Neuropharmacology*, 57, 640-652.
- Rubia, K., Smith, A. B., Woolley, J., Nosarti, C., Heyman, I., Taylor, E., & Brammer, M. (2006). Progressive increase of frontostriatal brain activation from childhood to adulthood during event-related tasks of cognitive control. *Human Brain Mapping*, 27, 973-993.
- Sasayama, D., Hayashida, A., Yamasue, H., Harada, Y., Kaneko, T., Kasai, K., . . . Amano, N. (2010). Neuroanatomical correlates of attention-deficit-hyperactivity disorder accounting for comorbid oppositional defiant disorder and conduct disorder. *Psychiatry and Clinical Neurosciences*, 64, 394-402.

- Schulz, K. P., Fan, J., Tang, C. Y., Newcorn, J. H., Buchsbaum, M. S., Cheung, A. M., & Halperin, J. M. (2004). Response inhibition in adolescents diagnosed with attention deficit hyperactivity disorder during childhood: An event-related fMRI study. *American Journal of Psychiatry*, *161*, 1650-1657.
- Seidman, L. J., Biederman, J., Liang, L., Valera, E. M., Monuteaux, M. C., Brown, A., . . . Makris, N. (2011). Gray matter alterations in adults with attention-deficit/hyperactivity disorder identified by voxel based morphometry. *Biological Psychiatry*, *69*, 857-866.
- Seidman, L. J., Valera, E. M., & Makris, N. (2005). Structural brain imaging of attention-deficit/hyperactivity disorder. *Biological Psychiatry*, *57*, 1263-1272.
- Seidman, L. J., Valera, E. M., Makris, N., Monuteaux, M. C., Boriell, D. L., Kelkar, K., . . . Biederman, J. (2006). Dorsolateral prefrontal and anterior cingulate cortex volumetric abnormalities in adults with attention-deficit/hyperactivity disorder identified by magnetic resonance imaging. *Biological Psychiatry*, *60*, 1071-1080.
- Semrud-Clikeman, M., Fine, J. G., Bledsoe, J., & Zhu, D. C. (2014). Regional volumetric differences based on structural MRI in children with two subtypes of ADHD and controls. *Journal of Attention Disorders*. Advance online publication. doi:10.1177/1087054714559642
- Shaw, P., De Rossi, P., Watson, B., Wharton, A., Greenstein, D., Raznahan, A., . . . Chakravarty, M. M. (2014). Mapping the development of the basal ganglia in children with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, *53*, 780-789. e711.
- Shaw, P., Eckstrand, K., Sharp, W., Blumenthal, J., Lerch, J. P., Greenstein, D., . . . Rapoport, J. L. (2007). Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proceedings of the National Academy of Sciences*, *104*, 19649-19654.
- Shaw, P., Gilliam, M., Liverpool, M., Weddle, C., Malek, M., Sharp, W., . . . Giedd, J. (2011). Cortical development in typically developing children with symptoms of hyperactivity and impulsivity: Support for a dimensional view of attention deficit hyperactivity disorder. *American Journal of Psychiatry*, *168*, 143-151.
- Shaw, P., Lalonde, F., Lepage, C., Rabin, C., Eckstrand, K., Sharp, W., . . . Rapoport, J. (2009). Development of cortical asymmetry in typically developing children and its disruption in attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*, *66*, 888-896.
- Shaw, P., Lerch, J., Greenstein, D., Sharp, W., Clasen, L., Evans, A., . . . Rapoport, J. (2006). Longitudinal mapping of cortical thickness and clinical outcome in children and adolescents with attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*, *63*, 540-549.
- Shaw, P., Malek, M., Watson, B., Sharp, W., Evans, A., & Greenstein, D. (2012). Development of cortical surface area and gyrification in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, *72*, 191-197.
- Shaw, P., Sudre, G., Wharton, A., Weingart, D., Sharp, W., & Sarlls, J. (2014). White matter microstructure and the variable adult outcome of childhood attention deficit hyperactivity disorder. *Neuropsychopharmacology*, *40*(3): 746-754.
- Smith, A. B., Taylor, E., Brammer, M., Toone, B., & Rubia, K. (2006). Task-specific hypoactivation in prefrontal and temporoparietal brain regions during motor inhibition and task switching in medication-naïve children and adolescents with attention deficit hyperactivity disorder. *American Journal of Psychiatry*, *163*, 1044-1051.
- Stoodley, C. J., & Schmahmann, J. D. (2009). Functional topography in the human cerebellum: A meta-analysis of neuroimaging studies. *NeuroImage*, *44*, 489-501.
- Stoodley, C. J., & Schmahmann, J. D. (2010). Evidence for topographic organization in the cerebellum of motor control versus cognitive and affective processing. *Cortex*, *46*, 831-844.
- Strick, P. L., Dum, R. P., & Fiez, J. A. (2009). Cerebellum and nonmotor function. *Annual Review of Neuroscience*, *32*, 413-434.
- Tamm, L., Menon, V., Ringel, J., & Reiss, A. L. (2004). Event-related fMRI evidence of frontotemporal involvement in aberrant response inhibition and task switching in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, *43*, 1430-1440.
- Valera, E. M., Faraone, S. V., Murray, K. E., & Seidman, L. J. (2007). Meta-analysis of structural imaging findings in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, *61*, 1361-1369.
- van Veen, V., Cohen, J. D., Botvinick, M. M., Stenger, V. A., & Carter, C. S. (2001). Anterior cingulate cortex, conflict monitoring, and levels of processing. *NeuroImage*, *14*, 1302-1308.
- Verster, J. C., Bekker, E. M., Kooij, J. J., Buitelaar, J. K., Verbaten, M. N., Volkerts, E. R., & Olivier, B. (2010). Methylphenidate significantly improves declarative memory functioning of adults with ADHD. *Psychopharmacology*, *212*, 277-281.
- Weiss, R. H. (1997). *Intelligence test 2nd scale (CFT 20)*. Göttingen, Germany: Hogrefe.
- Weissman, D. H., Giesbrecht, B., Song, A. W., Mangun, G. R., & Woldorff, M. G. (2003). Conflict monitoring in the human anterior cingulate cortex during selective attention to global and local object features. *NeuroImage*, *19*, 1361-1368.
- Weissman, D. H., Gopalakrishnan, A., Hazlett, C. J., & Woldorff, M. G. (2005). Dorsal anterior cingulate cortex resolves conflict from distracting stimuli by boosting attention toward relevant events. *Cerebral Cortex*, *15*, 229-237.
- Wilke, M., Holland, S. K., Altaye, M., & Gaser, C. (2008). Template-O-Matic: A toolbox for creating customized pediatric templates. *NeuroImage*, *41*, 903-913.
- Wilke, M., Schmithorst, V. J., & Holland, S. K. (2003). Normative pediatric brain data for spatial normalization and segmentation differs from standard adult data. *Magnetic Resonance in Medicine*, *50*, 749-757.
- Yang, P., Wang, P. N., Chuang, K. H., Jong, Y. J., Chao, T. C., & Wu, M. T. (2008). Absence of gender effect on children with attention-deficit/hyperactivity disorder as assessed by optimized voxel-based morphometry. *Psychiatry Research*, *164*, 245-253.

Author Biographies

Björn Bonath, PhD, is a Postdoctoral Fellow at the Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, Otto-von-Guericke University Magdeburg.

Jana Tegelbeckers, PhD, is a Student at the Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, Otto-von-Guericke University Magdeburg.

Marko Wilke, MD, is Leader of the Experimental Pediatric Neuroimaging Group, Pediatric Neurology and Department of Neuroradiology, University Hospital Tuebingen.

Hans-Henning Flechtner, MD, is Director of the Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, Otto-von-Guericke University Magdeburg.

Kerstin Krauel, PhD, is Leader of the ADHD Research Group at the Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, Otto-von-Guericke University Magdeburg.