

Hypothalamic Modulation of Thermoregulation

The hypothalamus also plays a central role in coordinating thermoregulatory responses and behavior patterns related to this [310]. Afferent input for this regulatory function is from temperature-sensitive neurons in the brain and particularly within the hypothalamus as well as from peripheral thermoreceptors in the skin, abdomen, peritoneum, and spinal cord. The information from all these sources is relayed centrally to the coordinating nuclei in the hypothalamus and other locations to result on autonomic nervous system, behavioral, and neuroendocrine response geared to maintain core body temperature within a narrow preset “normal” range. Thyroid hormones (both LT4 and LT3) are important effector mediators of this normal regulative effect, and their absence in the clinical syndrome of myxedema coma is typically among other features characterized by dramatic hypothermia.

While thermosensitive neurons are virtually ubiquitous in the hypothalamus, the most important sensory locations are the medial and lateral portions of the PON, portions of the anterior hypothalamus including the perifornical region, and the dorsomedial nucleus (DMN) [311–313]. Reduced temperature sensing in the PON results in increased heat generation by induction of the shivering reflex and initiation of non-shivering thermogenesis induced by sympathetic nervous neuron activation of UCP-1 brown adipose tissue (BAT) and by increased metabolic activity in skeletal muscle and other organs. Furthermore, cooling sensing in the PON and other hypothalamic thermo-sensors also induce various heat retention responses including cutaneous vasoconstriction and redirection of blood flow from peripheral to central core depots as well as behavioral responses geared to increased heat generation including moving to warmer places, increased layered clothing, and increased caloric intake. In some animals (though not clearly in humans), there is also an activation of the HPT axis with increased thyroid hormone-induced thermogenesis [314–316]. Contrawise warming

in the PON induces opposite behavioral and autonomic responses including hyperventilation and panting in some animal species, increased sweating, cutaneous vasodilatation, UCP-1 inhibition in BAT, and behavioral response to encourage increased heat loss [311]. The PON has both warm and cold thermosensitive neurons, but the warm thermosensitive neurons are dominant and more numerous in resting environmental states. Typically PON lesions result in anomalies of heat dissipation resulting in hyperthermic clinical states [312]. While the full details of the chemical mediators of the hypothalamic cold and warm neurosecretory responses are still not fully elucidated, Table 5.3 details some of the most prominent mediators thus far identified.

Body temperature in humans in steady state is associated with an intrinsic diurnal rhythm with peak temperature in the late afternoon and early

Table 5.3 Chemical mediators of hypothalamic and CNS thermoregulatory response

Chemical mediators of hypothalamic and CNS-related thermoregulatory signaling	
Hypothermic mediators	Hyperthermic mediators
Estrogens	GABA
Angiotensin 2 (AG2)	Opioid peptides ^a
Acetylcholine (ACH)	Progesterone
CCK	CRH
Dopamine	Prostaglandins
α-MSH	TRH (+ T4 and T3)
Noradrenaline (norepinephrine)	5HT
Opioid peptides ^a	UCP-1
Neurotensin	Irisin
Somatostatin	Endotoxin ^b
Substance P	Lipopolysaccharide (LPS) ^b
ADH (vasopressin)	Tumor necrosis factor-α (TNF-α) ^b
	Interleukin-1 (IL-1) ^b
	Interleukin-6 (IL-6) ^b
	Other proinflammatory cytokines ^b

^aThe variable thermogenic effects of opioid peptides are dependent on the specific peptide(s) involved (viz., endorphins, dynorphins, or enkephalins whether met or leu subtypes) and the receptor types they mediate their actions through (μ, μ; delta, δ; or kappa, κ)

^bThese mediators are only relevant to pathologic states of systemic infections or chronic inflammation and not to steady-state physiologic homeostatic regulation

evenings, while the nadir is in the early morning soon after waking up from sleep. This diurnal rhythm is mainly modulated by the SCN. The SCN has extensive projections to the PON, thus enabling thermo-sensing cross talk to regulate the diurnal rhythm [317, 318]. The set point for body temperature regulation is also sensitive to sex steroidal milieu with the core body temperature typically reaching a nadir just prior to the midcycle LH surge and then rising during the luteal phase of the menstrual cycle mediated by progesterone [319–321]. Estrogen appears to be the major mediator of the midcycle temperature dip which it initiates by stimulation of the heat dissipation response neurons in the PON. Progesterone has the opposite effect on these same neuronal groups of the PON [322–324]. Expectedly, the hypoestrogenic milieu of the human postmenopausal state increases central hypothalamic thermic responses resulting in the well-known menopausal vasomotor syndrome of hot flashes which are also typically well ameliorated with exogenous estrogen administration via defined estrogen replacement regimens [325, 326]. Also notable is that the vasomotor hot flashes are most prevalent in the afternoon and early evenings when the superimposed diurnal rhythm is associated with peaks related to neurosecretory stimulation from the SCN's circadian pacemaker [325, 326]. The fact that some postmenopausal women with hot flashes have a positive clinical response to clonidine (a centrally acting α -2 adrenergic agonist) suggests a possible role for catecholamines in mediating the vasomotor responses [327].

The recent recognition that adult humans have significant depots of BAT as demonstrated by FDG PET imaging demonstrated capacity of transformation of some white adipose tissue (WAT) to BAT under certain humoral and environmental stimuli in addition to the recognition of the intermediate fat tissue depot; beige adipose tissue (BeAT) has opened up another exciting area of investigation into the thermoregulatory and energy expenditure modulatory effects of the hypothalamus [328–331]. There is suggestion that the small amount of BAT present in adult humans (a few grams) when stimulated to induce

thermogenesis by UCP1-dependent means is able to increase total energy expenditure by up to 20%. Animal models lacking this capacity have an obese phenotype suggesting that beyond its role in thermogenesis and energy expenditure, BAT may also have a role in long-term weight management as well [328, 332, 333]. BAT activation physiologically occurs in the setting of cold ambient temperatures and involves a sympatho-excitatory pathway that involves gamma-aminobutyric acid (GABA) release in the PON thermo-sensor and the VMN of the hypothalamus [334]. In addition there appears to be a unique hypothalamic-mediated neurosecretory pathway from the VMN for inducing the beige transformation of WAT that appears mediated via the sympathetic nervous system [335]. There is also some evidence suggesting that brain GLP-1 expressed primarily in the VMN plays a central role both in the browning and beige transformation of WAT but also in BAT-induced thermogenesis. This GLP-1 effect appears to be mediated via AMPK [331, 336–339].

The hyperpyrexia response associated with infections and other systemic inflammatory states represents a unique situation where the normally tightly regulated temperature set point is upregulated for survival benefit of the organism so as to optimize recovery mechanisms from the immunomodulatory system while making the organism as inhospitable as possible for infective agents [340, 341]. This unique pathophysiologic response is achieved by upregulation of the body temperature set point in the thermo-sensors of the PON. This is apparently mediated by endotoxin and other proinflammatory cytokines like tumor necrosis alpha (TNF- α) and interleukin-6 (IL6). These mediators act via receptors in the vascular endothelial cells and subendothelial microglia of the OVLT. This consequently results in activation of the hypothalamic cyclooxygenase activity with increased production of prostaglandin E2 (PGE2) [342–344].

Finally it is also worthwhile to be aware of the potential role that irisin (the relatively recently identified adipo-myokine produced by both adipocytes and skeletal muscle) may have in modulation of thermogenesis by its central action on as

yet unclear hypothalamic nuclei targets. Its roles in centrally mediated reduction in caloric intake as well as modulation of the HPG axis at the hypothalamic level are also still in the early stages of full elucidation [345–352].

Hypothalamic Mediation of Circadian Rhythms

Circadian rhythms are genetically encoded cyclic physiologic functions, systems, and/or behavior patterns generated by internal mechanisms within the living entity. In mammals including humans, the hypothalamus is the most important modulator of such systems [353, 354]. The circadian regulatory system consists of an endogenous rhythm generator which functions akin to the cardiac pacemaker. This is sometimes referred to as the endogenous clock (aka *zeitgeber*), a photo-sensitive sensor system to encode light and dark and to thus link the endogenous clock to the time of day. In humans this is mediated mainly via the retinal photoreceptors (predominantly the cones). The afferent neuronal circuit linking the external environment to the endogenous clock is the retino-hypothalamic pathway, while efferent effector pathway from the endogenous clock is myriad and variable resulting in various modified physiologic functions and behavioral responses. The major endogenous clock in mammals (humans included) as earlier indicated is the hypothalamic SCN. The neurons of the SCN have autonomous synchronously repetitive neuronal firing mediated by local GABA release [355, 356]. Bilateral SCN lesions result in disruption of normal circadian rhythmicity characterized by disrupted sleep-wake cycling on abrogation of the typical daily routines associated with feeding, drinking, and the diurnal rhythms of melatonin and several adeno-hypophyseal hormones. This disruption can then be corrected by subsequent SCN transplantation [357–360]. While the full details of the genetic and molecular underpinnings of the SCN's neuronal cells endogenous pacemaker capacity are not completely understood at present, among the identified so-called “Clock” genes that are

involved in this are the Period (PER), Clock, Cryptochrome (CRY), and Bmal genes [353, 354]. The SCN has two major cellular subgroups: a ventrolateral and a dorsomedial group. The ventrolateral SCN neurons receive the majority of the SCN's afferent inputs which are predominantly via the retino-hypothalamic pathway. These neurons' main neurotransmitter products are pituitary adenylyl cyclase-activating peptide (PACAP) and nitric oxide [353, 354, 361]. The ventrolateral SCN neurons also receive afferent inputs from the intergeniculate leaflet of the thalamus and the midbrain raphe. These sets of afferent neurons utilize GABA, NPY, and 5HT as their main neurotransmitters. These inputs are important in modulating SCN pacemaker variability based on day-night oscillations. The dorsomedial SCN neuronal cells receive the bulk of their afferent input from the limbic system as well as from other hypothalamic nuclei [362, 363]. The efferent neurons of the two nuclear subgroups also have unique neurotransmitters. While the dominant ventrolateral nuclei neurons mainly release VIP, gastrin-releasing peptide, and GABA, the dorsomedial nuclei neurons mainly release arginine vasopressin, AG2, somatostatin, prokineticin 2, and GABA [354, 363, 364].

The SCN has three major diencephalic projections: (a) the hypothalamic subparaventricular zone (SPVZ), (b) the PVN, and (c) the medial and lateral tuberal hypothalamus.

From the SPVZ there are projections to the PON, and as expected these mediate circadian regulation of body temperature set points and food-dependent energy intake [354, 363, 364]. Additional efferent projections from the SPVZ extend to the dorsomedial nucleus of the hypothalamus, midline thalamus, midbrain reticular formation, and basal forebrain. These are involved in modulation of food intake, rest vs movement, sleep-wake cycling, and various pituitary hormone secretions based on light and photic stimuli.

The afferents to the PVN are GABA producing and are mainly involved in melatonin production pineal gland as previously discussed [363, 365, 366]. Beyond melatonin's well-documented role in modulating sleep-wake cycling,

it also has important immunomodulatory effects and is of considerable importance in reproductive functioning behavior among animals with seasonal breeding patterns though this is largely unimportant in humans [367–371]. The PVN projections from the SCN also appear to be central in the circadian night-day alteration of various adenohypophyseal hormone secretory patterns.

The afferents to the medial and lateral tuberal hypothalamus from the SCN are mainly to the VMN, the AN, and the lateral hypothalamic area. These projections are presumed to mainly regulate circadian secretion of neuroendocrine tuberoinfundibular and neurohypophyseal hormonal secretions.

Figure 5.17 provides a schematic overview of the main elements of the hypothalamic circadian regulatory network.

Hypothalamic Modulation of the Sleep-Wake Cycle

The sleep-wake cycle is one of the most important circadian rhythms and is closely regulated by various hypothalamic nuclei. More in-depth description and detailing of the role of the hypothalamus in this will be covered in a separate chapter of this volume. The following provides a basic overview and outline of the elements involved in the hypothalamus' regulatory input into sleep and wakefulness.

While wakefulness is characterized by asynchronous rapid low-voltage beta wave EEG-displayed cerebral cortical activity, the sleep state is more synchronized and organized into stages forms 1 to 4 based on depth of subconsciousness and progressive slowing and increase in amplitude of the detected cortical activity. While stage 1 sleep is characterized by alpha waves, stage 2 is characterized by spindle waves and k-complexes, while stage 3 and 4 sleep are characterized by high-voltage, low-frequency delta waves. States of partial arousal not associated with wakefulness can occur in the course of sleep and are characterized by asynchronous low-voltage electrical cortical activity akin to the

beta waves associated with wakefulness. These are theta waves and are also apparent often soon after sleep initiation. Sleep architecture is further subcategorized depending on the absence or presence of rapid eye movement into rapid eye movement (REM) sleep or non-rapid eye movement (N-REM) sleep. While REM sleep is associated with loss of muscle tone except for respiratory and inner ear muscle activity, N-REM sleep is associated with normal muscle tone and an increased depth of subconsciousness during sleep [372, 373].

The sleep-wake cycle's neuronal regulation involves a balanced coordination between diencephalon basal forebrain and the mid-rostral brainstem. These dual systems subserve states of rest (sleep) versus arousal (wakefulness). There are thus “sleep”- and “wakefulness”-promoting centers. The main sleep center is in the PON and includes neurons in the ventrolateral portions of the PON that have GABA and galanin as their main neurosecretory products [374]. A secondary sleep center has been identified in the midline thalamus (aka the visceral or limbic thalamus) mainly in the dorsomedial and reticular thalamic nuclei [375, 376]. Wakefulness centers include the ascending reticular system of Moruzzi and Magoun [377]. The system consists of two major subdivisions: a monoaminergic and a cholinergic producing set of neuronal nuclei. These are located in the mesencephalic (limbic midbrain) and pontine reticular formation. The monoaminergic system of neurons includes the locus coeruleus which produces noradrenaline, the medial and dorsal raphe which produces 5HT, and the parabrachial nucleus that produces glutamate. In addition, neurons from the ventral periaqueductal gray which produce dopamine are included in this group.

The cholinergic group of neurons which are located in the pontine tegmentum include the laterodorsal and pedunculopontine tegmental nuclei.

Other wakefulness centers have been demonstrated in the tuberomammillary nucleus (producing histamine), nuclei of the lateral hypothalamus, and preformical area that produce orexin, hypocretin, and MCH. Other less prominent but never-




Fig. 5.17 Schema of the hypothalamic circadian regulatory system

theless notable “wakefulness centers” include cholinergic- and GABA-producing neurons of the basal forebrain (aka nucleus basalis of Meynert), portions of the PON, medial septal nucleus, and the nucleus of the diagonal band of Broca [373].

The main sleep center in the PON has efferent neuronal connections with the wakefulness centers in the posterior hypothalamus via the medial forebrain bundle and serves to enable intimate interaction and coordination between sleep and wakefulness cycling. In addition, neuronal connections from the secondary sleep center in the dorsomedial thalamus and reticular formation travel via the medial forebrain bundle to the PeVN and PON as well as to the mesopontine tegmental nuclei.

The wakefulness centers also have efferent outputs of note. These again utilize the medial forebrain bundle, and there are two main such projections: a dorsal route that is mainly cholinergic and GABA secretory as well as a ventral route. The dorsal route projects mainly to the intralaminar and reticular thalamic nuclei and from there diffusely to the cerebral cortex. The ventral route is mainly aminergic and glutamine secreting arising mainly from pontine and mesencephalic nuclei and projects diffusely to the cerebral cortex.

Other Behavioral Modulatory Responses of the Hypothalamus

The hypothalamus plays roles in various other behavioral responses and patterns. This has been aptly demonstrated by various lesional and stimulatory experiments of various hypothalamic nuclei mostly in animal models. More in-depth discussion of these effects and the relevant clinical significance to human behavior and psychopathology will be covered in other chapters of this book. Suffice it to indicate that portions of the medial hypothalamus appear integral to controlling motivated behavior patterns including defensive behavior. Fos labeling has helped identify a group of hypothalamic nuclei in this regard referred to as the “behavior control column” that

are integral to expression of both innate and conditioned defensive behavior patterns [378–380]. Antipredatory defensive behavior has been demonstrated in various animal models with activation of the anterior hypothalamic nucleus, dorsomedial portion of the VMN, and ventrolateral portions of the premamillary nucleus [381–384]. The premamillary nucleus appears to play a critical role in such defensive behavior as destructive lesions of the nucleus abrogate such behaviors [384, 385]. The dorsal periaqueductal gray is an additional important relay station for mediating fear response behaviors [386–388].

The hypothalamus also has a role in the phenotypic behavior response of “social defeat” and “docility.” Animal models demonstrating this behavioral phenotype have been shown to have activated hypothalamic nuclei in sexually dimorphic locations including the PON, the ventrolateral portions of the VMN, and the ventral premamillary nucleus [389, 390]. These nuclei are also involved in modulating and regulating other social behaviors including sexual and aggressive behaviors.

Neuroendocrinology of the Pituitary Axis

Introduction/History

Early recognition for the role of the hypothalamus as a site for integration of endocrine, autonomic, and behavioral responses can be dated to the second to eighteenth centuries A.D [391]. The pituitary gland derives its name from the Greek *ptuo* and Latin *pituita*, meaning phlegm, reflecting its nasopharyngeal origin. Galen in the second century AD hypothesized that nasal phlegm originated from the brain and drained via the pituitary gland [391]. Andreas Vesalius in the sixteenth century included the first anatomical depiction of the infundibular-pituitary stalk. In 1733, Morgagni recorded the absence of adrenal glands in an anencephalic neonate, providing early evidence for a developmental and functional connection between the brain and the adrenal glands [392]. In 1778, Sommering introduces

the term “hypophysis.” In 1849, Claude Bernard set the stage for advances in neuroendocrinology by demonstrating that central lesions to the area of the fourth ventricle resulted in polyuria [392]. In 1948, Geoffrey Harris, the father of modern neuroendocrinology, showed that the anterior pituitary is regulated by the hypothalamus via the hypophyseal portal system [392]. It is now clear that with the hypothalamus, the pituitary orchestrates function of endocrine glands including the thyroid, adrenal, and gonads. The pituitary stalk serves as an anatomic and functional connection to the hypothalamus. Preservation of the hypothalamic-pituitary axis is critical for anterior pituitary control of sexual function and fertility, linear and organ growth, lactation, stress responses, energy, appetite, and temperature regulation and secondarily for carbohydrate and mineral metabolism.

In general, hypothalamus can be considered the coordinating center of the endocrine system. It consolidates signals derived from upper cortical inputs, autonomic function, environmental cues such as light and temperature, and peripheral endocrine feedback. In turn, the hypothalamus delivers precise signals to the pituitary gland, which then releases hormones that influence most endocrine systems in the body that form the hypothalamic-pituitary axis.

Blood Supply

The hypothalamic hormones are small peptides that are active only at the relatively high concentrations achieved in the pituitary portal blood system. Their small size and lack of known binding proteins results in rapid degradation and very low concentrations in the peripheral circulation. Therefore, the unique blood supply of the hypothalamic-pituitary axis is essential to its function. It is why the pituitary gland enjoys an abundant blood supply derived from several sources. Both the anterior lobe and posterior lobe of the pituitary have the same venous drainage, the anterior and posterior hypophyseal veins. The venous plexus empties into the petrosal sinuses

and then into the peripheral circulation via the internal jugular veins [393]. Thus, venous blood in the petrosal sinuses has a relatively high concentration of pituitary hormones and is a useful site to catheterize and sample when assessing pituitary function [391]. But they both have individual arterial supplies. The anterior pituitary gland receives arterial supply from the superior hypophyseal artery (a branch of the internal carotid artery). This vessel first forms a capillary network around the hypothalamus; blood from this network is then transported to a secondary capillary plexus surrounding the anterior pituitary known as the hypothalamo-hypophysial portal system [391]. This structure allows the hypothalamus to communicate with the anterior pituitary via the release of neurotransmitters into the bloodstream. The infundibulum and posterior pituitary gland receive a rich blood supply from many arteries; the major vessels are the superior hypophyseal artery, infundibular artery, and inferior hypophyseal artery.

Anatomy

The *hypothalamus* rests in the sella turcica at the base of the brain, below the third ventricle and above the optic chiasm and pituitary gland. This location is the intersection of the cortex, the cerebellum, and the brainstem [391]. A number of nuclear groups and fiber tracts are recognized in the hypothalamus; they are organized into three major regions including the lateral, medial, and periventricular hypothalamus, each having distinct morphological and functional features. The medial and periventricular subdivisions of the hypothalamus contain a high density of neuronal cell bodies organized into nuclear groups that are crucial for the regulation of the anterior and posterior pituitary gland. Some groups serve as relay centers for highly differentiated neural information coming from the neocortex, limbic system, and autonomic sensory centers in the brainstem of specific homeostatic behaviors such as thirst, hunger, thermoregulation, the sleep-wake cycle, and reproductive behavior. The lateral hypothala-

mus occupies the largest portion of the hypothalamus by volume and has relatively fewer neurons compared to the medial hypothalamus. But the lateral hypothalamus does interact with a massive fiber system, medial forebrain bundle (MFB), through this fiber system that information from the medial forebrain (amygdala, hippocampus, septum, olfactory system, neocortex) and the brainstem is carried to the medial and periventricular hypothalamic subdivisions [391].

One of the most important regions in the hypothalamus, essential for regulation of the pituitary gland, is the median eminence. It is a midline structure located in the basal hypothalamus ventral to the third ventricle and adjacent to the AN [391]. All hypophysiotropic hormones converge there before they are conveyed to the pituitary gland. The median eminence is one of seven circumventricular organs situated as midline structures in the walls of the lateral, third or fourth ventricles. These circumventricular organs contain a rich capillary plexus and have a fenestrated endothelium rendering the structures outside of the blood-brain barrier. This morphologic feature allows the secretion of brain-derived products into the peripheral circulation and/or makes circumventricular organs targets for blood-borne information which can then be transmitted to the brain [391].

Pituitary Gland

The adult human pituitary weighs approximately 600 mg (range 400–900 mg) and measures approximately 13 mm in the longest transverse diameter, 6–9 mm in vertical height, and about 9 mm anteroposteriorly. It is thus the size of average pea [394]. The pituitary gland is comprised of the anterior lobe and posterior lobe along with the pituitary stalk, which connects the median eminence to the pituitary gland. The posterior lobe of the pituitary gland is smaller than the anterior lobe and embryologically derives from the neural primordia as an outpouching from the floor of the third ventricle, a direct anatomic extension of the central nervous system. The posterior pituitary is composed primarily of

unmyelinated axons and axon terminals as well as specialized glial cells called pituicytes [391]. The anterior pituitary derives from the oral ectoderm as Rathke's pouch, first seen by the third week of pregnancy. The anterior lobe contains three subdivisions named pars distalis, pars intermedia, and pars tuberalis. The pars distalis makes up the bulk of the anterior pituitary and is responsible for the secretion of anterior pituitary hormones into the peripheral circulation. The pars intermedia lies between the pars distalis and the posterior pituitary, representing what remains of the original Rathke's pouch cleft and is considered vestigial [391]. Other important boundaries to the pituitary gland are the cavernous sinus laterally, which contain the internal carotid artery surrounded with sympathetic fibers, and the cranial nerves III, IV, V, and VI.

The anterior pituitary is composed of nests of cuboidal cells organized near venous sinusoids lined with a fenestrated epithelium where secretory products are collected. Classically, there are five cell types and six secretory products of the anterior pituitary gland: somatotrophs (growth hormone), lactotrophs (prolactin), corticotrophs (adrenocorticotrophic hormone), thyrotrophs (thyroid-stimulating hormone), and gonadotrophs (luteinizing hormone and follicle-stimulating hormone). However the anterior pituitary can also synthesize numerous other nonclassical peptides and substances that are important for paracrine and/or autocrine control of anterior pituitary secretion and/or cell proliferation under defined physiological conditions [391].

Table 5.4 details the other nonclassical peptides and other chemical mediators thus far identified to be produced by the anterior pituitary gland.

Physiology of Anterior Pituitary Hormone

Prolactin (PRL)

Physiology Prolactin (PRL) is synthesized in lactotroph cells which makes up 15–25% of func-

Table 5.4 List of nonclassical peptides and other chemical mediators from the anterior pituitary: adenohypophysis

List of nonclassical peptides and substances produced by anterior pituitary	
Substances	Cell types
<i>Peptides</i>	
Activin B, inhibin, follistatin	F, G
Aldosterone-stimulating factor	UN
Angiotensin II (angiotensinogen, angiotensin I)	—
Converting enzyme, cathepsin B, renin	C, G, L, S
Atrial natriuretic peptide	G
Corticotropin-releasing hormone-binding protein	C
Dynorphin	G
Galanin	L, S, T
Gaw (chromogranin B)	G
Growth hormone-releasing hormone	UN
Histidyl-proline diketopiperazine	UN
Motilin	S
Neuromedin B	T
Neuromedin U	C
Neuropeptide Y	T
Neurotensin	UN
Protein 7B2	G, T
Somatostatin 28	UN
Substance P (substance K)	G, L, T
Thyrotropin-releasing hormone	G, L, S, T
Vasoactive intestinal polypeptide	G, L, T
<i>Growth factors</i>	
Basic fibroblast growth factor	C, F
Chondrocyte growth factor	UN
Epidermal growth factor	G, T
Insulin-like growth factor I	S, F
Nerve growth factor	UN
Pituitary cytotropic factor	UN
Transforming growth factor alpha	L, S, G
Vascular endothelial growth factor	F
<i>Cytokines</i>	
Interleukin-1 beta	T
Interleukin-6	F
Leukemia inhibitory factor	C, F
<i>Neurotransmitters</i>	
Acetylcholine	C, L
Nitric oxide	F

C corticotropin, F folliculostellate cell, G gonadotroph, L lactotroph, S somatotroph, T thyrotroph, UN unknown

tioning anterior pituitary cells. Human PRL gene, located on chromosome 6, arose from a common ancestral gene, which gave rise to PRL, growth

hormone, and placental lactogen-related proteins [395]. PRL-expressing cells appear to arise from GH-producing cells. The identification of PRL was elusive because human GH is highly lactogenic and active in bioassays used to isolate and measure PRL. GH is present in human pituitary glands in much higher concentrations (5–10 mg) than PRL (approximately 100 µg) [396]. Occasional mammosomatotroph cells co-secrete both PRL and GH, often stored within the same granule. Prolactin structure is a 199-amino acid polypeptide containing three intramolecular disulfide bonds. It circulates in the blood in various sizes: monomeric PRL (“little prolactin”; 23 kDa), dimeric PRL (“big prolactin”; 48–56 kDa), and polymeric forms (also known as “big, big prolactin”; >100 kDa) [397]. The monomeric form is the most bioactive PRL. In response to TRH, the proportion of the monomeric form will increase. Monomeric PRL is cleaved into 8- and 16-kDa forms and the 16-kDa variant is anti-angiogenic. A glycosylated form is less biologically active than little PRL [398]. PRL receptor gene is a member of the cytokine receptor superfamily, localized to chromosome 5p13, creating receptor with an extracellular domain, a hydrophobic transmembrane domain, and an intracytoplasmic region homologous to the GH receptor [399]. PRL receptor dimerization occurs in both ligand-dependent and ligand-independent manners, a single PRL molecule binding to both components of the receptor dimer for formation of the trimeric ligand-receptor complex and subsequent activation by phosphorylation of intracellular Janus kinase/signal transducers (JAK-STAT) [400]. PRL receptors are expressed in many tissues from the pituitary, liver, adrenal cortex, kidneys, prostate, ovary, testes, intestine, epidermis, pancreatic islets, lung, myocardium, brain, and lymphocytes and most recognizable in breast tissue.

Regulation Prolactin (PRL) regulation is under the inhibitory control of dopamine, which is produced by the tuberoinfundibular (TIDA) cells and the hypothalamic tuberohypophyseal dopaminergic system [401]. Dopamine reaches the lactotrophs via the hypothalamic-pituitary portal

system and inhibits PRL secretion by binding to the type 2 dopamine receptors on pituitary lactotrophs. Prolactin participates in negative feedback by increasing tyrosine hydroxylase activity and thereby dopamine synthesis in the TIDA neurons [402]. Several other factors influence PRL gene expression, including estrogen, dopamine, thyrotropin-releasing hormone (TRH), and thyroid hormones [392]. Many factors have been shown to inhibit PRL synthesis and secretion which are endothelin-1, transforming growth factor beta 1 (TGF- β 1), fibroblast growth factors (FGF), vasoactive intestinal polypeptide (VIP), growth hormone-releasing hormone (GHRH), histamine, serotonin, calcitonin, and opiates [392].

Secretion Prolactin (PRL) is cleared rapidly, with a calculated disappearance half-life ranging from 26 to 47 min. PRL secretion occurs episodically in 4–14 secretory pulses over 24 h, each lasting 67–76 min, with the highest levels achieved during sleep and the lowest occurring between 10 am and noon [403]. Prolactin levels decline with age in both men and women. In older men, less PRL is produced with each secretory burst. Postmenopausal women have lower mean serum PRL levels and PRL pulse frequency than do premenopausal women, suggesting a stimulatory effect of estrogen on both these parameters [404].

Function PRL is essential for human survival, because it is responsible for milk production during pregnancy and lactation. Additional biologic functions include reproductive and metabolic effects, mammary development, and melanin synthesis [405]. Serum PRL concentrations rise to 10 times normal during pregnancy. Active lactation is due in part to estrogen and progesterone levels and elevation of PRL levels after delivery. Suckling increases milk production after parturition and is essential for continued lactation. In the absence of suckling, PRL concentrations return to normal by 7 days postpartum. Amenorrhea and infertility result from PRL-mediated inhibitory effects on hypothalamic gonadotropin-releasing hormone (GnRH) neu-

rons and on the pituitary to reduce secretion of the gonadotropins, LH, and FSH, resulting in a reduction in both amplitude and frequency of LH pulses [406].

Growth Hormone (GH)

Physiology Growth hormone (GH) is a single-chain polypeptide hormone consisting of 191 amino acids, which is synthesized, stored, and secreted by somatotroph cells. GH is the most abundant anterior pituitary hormone as somatotroph cells constitute up to 50% of the total anterior pituitary cell population and they are located predominantly in the lateral wings of the anterior pituitary gland. The pituitary gland contains a total of 5–15 mg of GH. Human growth hormone (hGH) genome locus spans approximately 66 kb and contains a cluster of five highly conserved genes located on the long arm of human chromosome 17q22–24 [407]. It encodes the many forms of hGH and human chorionic somatomotropin (hCS); one is hGH-N. The hGH-N gene is selectively transcribed in pituitary somatotrophs and codes for a 22-kDa (191-amino acid) protein which is the predominate GH variant, accounting for 75% of pituitary GH secretion [407]. But 10% of pituitary GH is a 20-kDa variant lacking amino acid residues formed from alternate splicing. Circulating GH molecules comprise several heterogeneous forms: 22- and 20-kDa monomers, an acetylated 22-kDa form, and two desamino GH molecules. The 22-kDa peptide is the major physiologic GH component, while the 20-kDa GH has a slower metabolic clearance [408]. GH elicits intracellular signaling through a peripheral growth hormone receptor (GHR), initiating a phosphorylation cascade involving the JAK-STAT pathway. GHR is a 620-amino acid 70-kDa protein of the class I cytokine/hematopoietin receptor superfamily consisting of an extracellular ligand-binding domain, a single membrane-spanning domain, and a cytoplasmic signaling component [409]. JAK2 (tyrosine kinase) activation leads to phosphorylation of intracellular signaling molecules, signal-transducing activators of transcription proteins

(STATs 1, 3, and 5), and critical signaling components for GH action [410].

Regulation Growth hormone synthesis and release are under control of a variety of hormonal agents, including growth hormone-releasing hormone (GHRH), somatostatin, ghrelin, IGF-1, thyroid hormone, and glucocorticoids. The somatotroph cell expresses specific receptors for growth hormone-releasing hormone (GHRH), GH secretagogues, and somatotropin release-inhibiting factor (SRIF) receptor subtypes 2 and 5, which all mediate GH secretion. Hypothalamic SRIF and GHRH are secreted in independent waves and interact together with additional GH secretagogues to generate pulsatile GH release. GH secretion is further regulated by IGF-1, which participates in a hypothalamic-pituitary peripheral regulatory feedback system [411]. Norepinephrine, insulin-induced hypoglycemia, clonidine, arginine administration, and exercise all facilitate GH secretion [412]. Emotional deprivation and endogenous depression suppress GH secretion. Acute glucocorticoid administration stimulates GH secretion, but chronic steroid treatment inhibits GH. Hypothyroidism impaired GH response to stimulation and normalized when thyroid status is restored. Gonadal steroids regulate GH secretion and GH action in men and women. Nutrition plays a major role in GH regulation. Malnutrition increases GH secretion, whereas obesity decreases GH secretion. These nutritional effects occur acutely, exemplified by fasting state [413].

Secretion GH secretion is episodic and exhibits a diurnal rhythm with approximately two thirds of GH secretion produced at night by the onset of slow-wave sleep. This accounts for 70% of daily GH secretion [414]. Normal GH secretion is characterized by secretory episodes separated by troughs of minimal basal secretion during which GH is undetectable. GH concentration is highest in the fetal circulation peaking at approximately 150 $\mu\text{g/L}$, and neonatal levels are lower ($\sim 30 \mu\text{g/L}$) reflecting the negative feedback control by rising levels of circulating IGF-1. GH output falls to a stable level during childhood

(prepubertal state levels 200–600 $\mu\text{g/day}$), rising to peak at a twofold to threefold level at the peak of puberty (1000–1800 $\mu\text{g/day}$) [412]. GH output declines exponentially in both sexes at the young adulthood transition, declining to one quarter of the values achieved in late puberty. The decline in GH status occurs by a change in pulse amplitude rather than frequency. Adiposity accounts for a significant component of declining GH output with increasing age [398].

Function GH is the most abundant hormone in the adult pituitary gland, and GH functions as a major metabolic hormone. Metabolic actions of GH promote fat metabolism by enhancing lipolysis and fatty acid oxidation and regulate lipoprotein metabolism by enhancing low-density lipoprotein (LDL) clearance [415]. GH enhances glucose uptake and utilization in cells and suppresses glucose oxidation and utilization while enhancing hepatic glucose production [416]. GH reduces protein oxidation and stimulates protein synthesis. Growth hormone (GH) stimulates the production of insulin-like growth factor 1 (IGF-1), also called somatomedin C; it is a primary mediator of the effects of growth hormone (GH) systemically [411]. IGF-1 stimulates systemic body growth and has growth-promoting effects on almost every cell in the body, especially the skeletal muscle, cartilage, bone, liver, kidney, nerve, skin, hematopoietic, and lung cells.

Adrenocorticotrophic Hormone (ACTH)

Physiology Corticotroph cells comprise about 20% of functional anterior pituitary cells; they are clustered mainly in the central median pituitary wedge. They are the earliest detectable human fetal pituitary cell type, appearing by the eighth week of gestation. These cells produce POMC peptide which is the precursor for ACTH, which acts on the adrenal glands to induce synthesis and secretion of adrenal steroids. Proopiomelanocortin (POMC) is a 266-amino acid pre-prohormone molecule encoding ACTH (1–39), β -lipotropin (LPH), and endorphins. The 8-kb human POMC gene is located on chromo-

some 2p23 [417]. It consists of three exons, first exon encodes a leader sequence, the second encodes the signal initiation sequence along with N-terminal residues of the precursor peptide, and the third exon encodes most of the mature peptide sequences including ACTH and β -LPH [418]. The POMC gene is expressed in pituitary and nonpituitary tissues including the brain, skin, placenta, gonads, gastrointestinal tissues, liver, kidney, adrenal medulla, lung, and lymphocytes. Corticotroph expression is determined by the pituitary promoter interaction (upstream) with exon 1, whereas peripheral expression of the short POMC mRNA is determined by exon 3 (downstream) promoter [419]. Multiple signals act to activate POMC gene expression, including corticotropin-releasing hormone (CRH), cytokines, arginine vasopressin (AVP), catecholamines, and vasoactive intestinal peptide (VIP), cAMP, activator protein-1 (AP1), and glucocorticoids [420]. The CRH type 1 receptor is predominantly expressed on the corticotroph, and receptor activation increases cAMP, protein kinase A, and CREB induction of CRH-binding protein (CRHBP) leading to POMC transcription [421]. After transcription POMC required several post-translational modifications before hormone secretion, including removal N-terminal signal sequence, glycosylation via an *O*-linkage, and serine phosphorylation [422]. POMC cleaved by prohormone convertase 1 (PC1) and prohormone convertase 2 (PC2) into ACTH, melanocyte-stimulating hormone, β -lipotropin (LPH), and endorphins. PC1 is most abundant in the pituitary and hypothalamus, whereas PC2 is present in the CNS, skin, and pancreatic islets but is absent in the pituitary. PC1 expression results in cleavage of POMC limited to four sites, with ACTH being a major product. In the hypothalamus and CNS, both PC1 and PC2 allow coordinated proteolysis, resulting in the generation of smaller fragments such as melanocyte-stimulating hormone (α -, β -, and γ -MSH), β -endorphin, and corticotropin-like intermediate lobe peptide (CLIP) [422].

Function ACTH is a polypeptide of 39 amino acids with a molecular weight of 4.5 kDa. Full-length ACTH is the only POMC-derived peptide

with adrenocorticotroph function; it is the ligand of the melanocortin receptor type 2 receptor (MC2R) [392]. MC2R activation results in production of adrenal glucocorticoids, androgenic steroids, and mineralocorticoids. ACTH signals via adenylyl cyclase to regulate P450 enzyme transcription of cortisol, aldosterone, 17-hydroxyprogesterone, and, to a lesser extent, adrenal androgens [423].

Regulation ACTH regulation involves at least three tiers of control: First, the brain and hypothalamus release regulatory molecules (including CRH, vasopressin, and dopamine) that directly regulate corticotroph function. Second, intrapituitary cytokines and growth factors act locally to regulate ACTH. Third, glucocorticoids maintain regulatory feedback control of corticotroph secretion by rapidly inhibiting hypothalamic CRH and pituitary ACTH secretion [392]. A tightly controlled immuno-neuroendocrine interface regulates the ACTH response to peripheral stressors, like pain, infection, inflammation, hemorrhage, hypovolemia, trauma, psychological stress, and hypoglycemia. These signals vary in ability to generate ACTH secretion and glucocorticoid response to ACTH.

Secretion ACTH is secreted with both circadian periodicity and ultradian pulsatility under the control of the suprachiasmatic nucleus [392]. This centrally controlled pattern is influenced by peripheral corticosteroids, given chronic glucocorticoid exposure (>24 h) with lead to HPA suppression persisting for days or longer. The circadian pattern of ACTH secretion typically begins at about 4 am, peaking before 7 am, with both ACTH and adrenal steroid levels reaching their nadir between 11 pm and 3 am [392]. In this 24-hour diurnal cycle, periodic ACTH secretory bursts occur at a frequency of 40 pulses per 24 h, with amplitude rather than frequency modulation which contributes to diurnal changes in ACTH [424]. ACTH circadian rhythm is controlled by visual cues and the light-dark cycle and is centrally controlled by CRH. CRH signal determines ACTH response, continuous signal desensitizing the ACTH response while a pulsatile CRH signal

restoring cortisol secretion. Endogenous and exogenous stress, including hypoglycemia, act centrally to increase ACTH pulse amplitude [425]. Daily ACTH secretion is higher in males, who also exhibit higher pulse frequency and peak amplitudes.

Gonadotropins Luteinizing hormone (LH) and follicle-stimulating hormone (FSH)

Physiology Gonadotroph cells secrete luteinizing hormone (LH) and follicle-stimulating hormone (FSH); it is part of the hypothalamic-pituitary-gonadal (HPG) axis. Gonadotroph cells comprise about 10–15% of the functional anterior pituitary cells [392]. LH and FSH act on the ovaries and testes to direct gametogenesis and sex steroid hormone synthesis. The synthesis and secretion of LH and FSH are under complex regulation by hypothalamic input (gonadotropin-releasing hormone, or GnRH), by positive and negative feedback from gonadal sex steroid and peptide hormones, and by paracrine modulation from local factors produced within the pituitary gland. The four heterodimeric glycoprotein hormones, LH, FSH, TSH, and hCG, share structural homology, having evolved from a common ancestral gene. While both the homologous LH and FSH molecules are co-secreted by gonadotrophs, their regulatory mechanisms are not similar. The α GSU, LH β , and FSH β subunits are encoded by different genes, located on chromosomes 6, 11, and 19. The LH/CG β gene cluster comprises seven genes, one gene encodes LH β , one encodes CG β , and the remainder are pseudogenes. The LH β and CG β genes have different promoters and transcriptional start sites. LH β includes a 24-amino acid signal peptide followed by a 121-amino acid mature protein. The FSH β gene, on chromosome 11, is organized similarly to the other glycoprotein hormone β genes, encoding a mature peptide of 111 amino acids, with 2 glycosylation sites, and like LH β , it is expressed only in gonadotrophs. The heterodimeric structure of the subunit of LH and FSH is essential for biologic activity. Disulfide linkages within each subunit result in a tertiary structure that allows and maintains non-

covalent heterodimerization, forming the ultra-structure of the mature folded molecule to facilitate specific ligand-receptor interaction [426].

Regulation LH and FSH secretion patterns and in general the hypothalamic-pituitary-gonadal (HPG) axis are mediated primarily via GnRH, paracrine intrapituitary factors (primarily activin and follistatin), and peripheral feedback (both gonadal sex steroids and gonadal peptide hormones) [392]. Hypothalamic control of gonadotropin secretion occurs primarily through actions of GnRH. The mechanisms that regulate GnRH secretion have been provided by the study of genetic abnormalities that manifest as GnRH deficiency, defects in GnRH secretion, and defects in GnRH action, all of which lead to pubertal disorders or infertility. Conversely, a genetic defect in MKRN3 was noted to cause premature reactivation of GnRH secretion leading to precocious puberty [427]. Many neurotransmitters directly or indirectly modulate GnRH secretion, including norepinephrine, dopamine, serotonin, γ -aminobutyric acid (GABA), glutamate, opiates, neuropeptide Y (NPY), and galanin. Glutamate and norepinephrine provide stimulatory drive, whereas GABA and opioid peptides are inhibitory [392]. Kisspeptins, neurokinin B (NKB), and substance P are key to modulate GnRH secretion. Three gonadal peptides (inhibins, activins, and follistatin) are selective for FSH in terms of their effects on gonadotropins and serve as an additional mechanism for the differential control of FSH and LH. Inhibins, originating in the gonads and acting on the pituitary downregulate FSH synthesis and inhibit FSH secretion [428]. Activins act locally in the pituitary as autocrine/paracrine factors and play an important role to enhance FSH biosynthesis and secretion [428]. Follistatin, an activin-binding protein, inhibits activin action by interfering with activin binding to its receptor [429]. Leptin, a product of peripheral adipose tissue, is a positive regulator of the HPG axis and enables a pivotal link between body fat and reproduction [430]. Nutritional, metabolic, stress, and circadian inputs all appear

to act through these peptides to modulate GnRH, gonadotropin secretion, and the activity of the HPG axis. Gonadal steroid hormones (estrogens, progesterones, and androgens) all have effects on gonadotropins which occur both directly at the level of the gonadotroph and indirectly via effects at the hypothalamus that modulate GnRH secretion.

Secretion GnRH secretion into the hypophyseal portal circulation is pulsatile, resulting in episodic stimulation of the gonadotroph. GnRH signaling initiates with recognition by its receptor, GnRHR. GnRHR activation increases calcium mobilization and stimulates influx of extracellular calcium to induce pituitary LH and FSH secretion [431]. The patterns of GnRH signaling (amplitude, frequency, and contour) are important as each of these characteristics can influence gonadotroph responses allowing for two functionally distinct gonadotropins to be differentially regulated by a single hypothalamic-releasing hormone. The characteristic secretory episodes for LH and FSH indicate daily production rates of 1000 IU and 200 IU, with a half-life of 90 and 500 min for each respective β -subunit [432], with FSH having a longer circulating half-life than that of LH.

Function The primary targets of FSH and LH are the gonads; therefore the effects differ in the male and female. In female, FSH acts on FSH receptors in granulosa cells to facilitate follicular growth and estradiol biosynthesis. In response to FSH, follicles convert androstenedione to estradiol by aromatase activity [433]. FSH also controls granulosa cell production of inhibin during the follicular phase and induces LH receptor expression in granulosa cells of large preovulatory follicles. FSH also initiates the recruitment of the next generation of follicles for subsequent cycles. LH is a major regulator of ovarian steroid synthesis. LH stimulates estrogen production by promoting synthesis of androgen precursors in theca cells, which then diffuse into neighboring granulosa cells, where they are aromatized into estrogens under the control of FSH [434]. The midcycle LH surge initiates the rupture of the

ovulatory follicle and ovulation and induces conversion of the follicle wall into the corpus luteum. LH helps to sustain luteinization by stimulating progesterone synthesis [433].

In male, LH acts on LH receptors in Leydig cells to induce intratesticular testosterone synthesis. FSH binds to FSH receptors on Sertoli cells and stimulates the production of inhibins, androgen-binding protein, androgen receptor, and other proteins. FSH mediates the maturation of spermatids into mature spermatozoa in concert with testosterone [435].

Thyroid-Stimulating Hormone (TSH)

Physiology Thyroid-stimulating hormone (TSH) is made by the thyrotroph cells, which comprise approximately 5% of the functional anterior pituitary cells. They are situated predominantly in the anteromedial areas of the gland. They are smaller than the other cell types and are irregularly shaped. TSH is a glycoprotein hormone comprising a 28-kDa heterodimer of two noncovalently linked α - and β -subunits. The tertiary TSH structure comprises three hairpin loops separated by central disulfide bonds [436]. The 13.5-kb α -subunit gene is located on chromosome 6 and comprises four exons and three introns. The α -subunit is common to TSH, LH, FSH, and hCG, but its regulation is uniquely cell-specific; upstream promoter elements are required for thyrotroph-specific expression [437]. The 4.9-kb TSH β -subunit gene located on chromosome 1 comprises three exons and two introns; it is unique and confers specificity of action [437]. Production of the mature heterodimeric TSH molecule requires complex cotranslational glycosylation and folding of nascent α - and β -subunits [438].

Regulation The thyrotropin-releasing hormone (TRH) neuron plays a central role in determining the set point of the hypothalamic-pituitary-thyroid axis by regulating pituitary TSH release [439]. Three main neuronal groups mediate the effects on hypothalamic TRH neurons: First is adrenergic input from the medulla

in response to the effects of cold exposure, as catecholamines increase the set point for inhibition of TRH gene expression by T₃ [440]. The second group, TRH neurons, receive projections from the arcuate nucleus and contain two leptin-responsive groups: POMC system and NPY/agouti-related protein (AGRP) system regulating energy homeostasis [440]. Third, the hypothalamic dorsomedial nucleus projects to the paraventricular nucleus which represents alternative pathways by which leptin acts to regulate TRH neurons [440].

Intrapituitary TSH is stored in secretory granules and the mature hormone is released primarily in response to hypothalamic TRH. Both α - and β -TSH subunit gene transcriptions are induced by TRH. Feedback regulation by thyroid hormones on TRH and TSH is elaborated through a complex system of paracrine control. The effects of thyroid hormones are mediated by thyroid hormone receptors (TRs). TRs exist as two major isoforms, TR α and TR β . TR α is the key isoform responsible for T₃-mediated negative-feedback regulation by hypophysiologic TRH neurons [441]. The local availability of T₃ is determined by deiodinase 2, which generates T₃ from circulating thyroxine (T₄) [439].

Other factors affecting TSH: Somatotropin release-inhibiting factor (SRIF) inhibits TSH pulse amplitude and blocks the nocturnal TSH surge directly at the pituitary level [442]. Dopamine infusions suppress TSH pulse amplitude by 70% and abrogate the nocturnal TSH surge; however prolonged use of dopamine agonists, however, does not result in hypothyroidism [443]. Certain medications, such as glucocorticoids and NSAIDs, also play a role in suppressing TSH [444].

Secretion Daily TSH production is approximately 100–400 mU with a calculated circulating half-life of approximately 50 min [445]. Secretion rates are enhanced up to 15-fold in hypothyroidism and are suppressed in hyperthyroidism. The degree of TSH glycosylation determines both metabolic clearance rate and bioactivity, and in hypothyroidism, the molecule appears highly sialylated, enhancing bioactivity [438]. TSH

secretion is pulsatile with the low pulse amplitudes along with long TSH half-life result in modest circulating variances; it is amplified in hypothyroidism and abrogated in critical illness [446]. Secretory pulses every 2–3 h are interspersed with periods of tonic, nonpulsatile TSH secretion. Circadian TSH secretion peaks between 11 pm and 5 am, mainly due to increased pulse amplitude that is not sleep-entrained [447]. The 24-hour TSH secretion is stable and robust and not influenced by sex, body mass index, and age [448].

Function Hypothalamic-pituitary-thyroid system plays a critical role in development, growth, and cellular metabolism with thyroid hormone. TSH induces thyroid hormone synthesis and release and maintains trophic thyroid cell integrity.

Posterior Pituitary and Stalk

Anatomy

The posterior pituitary or neurohypophysis is composed of the pars nervosa (also known as the neural or posterior lobe), the infundibular stalk, and the median eminence. The infundibular stalk is surrounded by the pars tuberalis, and together they constitute the *hypophyseal stalk*. The pituitary stalk serves as an anatomic and functional connection to the hypothalamus by connecting the pituitary gland to the median eminence of the hypothalamus. The posterior pituitary is not glandular in nature like the anterior pituitary. It is neural tissue that is made up by the conglomeration of distal axon terminals from the hypothalamic magnocellular neurons. The cell bodies of these neurons are located in paired SON and paired PVN of the hypothalamus. During embryogenesis neuroepithelial cells of the lining of the third ventricle mature into magnocellular neurons and migrate laterally to and above the optic chiasm forming the SON and to both walls of the third ventricle to form the PVN [449]. Axons of the SON join axons of the PVN and course to the basal hypothalamus where they join

the axons from the other side and course through the infundibular stalk to the axon terminals in the posterior pituitary.

Axon terminals of the magnocellular neurons contain neurosecretory granules, membrane-bound packets of precursor hormones stored for subsequent release. The precursor proteins traverse the endoplasmic reticulum and the Golgi apparatus to be packaged in secretory granules [450]. The neurosecretory granules then travel along microtubules down the long axons through the stalk of the infundibulum to the posterior pituitary where the granules are stored. During transport, peptide enzymes (peptidases) within the neurosecretory granules cleave the prohormone into the hormone: vasopressin (ADH) or oxytocin, the carrier protein (neurophysin), and (for vasopressin) the glycopeptide. Blood supply for the posterior pituitary comes directly from the inferior hypophyseal arteries and branches of the posterior communicating and internal carotid arteries. The drainage is into the cavernous sinus and internal jugular vein.

The hormones of the posterior pituitary, oxytocin and vasopressin, are for the most part synthesized in individual hormone-specific magnocellular neurons, within the paraventricular nuclei and supraoptic nuclei, allowing stimulation of hormone-specific neurons. The supraoptic nucleus magnocellular neurons mainly produce vasopressin, around 80–90% and virtually all axons projecting to the posterior pituitary [449]. The organization of the paraventricular nuclei, however, is much more complex. There are five subnuclei and parvocellular divisions that synthesize other peptides, such as corticotropin-releasing hormone (CRH), thyrotropin-releasing hormone (TRH), somatostatin, and opioids [451–453]. The parvocellular neurons project to the median eminence, brainstem, and spinal cord, where they play a role in a variety of neuroendocrine autonomic functions [380]. The SCN, which is located in the midline at the base of and anterior to the third ventricle, also synthesizes vasopressin and controls circadian rhythms as well as seasonal rhythms [451].

Posterior Pituitary Hormones: Synthesis and Secretion

Vasopressin and oxytocin are nonapeptides that are synthesized in the cell bodies of the magnocellular neurons as part of a larger precursor molecule consisting of 6-amino acid ring with a cysteine-to-cysteine bridge and a 3-amino acid tail [454]. The hormones consist of the precursor molecule and a hormone-specific neurophysin and for vasopressin a glycopeptide. Subsequently the precursor molecule is packaged in neurosecretory granules and cleaved to the products during transport to the posterior pituitary. When a stimulus for secretion of vasopressin or oxytocin acts on the appropriate magnocellular cell body, an action potential is generated and propagates down the long axon to the posterior pituitary. The action potential causes an influx of calcium, which induces neurosecretory granules to fuse with the cell membrane, extruding the contents of the neurosecretory granule into the perivascular space and later into the capillary system of the posterior pituitary. Thus, there is coordination of stimulated release of hormone, transport of hormone, and synthesis of new hormone. The control of hormone synthesis is at the level of transcription. Stimuli for secretion of vasopressin or oxytocin also stimulate transcription and increase the mRNA content in the magnocellular neurons. When synthesis is turned off, transport stops, and when synthesis is increased, transport is upregulated [455].

Antidiuretic Hormone (ADH)/ Vasopressin

The physiologic regulation of vasopressin synthesis and secretion involves two systems: osmotic and pressure/volume. There are separate systems at the level of the receptors on the end organs of response. The V_{1a} receptors on blood vessels are distinct from V_2 receptors on renal collecting duct epithelia. V_2 receptors also regulate the nontraditional action of vasopressin to stimulate factor VIII and von Willebrand fac-

tor production. A third receptor, V_{1b} , is responsible for the nontraditional biologic action of vasopressin to stimulate ACTH secretion from the anterior pituitary and has been found in numerous peripheral tissues and areas of the brain [456].

Physiology of Volume and Pressure Regulation

Vasopressin is the main hormone involved in the regulation of water homeostasis and osmolality along with thirst. The renin-angiotensin-aldosterone system (RAAS) is mainly responsible for the regulation of blood pressure and volume [450]. Vasopressin plays a smaller role in regulation of blood pressure and volume when compared to the RAAS. High-pressure arterial baroreceptors are located in the carotid sinus and aortic arch, and low-pressure volume receptors are located in the atria and pulmonary venous system [457]. The afferent signals from these receptors are carried to the brainstem through cranial nerves IX and X. Volume/pressure regulation by vasopressin operates through V_1 receptors on blood vessels. When blood pressure is low and volume is low, activation of V_1 receptors causes contraction of vascular smooth muscle to raise blood pressure and constrict intravascular volume around the available fluid volume, to effectively increase plasma volume and reestablish the inhibition of secretion of vasopressin. Baroreceptors and volume receptors normally inhibit the magnocellular neurons causing decreased secretion of vasopressin. The opposite is true for a decrease in tonic inhibition, blood pressure or volume, will result in stimulation of the magnocellular neurons and release of vasopressin.

Vasopressin's action at the kidney to retain water will help replace volume, but to a lesser extent when compared to the RAAS which is the major hormonal regulation for volume control by stimulating sodium reabsorption in the kidney. This is in part due to volume/baroreceptor responses which are much less sensitive than are the osmoreceptors. The lesser response is attrib-

uted to the fact that changes in blood volume and central venous pressure have little effect to increase vasopressin as long as arterial pressure is maintained by alternative regulatory mechanisms such as RAAS and sympathetic reflexes [457]. A decrease in volume or pressure of around 10–15% is necessary before there is a measurable increase in plasma vasopressin. By counterregulatory effects increases in pressure and central volume will decrease vasopressin secretion [458].

Physiology of Osmotic Regulation

The primary receptors for sensing changes in osmolality are located in the brain. Experimental brain lesions in animals have strongly implicated cells in the OVLT and areas of the anterior hypothalamus near the anterior wall of the third cerebral ventricle as the primary osmoreceptors [454]. Surgical destruction or patients with brain damage affecting the OVLT have inability to maintain normal osmolality due to abolished vasopressin secretion and thirst responses to hyperosmolality but no alteration in their responses to hypovolemia [459]. In contrast, destruction of the magnocellular neurons of the SON and PVN eliminates dehydration-induced secretion of vasopressin but does not alter thirst, indicating that osmotically stimulated thirst is generated at a site proximal to the magnocellular cells.

Extracellular fluid (ECF) osmolality is predominantly determined by sodium concentration. Osmolality varies from 280 to 295 mOsm/kg H_2O in normal subjects, but in any individual it is maintained within a narrower range by three different mechanisms: sensitive response of plasma vasopressin to changes in plasma osmolality, the sensitive response of urine osmolality to changes in plasma vasopressin, and then the gain in the system by the response of urine volume to change in plasma vasopressin [454]. Basal plasma vasopressin is in the range of 0.5–2 pg/mL. As little as a 1% increase or decrease in plasma osmolality will cause a rapid increase or decrease of vasopressin released from the store of hormone in the

posterior pituitary [460]. Rapid metabolism of vasopressin is also characteristic of the hormone, which circulates in plasma with a half-life of approximately 15 min, and this allows rapid changes in levels of vasopressin in plasma. In humans a direct linear relationship exists between plasma osmolality and plasma vasopressin. The opposite is true for the relationship of plasma vasopressin to urine volume. Thus, small increases in plasma osmolality produce a concentrated urine, and small decreases produce a water diuresis.

The kidney conserves water by the combined functions of the loop of Henle and the collecting duct. The loop of Henle generates a high osmolality in the renal medulla via the countercurrent multiplier system. Vasopressin acts in the collecting duct and increases water and urea permeability, allowing osmotic equilibration between the urine and the hypertonic medullary interstitium. The net effect of this process is to extract water from urine from the medulla, through interstitial blood vessels, the vasa recta resulting in increased urine concentration and decreased urine volume. The antidiuretic effect of vasopressin occurs when binding to V_2 receptors on the epithelial principal cells of the renal collecting tubule. Binding activates adenylate cyclase, increasing cyclic adenosine monophosphate (cAMP), which then stimulates protein kinase A. This process leads to phosphorylation and activation of aquaporin 2 and movement of the water channels into the luminal membrane. In addition to shifting water from the collecting duct into the hypertonic inner medulla and concentration of urine, binding also promotes activation of the V_2 receptor increasing the synthesis of aquaporin 2 and the permeability of aquaporin 2 to water [461]. Aquaporins 3 and 4 are constitutively synthesized and are expressed at high levels in the basolateral plasma membranes of principal cells, where they are responsible for the high water permeability of the basolateral plasma membrane [462, 463]. Dissociation of vasopressin from the V_2 receptor allows intracellular cAMP levels to decrease, and the water channels are then reinternalized, terminating the increased water permeability. The aquaporin-containing vesicles remain

just below the apical membrane and can be quickly shuttled into and out of the membrane in response to changes in intracellular cAMP levels. This mechanism allows minute-to-minute regulation of renal water excretion in response to changes in ambient levels of vasopressin in plasma.

During prolonged periods of dehydration (at least 24 h), long-term regulation of collecting duct water permeability occurs in response to chronically elevated levels of circulating vasopressin. Persistently elevated levels of vasopressin upregulate synthesis of aquaporin 2 and aquaporin 3 water channels in the collecting duct principal cells, thus achieving maximum permeability and water conservation. In response urine volume can be reduced to a minimum but not eliminated. To maintain water homeostasis, water must also be consumed to replace the obligate urinary and insensible fluid losses. This is regulated by thirst. Similar to vasopressin, thirst can be stimulated by increases in osmolality of the ECF or by decreases in intravascular volume.

Oxytocin

The classic roles of oxytocin are smooth muscle activation promoting milk let-down with nursing and uterine myometrial contraction at parturition [450]. Difficulty to carry studies in pregnant women and human tissue makes physiologic regulation of oxytocin secretion and function a less well-known subject when comparing humans to other mammals. All mammals secrete oxytocin to stimulate milk let-down associated with nursing, unique characteristic among all mammals. The milk-producing unit of the breast is the alveolar system in which clusters of milk-producing cells are surrounded by specialized myoepithelial cells. Oxytocin receptors are localized on glandular cells of alveoli which synthesize milk. Glandular cells are surrounded by myoepithelial specialized cells that are connected to ductules and then ducts that lead to the nipple. Oxytocin in the systemic circulation acts on these receptors to cause myoepithelial contraction along with shortening and widening of the ducts

to causing milk secretion and enhancing flow. During lactation sucking at the breast triggers an afferent signal that is transmitted from the mechanoreceptors or tactile receptors in the breast to the spinal cord ascending to the oxytocinergic magnocellular neurons in the supraoptic nucleus and the paraventricular nucleus, where neurotransmitters trigger oxytocin secretion. Pulsatile release of the hormone produces a pulsatile pumping action on the alveoli, promoting maximum emptying of milk from the alveoli [464].

Oxytocin also plays a role in parturition, mainly uterine contraction during expulsion stage. During early labor, upregulation in the uterus of oxytocin receptor mRNA occurs, and oxytocin receptor numbers increase [465, 466]. Oxytocin receptors are prominent in the fundus of the uterus, where they stimulate myometrial contraction, and in decidual cells, where they stimulate the production of prostaglandins. At parturition increased oxytocin activity in the fundus will push the fetus toward the cervix, which is thinned and relaxed by the effects of prostaglandins.


Figures 5.18 and 5.19 provide a pictorial view of the various anatomic subsections and blood supply organization of the pituitary gland.

Other Aspects of the Hypophysis Cerebri

Comparative anatomical studies between various mammals as well as other vertebrates and human pituitary glands have provided some insight into other less well-understood and appreciated sections of the pituitary gland. Furthermore noted age-related differentials between fetal, neonatal, childhood, adult, and geriatric configurations of human pituitary have also provided additional insight in this regard [467–469].

The hypophysis (pituitary) is connected to the hypothalamus via the pituitary stalk which has a neurohypophyseal component: the *infundibulum*. This is a sheath of projecting axons from several hypothalamic nuclei as previously detailed above. The pituitary stalk however also has an adenohypophyseal component which forms and encircling tube around the infundibulum and is called the *pars tuberalis*. There is some suggestion that the *pars tuberalis* may have a specialty function beyond being a more superiorly located location for anterior pituitary hormonal secretory cells. It appears to in addition serve as the primary transducer for long-term internal timers and is sometimes referred to as the “circannual clock” (as opposed to the SCNs established role as the

Fig. 5.18 Pictorial of the human pituitary gland; hypophysis cerebri






Fig. 5.19 Schema of the blood supply of the pituitary gland

circadian biologic clock center) [470]. In this capacity the pars tuberalis appears to play a role in establishing and regulating seasonal rhythmicity of various autonomic and other physiologic functions [470–473].

The *pars intermedia* is the small sliver of tissue that lies between the adenohypophysis and neurohypophysis. It contains three main groups of cells: chromophobes, basophils, and colloid lining secretory cells that line several colloid-filled cysts which are vestiges of the developmental Rathke's pouch [467–473]. Human fetal pituitary glands as well as the pituitary from fish and amphibians produce MSH in abundance. While this is important for proper skin pigmentation development in human fetuses in the lower vertebrates, it has a more complex and important role in adult animals including environmental-induced skin pigmentation for camouflage purposes as well as for playing a role in sexual and reproductive mating behaviors. In human adults however, it is typically vestigial or completely absent, and it is unclear that it has any significant physiologic functional role in adults. The cells of the *pars intermedia* are highly metabolically

active and especially demonstrable in bovines demonstrate a triphasic cyclic activity routine characterized by (a) resting cellular phase, (b) desquamation with autolysis, and (c) cellular regeneration. The desquamated cuboidal secretory cells produce the colloid which fills the colloid cysts present in the *pars intermedia*, and from here the colloid is drained by the extensive capillaries traveling between the neuro- and adenohypophysis with subsequent venous drainage via the inferior pituitary venules.

To varying degrees human pituitary glands can also demonstrate a well-defined *interglandular cleft* that lies between the *pars intermedia* and the adenohypophysis. It is a true glandular lumen lined with secretory cuboidal cells. While well defined in some mammals and in the fetal pituitary, its presence in adult humans is variable and when excessive can induce pathology due to local mass effect and is then referred to as Rathke's cleft cysts which will be discussed in more depth in other chapters. The exact functional significance of any of the colloid of the interglandular cleft to adult human physiology remains unresolved.

Concluding Remarks

This chapter has attempted to give a comprehensive overview of the roles that the hypothalamus and pituitary play in the normal regulation of various voluntary and autonomic functions and behavior patterns and has sought to show the close integration between specialized neuronal and endocrine functions between the various cells and nuclei of both organs. As a unit the hypothalamic-pituitary axes serve as the highest center of systemic endocrine oversight and modulation.

While our knowledge of this complex system has grown significantly, there still remain unresolved questions from a molecular through cellular to anatomic level of inquiry as regards various aspects of the structure and function of this system. Even better understanding of the interactions between the hypothalamic-pituitary axes and its upward projections to the thalamus, limbic system, cerebellum, and neocortex as well as its downward projections to the midbrain, spinal cord, peripheral nervous system, and systemic circulation will have great implications for improving therapeutics in a wide range of clinical disciplines.

References/Bibliography

- Scharrer E, Scharrer B. Neurosecretion. *Physiol Rev*. 1945;25:171–81.
- Guillemin R. Peptides in the brain: the new endocrinology of the neuron. *Science*. 1978;202(4366):390–402. Epub 1978/10/27.
- Schally AV. Aspects of hypothalamic regulation of the pituitary gland. *Science*. 1978;202(4363):18–28. Epub 1978/10/06.
- Kurrasch DM, Cheung CC, Lee FY, Tran PV, Hata K, Ingraham HA. The neonatal ventromedial hypothalamus transcriptome reveals novel markers with spatially distinct patterning. *J Neurosci*. 2007;27(50):13624–34. Epub 2007/12/14.
- Tobet SA. Genes controlling hypothalamic development and sexual differentiation. *Eur J Neurosci*. 2002;16(3):373–6. Epub 2002/08/24.
- Markakis EA. Development of the neuroendocrine hypothalamus. *Front Neuroendocrinol*. 2002;23(3):257–91. Epub 2002/07/20.
- McNay DE, Pelling M, Claxton S, Guillemot F, Ang SL. *Mash1* is required for generic and subtype differentiation of hypothalamic neuroendocrine cells. *Mol Endocrinol*. 2006;20(7):1623–32. Epub 2006/02/14.
- Yu S, Francois M, Huesing C, Munzberg H. The hypothalamic preoptic area and body weight control. *Neuroendocrinology*. 2018;106(2):187–94. Epub 2017/08/05.
- Cornejo MP, Hentges ST, Maliqueo M, Coirini H, Becu-Villalobos D, Elias CF. Neuroendocrine regulation of metabolism. *J Neuroendocrinol*. 2016;28(7). Epub 2016/04/27.
- Schneeberger M, Gomis R, Claret M. Hypothalamic and brainstem neuronal circuits controlling homeostatic energy balance. *J Endocrinol*. 2014;220(2):T25–46. Epub 2013/11/14.
- van Dijk G, Evers SS, Guidotti S, Thornton SN, Scheurink AJ, Nyakas C. The lateral hypothalamus: a site for integration of nutrient and fluid balance. *Behav Brain Res*. 2011;221(2):481–7. Epub 2011/02/09.
- Donato J Jr, Cravo RM, Frazao R, Elias CF. Hypothalamic sites of leptin action linking metabolism and reproduction. *Neuroendocrinology*. 2011;93(1):9–18. Epub 2010/11/26.
- Plant TM. Hypothalamic control of the pituitary-gonadal axis in higher primates: key advances over the last two decades. *J Neuroendocrinol*. 2008;20(6):719–26. Epub 2008/07/08.
- Harrold JA. Hypothalamic control of energy balance. *Curr Drug Targets*. 2004;5(3):207–19. Epub 2004/04/03.
- Muller MB, Uhr M, Holsboer F, Keck ME. Hypothalamic-pituitary-adrenocortical system and mood disorders: highlights from mutant mice. *Neuroendocrinology*. 2004;79(1):1–12. Epub 2004/02/03.
- Bernardis LL, Bellinger LL. The lateral hypothalamic area revisited: neuroanatomy, body weight regulation, neuroendocrinology and metabolism. *Neurosci Biobehav Rev*. 1993;17(2):141–93. Epub 1993/01/01.
- Smith R, Thomson M. Neuroendocrinology of the hypothalamo-pituitary-adrenal axis in pregnancy and the puerperium. *Baillieres Clin Endocrinol Metab*. 1991;5(1):167–86. Epub 1991/03/01.
- Veldhuis JD. The hypothalamic pulse generator: the reproductive core. *Clin Obstet Gynecol*. 1990;33(3):538–50. Epub 1990/09/01.
- Reichlin S. Neuroendocrinology of the pituitary gland. *Toxicol Pathol*. 1989;17(2):250–5. Epub 1989/01/01.
- Arimura A, Fishback JB. Somatostatin: regulation of secretion. *Neuroendocrinology*. 1981;33(4):246–56. Epub 1981/01/01.
- Almeida OF, Hassan AH, Holsboer F. Intrahypothalamic neuroendocrine actions of corticotropin-releasing factor. *Ciba Found Symp*. 1993;172:151–69. Discussion 69–72. Epub 1993/01/01.
- Bonfiglio JJ, Inda C, Refojo D, Holsboer F, Arzt E, Silberstein S. The corticotropin-releasing hormone network and the hypothalamic-pituitary-adrenal

- axis: molecular and cellular mechanisms involved. *Neuroendocrinology*. 2011;94(1):12–20. Epub 2011/05/18.
23. Born J, Fehm HL. Hypothalamus-pituitary-adrenal activity during human sleep: a coordinating role for the limbic hippocampal system. *Exp Clin Endocrinol Diab Off J German Society of Endocrinology [and] German Diabetes Association*. 1998;106(3):153–63. Epub 1998/08/26.
24. Buckingham JC. Stress and the hypothalamo-pituitary-immune axis. *Int J Tissue React*. 1998;20(1):23–34. Epub 1998/04/30.
25. Balthazart J, Ball GF. Topography in the pre-optic region: differential regulation of appetitive and consummatory male sexual behaviors. *Front Neuroendocrinol*. 2007;28(4):161–78. Epub 2007/07/13.
26. Constantin S. Physiology of the gonadotrophin-releasing hormone (GnRH) neurone: studies from embryonic GnRH neurones. *J Neuroendocrinol*. 2011;23(6):542–53. Epub 2011/03/30.
27. Genazzani AR, Petraglia F, Gamba O, Sgarbi L, Greco MM, Genazzani AD. Neuroendocrinology of the menstrual cycle. *Ann N Y Acad Sci*. 1997;816:143–50. Epub 1997/06/17.
28. Harrold JA. Leptin leads hypothalamic feeding circuits in a new direction. *BioEssays*. 2004;26(10):1043–5. Epub 2004/09/24.
29. Plant TM. Neuroendocrine control of the onset of puberty. *Front Neuroendocrinol*. 2015;38:73–88. Epub 2015/04/29.
30. Belle MD. Circadian tick-talking across the neuroendocrine system and suprachiasmatic nuclei circuits: the enigmatic communication between the molecular and electrical membrane clocks. *J Neuroendocrinol*. 2015;27(7):567–76. Epub 2015/04/08.
31. Brown CH, Bourque CW. Mechanisms of rhythmogenesis: insights from hypothalamic vasopressin neurons. *Trends Neurosci*. 2006;29(2):108–15. Epub 2005/12/07.
32. Rubin RT, Poland RE, Rubin LE, Gouin PR. The neuroendocrinology of human sleep. *Life Sci*. 1974;14(6):1041–52. Epub 1974/03/16.
33. Hofman MA, Swaab DF. The sexually dimorphic nucleus of the preoptic area in the human brain: a comparative morphometric study. *J Anat*. 1989;164:55–72. Epub 1989/06/01.
34. Quinlins KM, Bonthuis PJ, Harris EP, Shetty SR, Rissman EF. Neural growth hormone: regional regulation by estradiol and/or sex chromosome complement in male and female mice. *Biol Sex Differ*. 2015;6:8. Epub 2015/05/20.
35. Castaneya-Ruiz L, Gonzalez-Marrero I, Castaneya-Ruiz A, Gonzalez-Toledo JM, Castaneya-Ruiz M, de Paz-Carmona H, et al. Luteinizing hormone-releasing hormone distribution in the anterior hypothalamus of the female rats. *ISRN Anat*. 2013;2013:870721. Epub 2013/01/01.
36. Isgor C, Cecchi M, Kabbaj M, Akil H, Watson SJ. Estrogen receptor beta in the paraventricular nucleus of hypothalamus regulates the neuroendocrine response to stress and is regulated by corticosterone. *Neuroscience*. 2003;121(4):837–45. Epub 2003/10/29.
37. McCarthy MM, Arnold AP, Ball GF, Blaustein JD, De Vries GJ. Sex differences in the brain: the not so inconvenient truth. *J Neurosci*. 2012;32(7):2241–7. Epub 2012/03/08.
38. Oldfield BJ, McKinley MJ. Circumventricular organs. In: Paxinos G, editor. *The rat nervous system*. Cambridge, Massachusetts: Academic Press, 1995. p. 391–403.
39. Weindl A. Neuroendocrine aspects of the circumventricular organs. In: Ganong W, Martini L, editors. *Frontiers in neuroendocrinology*. Oxford University Press, Oxford, England, UK; 1973. p. 3–32.
40. Johnson AK, Gross PM. Sensory circumventricular organs and brain homeostatic pathways. *FASEB J*. 1993;7(8):678–86. Epub 1993/05/01.
41. Fekete C, Freitas BC, Zeold A, Wittmann G, Kadar A, Liposits Z, et al. Expression patterns of WSB-1 and USP-33 underlie cell-specific posttranslational control of type 2 deiodinase in the rat brain. *Endocrinology*. 2007;148(10):4865–74. Epub 2007/07/14.
42. Kallo I, Mohacsik P, Vida B, Zeold A, Bardoczi Z, Zavacki AM, et al. A novel pathway regulates thyroid hormone availability in rat and human hypothalamic neurosecretory neurons. *PLoS One*. 2012;7(6):e37860. Epub 2012/06/22.
43. Tu HM, Kim SW, Salvatore D, Bartha T, Legradi G, Larsen PR, et al. Regional distribution of type 2 thyroxine deiodinase messenger ribonucleic acid in rat hypothalamus and pituitary and its regulation by thyroid hormone. *Endocrinology*. 1997;138(8):3359–68. Epub 1997/08/01.
44. Sanders NM, Dunn-Meynell AA, Levin BE. Third ventricular alloxan reversibly impairs glucose counterregulatory responses. *Diabetes*. 2004;53(5):1230–6. Epub 2004/04/28.
45. Weindl A, Sofroniew MV. Relation of neuropeptides to mammalian circumventricular organs. In: Martin JB, Reichlin S, Bick KL, editors. *Neurosecretion and brain peptides*. New York, USA: Raven Press, 1981. p. 303–320.
46. McKinley MJ, Burns P, Colvill LM, Oldfield BJ, Wade JD, Weisinger RS, et al. Distribution of Fos immunoreactivity in the lamina terminalis and hypothalamus induced by centrally administered relaxin in conscious rats. *J Neuroendocrinol*. 1997;9(6):431–7. Epub 1997/06/01.
47. Oldfield BJ, Badoer E, Hards DK, McKinley MJ. Fos production in retrogradely labelled neurons of the lamina terminalis following intravenous infusion of either hypertonic saline or angiotensin II. *Neuroscience*. 1994;60(1):255–62. Epub 1994/05/01.

48. Hoffman GE, Gibbs FP. LHRH pathways in rat brain: 'deafferentation' spares a sub-chiasmatic LHRH projection to the median eminence. *Neuroscience*. 1982;7(8):1979–93. Epub 1982/01/01.
49. Dellmann HD. Fine structural organization of the subfornical organ. A concise review. *Brain Res Bull*. 1985;15(1):71–8. Epub 1985/07/01.
50. Dellmann HD, Simpson JB. The subfornical organ. *Int Rev Cytol*. 1979;58:333–421. Epub 1979/01/01.
51. Gross PM. Circumventricular organ capillaries. *Prog Brain Res*. 1992;91:219–33. Epub 1992/01/01.
52. Swanson LW, Mogenson GJ. Neural mechanisms for functional coupling of autonomic, endocrine and somatomotor responses in adaptive behavior. *Brain Res Brain Res Rev*. 1981;3:2–34.
53. Bourque CW, Oliet SH, Richard D. Osmoreceptors, osmoreception, and osmoregulation. *Front Neuroendocrinol*. 1994;15(3):231–74. Epub 1994/09/01.
54. Ferguson AV. Neurophysiological analysis of mechanisms for subfornical organ and area postrema involvement in autonomic control. *Prog Brain Res*. 1992;91:413–21. Epub 1992/01/01.
55. Toni R, Mosca S, Ruggeri F, Valmori A, Orlandi G, Toni G, et al. Effect of hypothyroidism on vasoactive intestinal polypeptide-immunoreactive neurons in forebrain-neurohypophyseal nuclei of the rat brain. *Brain Res*. 1995;682(1–2):101–15. Epub 1995/06/05.
56. Kai A, Ono K, Kawano H, Honda E, Nakanishi O, Inenaga K. Galanin inhibits neural activity in the subfornical organ in rat slice preparation. *Neuroscience*. 2006;143(3):769–77. Epub 2006/10/10.
57. Sakai K, Agassandian K, Morimoto S, Sinnayah P, Cassell MD, Davisson RL, et al. Local production of angiotensin II in the subfornical organ causes elevated drinking. *J Clin Invest*. 2007;117(4):1088–95. Epub 2007/04/04.
58. Gross PM. The subfornical organ as a model of neurohumoral integration. *Brain Res Bull*. 1985;15(1):65–70. Epub 1985/07/01.
59. Hajdu I, Szentirmai E, Obal F Jr, Krueger JM. Different brain structures mediate drinking and sleep suppression elicited by the somatostatin analog, octreotide, in rats. *Brain Res*. 2003;994(1):115–23. Epub 2003/12/04.
60. Kadekaro M, Gross PM. Elevated glucose utilization in the subfornical organ during dehydration. *Brain Res Bull*. 1985;15(1):99–104. Epub 1985/07/01.
61. Price CJ, Hoyda TD, Samson WK, Ferguson AV. Nesfatin-1 influences the excitability of paraventricular nucleus neurones. *J Neuroendocrinol*. 2008;20(2):245–50. Epub 2007/12/20.
62. Price TO, Samson WK, Niehoff ML, Banks WA. Permeability of the blood-brain barrier to a novel satiety molecule nesfatin-1. *Peptides*. 2007;28(12):2372–81. Epub 2007/11/17.
63. Samson WK, White MM, Price C, Ferguson AV. Obestatin acts in brain to inhibit thirst. *Am J Physiol Regul Integr Comp Physiol*. 2007;292(1):R637–43. Epub 2006/08/26.
64. Young CN, Morgan DA, Butler SD, Mark AL, Davisson RL. The brain subfornical organ mediates leptin-induced increases in renal sympathetic activity but not its metabolic effects. *Hypertension*. 2013;61(3):737–44. Epub 2013/01/30.
65. Lee K, Tan J, Morris MB, Rizzoti K, Hughes J, Cheah PS, et al. Congenital hydrocephalus and abnormal subcommissural organ development in Sox3 transgenic mice. *PLoS One*. 2012;7(1):e29041. Epub 2012/02/01.
66. Saha S, Subhedar N. Calcitonin-like immunoreactivity in the subcommissural organ-Reissner's fiber complex of some freshwater and marine teleosts. *J Chem Neuroanat*. 2011;41(2):122–8. Epub 2010/12/28.
67. Elgot A, Ahboucha S, Bouyatas MM, Fevre-Montange M, Gamrani H. Water deprivation affects serotonergic system and glycoprotein secretion in the sub-commissural organ of a desert rodent *Meriones shawi*. *Neurosci Lett*. 2009;466(1):6–10. Epub 2009/09/01.
68. Borison HL. Area postrema: chemoreceptor circumventricular organ of the medulla oblongata. *Prog Neurobiol*. 1989;32(5):351–90. Epub 1989/01/01.
69. Brizzee KR, Klara PM. The structure of the mammalian area postrema. *Fed Proc*. 1984;43(15):2944–8. Epub 1984/12/01.
70. Lavezzi AM, Mecchia D, Maturri L. Neuropathology of the area postrema in sudden intrauterine and infant death syndromes related to tobacco smoke exposure. *Auton Neurosci*. 2012;166(1–2):29–34. Epub 2011/10/11.
71. Gross PM, Wainman DS, Shaver SW, Wall KM, Ferguson AV. Metabolic activation of efferent pathways from the rat area postrema. *Am J Phys*. 1990;258(3 Pt 2):R788–97. Epub 1990/03/01.
72. Shaver SW, Kadekaro M, Gross PM. High metabolic activity in the dorsal vagal complex of Brattleboro rats. *Brain Res*. 1989;505(2):316–20. Epub 1989/12/29.
73. Shaver SW, Kadekaro M, Gross PM. Focal metabolic effects of angiotensin and captopril on subregions of the rat subfornical organ. *Peptides*. 1990;11(3):557–63. Epub 1990/05/01.
74. Shaver SW, Kadekaro M, Gross PM. Differential rates of glucose metabolism across subregions of the subfornical organ in Brattleboro rats. *Regul Pept*. 1990;27(1):37–49. Epub 1990/01/01.
75. Feldberg W, Pyke D, Stubbs WA. Hyperglycaemia: imitating Claude Bernard's pique with drugs. *J Auton Nerv Syst*. 1985;14(3):213–28. Epub 1985/11/01.
76. Li MC. Effect of pique diabétique upon the blood sugar content of rabbits, adrenalectomized or splanchnicotomized. *Tohoku J Exp Med*. 1952;56(4):310. Epub 1952/10/01.
77. Arendt J, Skene DJ. Melatonin as a chronobiotic. *Sleep Med Rev*. 2005;9(1):25–39. Epub 2005/01/15

78. Macchi MM, Bruce JN. Human pineal physiology and functional significance of melatonin. *Front Neuroendocrinol.* 2004;25(3-4):177-95. Epub 2004/12/14.
79. Chen CY, Chen FH, Lee CC, Lee KW, Hsiao HS. Sonographic characteristics of the cavum velum interpositum. *AJNR Am J Neuroradiol.* 1998;19(9):1631-5. Epub 1998/11/05.
80. Arendt J. Melatonin and the mammalian pineal gland. 1st ed. London UK:Chapman and Hall, 1995.
81. Pritchard TC, Alloway KD. Medical neuroscience. Raleigh, NC; Hayes Barton Press, 1999. p. 76-77.
82. Moller M, Baeres FM. The anatomy and innervation of the mammalian pineal gland. *Cell Tissue Res.* 2002;309(1):139-50. Epub 2002/07/12.
83. Kleinschmidt-DeMasters BK, Prayson RA. An algorithmic approach to the brain biopsy -- part I. *Arch Pathol Lab Med.* 2006;130(11):1630-8. Epub 2006/11/02.
84. Prayson RA, Kleinschmidt-DeMasters BK. An algorithmic approach to the brain biopsy--part II. *Arch Pathol Lab Med.* 2006;130(11):1639-48. Epub 2006/11/02.
85. Schmidt F, Penka B, Trauner M, Reinsperger L, Ranner G, Ebner F, et al. Lack of pineal growth during childhood. *J Clin Endocrinol Metab.* 1995;80(4):1221-5. Epub 1995/04/01.
86. Sumida M, Barkovich AJ, Newmont TH. Development of the pineal gland: measurement with MR. *AJNR Am J Neuroradiol.* 1996;17(2):233-6. Epub 1996/02/01.
87. Tapp E, Huxley M. The weight and degree of calcification of the pineal gland. *J Pathol.* 1971;105(1):31-9. Epub 1971/09/01.
88. Tapp E, Huxley M. The histological appearance of the human pineal gland from puberty to old age. *J Pathol.* 1972;108(2):137-44. Epub 1972/10/01.
89. Lerner AB, Case JD, Takahashi Y. Isolation of melatonin and 5-methoxyindole-3-acetic acid from bovine pineal glands. *J Biol Chem.* 1960;235:1992-7. Epub 1960/07/01.
90. Axelrod J. The pineal gland. *Endeavour.* 1970;29(108):144-8. Epub 1970/09/01.
91. Lowrey PL, Takahashi JS. Genetics of the mammalian circadian system: photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation. *Annu Rev Genet.* 2000;34:533-62. Epub 2000/11/28.
92. Airaksinen MM, Sainio EL, Leppaluoto J, Kari I. 6-methoxy-tetrahydro-beta-carboline (pinoline): effects on plasma renin activity and aldosterone, TSH, LH and beta-endorphin levels in rats. *Acta Endocrinol.* 1984;107(4):525-30. Epub 1984/12/01.
93. Aranda M, Albendea CD, Lostale F, Lopez-Pingarron L, Fuentes-Broto L, Martinez-Ballarín E, et al. In vivo hepatic oxidative stress because of carbon tetrachloride toxicity: protection by melatonin and pinoline. *J Pineal Res.* 2010;49(1):78-85. Epub 2010/06/08.
94. Fuentes-Broto L, Miana-Mena FJ, Piedrafita E, Berzosa C, Martínez-Ballarín E, García-Gil FA, et al. Melatonin protects against taurolithocholic-induced oxidative stress in rat liver. *J Cell Biochem.* 2010;110(5):1219-25. Epub 2010/06/22.
95. Langer SZ, Lee CR, Segonzac A, Tateishi T, Esnaud H, Schoemaker H, et al. Possible endocrine role of the pineal gland for 6-methoxytetrahydro-beta-carboline, a putative endogenous neuromodulator of the [3H]imipramine recognition site. *Eur J Pharmacol.* 1984;102(2):379-80. Epub 1984/07/13.
96. Langer SZ, Raisman R, Tahraoui L, Scatton B, Niddam R, Lee CR, et al. Substituted tetrahydro-beta-carbolines are possible candidates as endogenous ligand of the [3H]imipramine recognition site. *Eur J Pharmacol.* 1984;98(1):153-4. Epub 1984/02/10.
97. Millan-Plano S, Piedrafita E, Miana-Mena FJ, Fuentes-Broto L, Martínez-Ballarín E, López-Pingarrón L, et al. Melatonin and structurally-related compounds protect synaptosomal membranes from free radical damage. *Int J Mol Sci.* 2010;11(1):312-28. Epub 2010/02/18.
98. Pakkila R, Zilmer M, Kullisaar T, Rago L. Comparison of the antioxidant activity of melatonin and pinoline in vitro. *J Pineal Res.* 1998;24(2):96-101. Epub 1998/03/24.
99. Pless G, Frederiksen TJ, Garcia JJ, Reiter RJ. Pharmacological aspects of N-acetyl-5-methoxytryptamine (melatonin) and 6-methoxy-1,2,3,4-tetrahydro-beta-carboline (pinoline) as antioxidants: reduction of oxidative damage in brain region homogenates. *J Pineal Res.* 1999;26(4):236-46. Epub 1999/05/26.
100. Siu AW, Reiter RJ, To CH. Pineal indoleamines and vitamin E reduce nitric oxide-induced lipid peroxidation in rat retinal homogenates. *J Pineal Res.* 1999;27(2):122-8. Epub 1999/09/25.
101. Herraiz T, Galisteo J. Endogenous and dietary indoles: a class of antioxidants and radical scavengers in the ABTS assay. *Free Radic Res.* 2004;38(3):323-31. Epub 2004/05/08.
102. Motta M, Fraschini F, Martini L. Endocrine effects of pineal gland and of melatonin. *Proc Soc Exp Biol Med.* 1967;126(2):431-5. Epub 1967/11/01.
103. Naidoo V, Naidoo S, Mahabeer R, Raidoo DM. Cellular distribution of the endothelin system in the human brain. *J Chem Neuroanat.* 2004;27(2):87-98. Epub 2004/05/04.
104. Sharan K, Lewis K, Furukawa T, Yadav VK. Regulation of bone mass through pineal-derived melatonin-MT2 receptor pathway. *J Pineal Res.* 2017;63(2). Epub 2017/05/18.
105. Chew BH, Weaver DF, Gross PM. Dose-related potent brain stimulation by the neuropeptide endothelin-1 after intraventricular administration in conscious rats. *Pharmacol Biochem Behav.* 1995;51(1):37-47. Epub 1995/05/01.

106. Jacobs RA, Satta MA, Dahia PL, Chew SL, Grossman AB. Induction of nitric oxide synthase and interleukin-1 β , but not heme oxygenase, messenger RNA in rat brain following peripheral administration of endotoxin. *Brain Res Mol Brain Res.* 1997;49(1–2):238–46. Epub 1997/12/05.
107. Jacobs RA, Schaad NC, Vanecek J, Leaver S, Aubry JM, Korf HW, et al. Pineal nitric oxide synthase, but not heme oxygenase, mRNA is suppressed by continuous exposure to light. *Brain Res Mol Brain Res.* 1999;70(2):264–72. Epub 1999/07/17.
108. Gross PM, Wainman DS, Chew BH, Espinosa FJ, Weaver DF. Calcium-mediated metabolic stimulation of neuroendocrine structures by intraventricular endothelin-1 in conscious rats. *Brain Res.* 1993;606(1):135–42. Epub 1993/03/19.
109. Del Brutto OH, Mera RM, Lama J, Zambrano M, Castillo PR. Pineal gland calcification is not associated with sleep-related symptoms. A population-based study in community-dwelling elders living in Atahualpa (rural coastal Ecuador). *Sleep Med.* 2014;15(11):1426–7. Epub 2014/10/04.
110. Doyle AJ, Anderson GD. Physiologic calcification of the pineal gland in children on computed tomography: prevalence, observer reliability and association with choroid plexus calcification. *Acad Radiol.* 2006;13(7):822–6. Epub 2006/06/17.
111. Tan DX, Xu B, Zhou X, Reiter RJ. Pineal calcification, melatonin production, aging, associated health consequences and rejuvenation of the pineal gland. *Molecules.* 2018;23(2). Epub 2018/02/01.
112. Vigh B, Szel A, Debreceni K, Fejer Z, Manzano e Silva MJ, Vigh-Teichmann I. Comparative histology of pineal calcification. *Histol Histopathol.* 1998;13(3):851–70. Epub 1998/08/05.
113. Whitehead MT, Oh C, Raju A, Choudhri AF. Physiologic pineal region, choroid plexus, and dural calcifications in the first decade of life. *AJNR Am J Neuroradiol.* 2015;36(3):575–80. Epub 2014/10/31.
114. Flament-Durand J. The hypothalamus: anatomy and functions. *Acta Psychiatr Belg.* 1980;80(4):364–75. Epub 1980/07/01.
115. Hrabovszky E. Neuroanatomy of the human hypothalamic kisspeptin system. *Neuroendocrinology.* 2014;99(1):33–48. Epub 2014/01/10.
116. Morin LP. Neuroanatomy of the extended circadian rhythm system. *Exp Neurol.* 2013;243:4–20. Epub 2012/07/07.
117. Patel CR, Fernandez-Miranda JC, Wang WH, Wang EW. Skull base anatomy. *Otolaryngol Clin N Am.* 2016;49(1):9–20. Epub 2015/11/29.
118. Swaab DF, Hofman MA, Lucassen PJ, Purba JS, Raadsheer FC, Van de Nes JA. Functional neuroanatomy and neuropathology of the human hypothalamus. *Anat Embryol.* 1993;187(4):317–30. Epub 1993/04/01.
119. Watts AG. 60 years of neuroendocrinology: the structure of the neuroendocrine hypothalamus: the neuroanatomical legacy of Geoffrey Harris. *J Endocrinol.* 2015;226(2):T25–39. Epub 2015/05/23.
120. Matsumoto H, Noguchi J, Takatsu Y, Horikoshi Y, Kumano S, Ohtaki T, et al. Stimulation effect of galanin-like peptide (GALP) on luteinizing hormone-releasing hormone-mediated luteinizing hormone (LH) secretion in male rats. *Endocrinology.* 2001;142(8):3693–6. Epub 2001/07/19.
121. Merchenthaler I, Lennard DE. The hypophysiotropic neurotensin-immunoreactive neuronal system of the rat brain. *Endocrinology.* 1991;129(6):2875–80. Epub 1991/12/01.
122. Merchenthaler I, Lopez FJ, Lennard DE, Negro-Vilar A. Sexual differences in the distribution of neurons coexpressing galanin and luteinizing hormone-releasing hormone in the rat brain. *Endocrinology.* 1991;129(4):1977–86. Epub 1991/10/01.
123. Niimi M, Takahara J, Sato M, Kawanishi K. Sites of origin of growth hormone-releasing factor-containing neurons projecting to the stalk-median eminence of the rat. *Peptides.* 1989;10(3):605–8. Epub 1989/05/01.
124. Niimi M, Takahara J, Sato M, Kawanishi K. Neurotensin and growth hormone-releasing factor-containing neurons projecting to the median eminence of the rat: a combined retrograde tracing and immunohistochemical study. *Neurosci Lett.* 1991;133(2):183–6. Epub 1991/12/09.
125. Niimi M, Takahara J, Sato M, Kawanishi K. Identification of dopamine and growth hormone-releasing factor-containing neurons projecting to the median eminence of the rat by combined retrograde tracing and immunohistochemistry. *Neuroendocrinology.* 1992;55(1):92–6. Epub 1992/01/01.
126. Takatsu Y, Matsumoto H, Ohtaki T, Kumano S, Kitada C, Onda H, et al. Distribution of galanin-like peptide in the rat brain. *Endocrinology.* 2001;142(4):1626–34. Epub 2001/03/17.
127. Dalcik H, Phelps CJ. Median eminence-afferent vasoactive intestinal peptide (VIP) neurons in the hypothalamus: localization by simultaneous tract tracing and immunocytochemistry. *Peptides.* 1993;14(5):1059–66. Epub 1993/09/01.
128. Ishikawa K, Taniguchi Y, Kurosumi K, Suzuki M, Shinoda M. Immunohistochemical identification of somatostatin-containing neurons projecting to the median eminence of the rat. *Endocrinology.* 1987;121(1):94–7. Epub 1987/07/01.
129. Kawano H, Daikoku S. Somatostatin-containing neuron systems in the rat hypothalamus: retrograde tracing and immunohistochemical studies. *J Comp Neurol.* 1988;271(2):293–9. Epub 1988/05/08.
130. Kawano H, Daikoku S, Shibasaki T. CRF-containing neuron systems in the rat hypothalamus: retrograde tracing and immunohistochemical studies. *J Comp Neurol.* 1988;272(2):260–8. Epub 1988/06/08.
131. Kawano H, Tsuruo Y, Bando H, Daikoku S. Hypophysiotrophic TRH-producing neurons

- identified by combining immunohistochemistry for pro-TRH and retrograde tracing. *J Comp Neurol.* 1991;307(4):531–8. Epub 1991/05/22.
132. Merchenthaler I. Enkephalin-immunoreactive neurons in the parvocellular subdivisions of the paraventricular nucleus project to the external zone of the median eminence. *J Comp Neurol.* 1992;326(1):112–20. Epub 1992/12/01.
 133. Merchenthaler I, Meeker M, Petrusz P, Kizer JS. Identification and immunocytochemical localization of a new thyrotropin-releasing hormone precursor in rat brain. *Endocrinology.* 1989;124(4):1888–97. Epub 1989/04/01.
 134. Merchenthaler I, Setalo G, Csontos C, Petrusz P, Flerko B, Negro-Vilar A. Combined retrograde tracing and immunocytochemical identification of luteinizing hormone-releasing hormone- and somatostatin-containing neurons projecting to the median eminence of the rat. *Endocrinology.* 1989;125(6):2812–21. Epub 1989/12/01.
 135. Niimi M, Takahara J, Hashimoto K, Kawanishi K. Immunohistochemical identification of corticotropin releasing factor-containing neurons projecting to the stalk-median eminence of the rat. *Peptides.* 1988;9(3):589–93. Epub 1988/05/01.
 136. De Mey J, Dierickx K, Vandesande F. Identification of neurophysin producing cells. III. Immunohistochemical demonstration of neurophysin I-producing neurons in the bovine infundibular nucleus. *Cell Tissue Res.* 1975;161(2):219–24. Epub 1975/08/18.
 137. De Mey J, Dierickx K, Vandesande F. Immunohistochemical demonstration of neurophysin I - and neurophysin II - containing nerve fibres in the external region of the bovine median eminence. *Cell Tissue Res.* 1975;157(4):517–9. Epub 1975/01/01.
 138. Vandesande F, Dierickx K. Identification of the vasopressin producing and of the oxytocin producing neurons in the hypothalamic magnocellular neurosecretory system of the rat. *Cell Tissue Res.* 1975;164(2):153–62. Epub 1975/12/02.
 139. Vandesande F, Dierickx K, De Mey J. Identification of separate vasopressin-neurophysin II and oxytocin-neurophysin I containing nerve fibres in the external region of the bovine median eminence. *Cell Tissue Res.* 1975;158(4):509–16. Epub 1975/05/20.
 140. Vandesande F, Dierickx K, DeMey J. Identification of the vasopressin-neurophysin producing neurons of the rat suprachiasmatic nuclei. *Cell Tissue Res.* 1975;156(3):377–80. Epub 1975/01/01.
 141. Vandesande F, Dierickx K, DeMey J. Identification of the vasopressin-neurophysin II and the oxytocin-neurophysin I producing neurons in the bovine hypothalamus. *Cell Tissue Res.* 1975;156(2):189–200. Epub 1975/01/01.
 142. Plotsky PM, Bruhn TO, Otto S. Central modulation of immunoreactive arginine vasopressin and oxytocin secretion into the hypophyseal-portal circulation by corticotropin-releasing factor. *Endocrinology.* 1985;116(4):1669–71. Epub 1985/04/01.
 143. Plotsky PM, Bruhn TO, Vale W. Hypophysiotropic regulation of adrenocorticotropin secretion in response to insulin-induced hypoglycemia. *Endocrinology.* 1985;117(1):323–9. Epub 1985/07/01.
 144. Plotsky PM, Bruhn TO, Vale W. Evidence for multi-factor regulation of the adrenocorticotropin secretory response to hemodynamic stimuli. *Endocrinology.* 1985;116(2):633–9. Epub 1985/02/01.
 145. Raff H. Interactions between neurohypophysial hormones and the ACTH-adrenocortical axis. *Ann N Y Acad Sci.* 1993;689:411–25. Epub 1993/07/22.
 146. Chowdrey HS, Lightman SL. Role of central amino acids and peptide-mediated pathways in neurohypophysial hormone release. *Ann N Y Acad Sci.* 1993;689:183–93. Epub 1993/07/22.
 147. Larsen PJ, Mikkelsen JD, Chowdrey HS, Jessop DS, Lightman SL. Osmotic regulation of neuro-peptide Y synthesis in magnocellular neurons of the hypothalamo-neurohypophysial system. *Ann N Y Acad Sci.* 1993;689:619–22. Epub 1993/07/22.
 148. Mikkelsen JD, Schmidt P, Sheikh SP, Larsen PJ. Non-vasopressinergic, non-oxytocinergic neuropeptides in the rat hypothalamo-neurohypophyseal tract: experimental immunohistochemical studies. *Prog Brain Res.* 1992;91:367–71. Epub 1992/01/01.
 149. Morris JF, Pow DV. New anatomical insights into the inputs and outputs from hypothalamic magnocellular neurons. *Annals of the New York Academy of Sciences.* 1993;689:16–33. Epub 1993/07/22.
 150. Nakamura S, Naruse M, Naruse K, Shioda S, Nakai Y, Uemura H. Colocalization of immunoreactive endothelin-1 and neurohypophysial hormones in the axons of the neural lobe of the rat pituitary. *Endocrinology.* 1993;132(2):530–3. Epub 1993/02/01.
 151. Stopa EG, LeBlanc VK, Hill DH, Anthony EL. A general overview of the anatomy of the neurohypophysis. *Ann N Y Acad Sci.* 1993;689:6–15. Epub 1993/07/22.
 152. Iliff JJ, Wang M, Liao Y, Plogg BA, Peng W, Gundersen GA, et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med.* 2012;4(147):147ra11. Epub 2012/08/17.
 153. Rennels ML, Gregory TF, Blaumanis OR, Fujimoto K, Grady PA. Evidence for a 'paravascular' fluid circulation in the mammalian central nervous system, provided by the rapid distribution of tracer protein throughout the brain from the subarachnoid space. *Brain Res.* 1985;326(1):47–63. Epub 1985/02/04.
 154. Tannenbaum GS, Martin JB. Evidence for an endogenous ultradian rhythm governing growth hormone secretion in the rat. *Endocrinology.* 1976;98(3):562–70. Epub 1976/03/01.

155. Tannenbaum GS, Martin JB, Colle E. Ultradian growth hormone rhythm in the rat: effects of feeding, hyperglycemia, and insulin-induced hypoglycemia. *Endocrinology*. 1976;99(3):720–7. Epub 1976/09/01.
156. Smith RG, Van der Ploeg LH, Howard AD, Feighner SD, Cheng K, Hickey GJ, et al. Peptidomimetic regulation of growth hormone secretion. *Endocr Rev*. 1997;18(5):621–45. Epub 1997/10/23 22:24.
157. Grouselle D, Chaillou E, Caraty A, Bluet-Pajot MT, Zizzari P, Tillet Y, et al. Pulsatile cerebrospinal fluid and plasma ghrelin in relation to growth hormone secretion and food intake in the sheep. *J Neuroendocrinol*. 2008;20(10):1138–46. Epub 2008/08/05.
158. Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*. 1999;402(6762):656–60. Epub 1999/12/22.
159. Muccioli G, Tschop M, Papotti M, Deghenghi R, Heiman M, Ghigo E. Neuroendocrine and peripheral activities of ghrelin: implications in metabolism and obesity. *Eur J Pharmacol*. 2002;440(2–3):235–54. Epub 2002/05/15.
160. Muller TD, Nogueiras R, Andermann ML, Andrews ZB, Anker SD, Argente J, et al. Ghrelin. *Mol Metab*. 2015;4(6):437–60. Epub 2015/06/05.
161. Tschop M, Flora DB, Mayer JP, Heiman ML. Hypophysectomy prevents ghrelin-induced adiposity and increases gastric ghrelin secretion in rats. *Obes Res*. 2002;10(10):991–9. Epub 2002/10/12.
162. Ukkola O, Ravussin E, Jacobson P, Perusse L, Rankinen T, Tschop M, et al. Role of ghrelin polymorphisms in obesity based on three different studies. *Obes Res*. 2002;10(8):782–91. Epub 2002/08/16.
163. Plotsky PM, Kjaer A, Sutton SW, Sawchenko PE, Vale W. Central activin administration modulates corticotropin-releasing hormone and adrenocorticotropin secretion. *Endocrinology*. 1991;128(5):2520–5. Epub 1991/05/01.
164. Sarkar S, Fekete C, Legradi G, Lechan RM. Glucagon like peptide-1 (7–36) amide (GLP-1) nerve terminals densely innervate corticotropin-releasing hormone neurons in the hypothalamic paraventricular nucleus. *Brain Res*. 2003;985(2):163–8. Epub 2003/09/12.
165. Sawchenko PE, Swanson LW. Central noradrenergic pathways for the integration of hypothalamic neuroendocrine and autonomic responses. *Science*. 1981;214(4521):685–7. Epub 1981/11/06.
166. Sawchenko PE, Swanson LW, Grzanna R, Howe PR, Bloom SR, Polak JM. Colocalization of neuropeptide Y immunoreactivity in brainstem catecholaminergic neurons that project to the paraventricular nucleus of the hypothalamus. *J Comp Neurol*. 1985;241(2):138–53. Epub 1985/11/08.
167. Sawchenko PE, Swanson LW, Rivier J, Vale WW. The distribution of growth-hormone-releasing factor (GRF) immunoreactivity in the central nervous system of the rat: an immunohistochemical study using antisera directed against rat hypothalamic GRF. *J Comp Neurol*. 1985;237(1):100–15. Epub 1985/07/01.
168. Besedovsky HO, del Rey A. Immune-neuroendocrine interactions: facts and hypotheses. *Endocr Rev*. 1996;17(1):64–102. Epub 1996/02/01.
169. Sawchenko PE, Li HY, Ericsson A. Circuits and mechanisms governing hypothalamic responses to stress: a tale of two paradigms. *Prog Brain Res*. 2000;122:61–78. Epub 2000/03/29.
170. Dimitrov EL, DeJoseph MR, Brownfield MS, Urban JH. Involvement of neuropeptide Y Y1 receptors in the regulation of neuroendocrine corticotropin-releasing hormone neuronal activity. *Endocrinology*. 2007;148(8):3666–73. Epub 2007/04/28.
171. Tuchelt H, Dekker K, Bahr V, Oelkers W. Dose-response relationship between plasma ACTH and serum cortisol in the insulin-hypoglycaemia test in 25 healthy subjects and 109 patients with pituitary disease. *Clin Endocrinol*. 2000;53(3):301–7. Epub 2000/09/06.
172. Levy BH, Tasker JG. Synaptic regulation of the hypothalamic-pituitary-adrenal axis and its modulation by glucocorticoids and stress. *Front Cell Neurosci*. 2012;6:24. Epub 2012/05/18.
173. Sanchez E, Vargas MA, Singru PS, Pascual I, Romero F, Fekete C, et al. Tanycyte pyroglutamate peptidase II contributes to regulation of the hypothalamic-pituitary-thyroid axis through glial-axonal associations in the median eminence. *Endocrinology*. 2009;150(5):2283–91. Epub 2009/01/31.
174. Blake NG, Eckland DJ, Foster OJ, Lightman SL. Inhibition of hypothalamic thyrotropin-releasing hormone messenger ribonucleic acid during food deprivation. *Endocrinology*. 1991;129(5):2714–8. Epub 1991/11/01.
175. Harris AR, Fang SL, Azizi F, Lipworth L, Vagenakis AG, Braverman LE. Effect of starvation on hypothalamic-pituitary-thyroid function in the rat. *Metab Clin Exp*. 1978;27(9):1074–83. Epub 1978/09/01.
176. Harris AR, Fang SL, Vagenakis AG, Braverman LE. Effect of starvation, nutrient replacement, and hypothyroidism on in vitro hepatic T4 to T3 conversion in the rat. *Metab Clin Exp*. 1978;27(11):1680–90. Epub 1978/11/01.
177. Legradi G, Emerson CH, Ahima RS, Flier JS, Lechan RM. Leptin prevents fasting-induced suppression of prothyrotropin-releasing hormone messenger ribonucleic acid in neurons of the hypothalamic paraventricular nucleus. *Endocrinology*. 1997;138(6):2569–76. Epub 1997/06/01.
178. Rondeel JM, Heide R, de Greef WJ, van Toor H, van Haasteren GA, Klootwijk W, et al. Effect of starvation and subsequent refeeding on thyroid function and release of hypothalamic thyrotropin-releasing hormone. *Neuroendocrinology*. 1992;56(3):348–53. Epub 1992/09/01.

179. Flier JS, Harris M, Hollenberg AN. Leptin, nutrition, and the thyroid: the why, the wherefore, and the wiring. *J Clin Invest*. 2000;105(7):859–61. Epub 2000/04/05.
180. Adler SM, Wartofsky L. The nonthyroidal illness syndrome. *Endocrinol Metab Clin N Am*. 2007;36(3):657–72. vi. Epub 2007/08/04.
181. Bartalena L. The dilemma of non-thyroidal illness syndrome: to treat or not to treat? *J Endocrinol Investig*. 2003;26(12):1162. Epub 2004/04/02.
182. DeGroot LJ. “Non-thyroidal illness syndrome” is functional central hypothyroidism, and if severe, hormone replacement is appropriate in light of present knowledge. *J Endocrinol Investig*. 2003;26(12):1163–70. Epub 2004/04/02.
183. Lee S, Farwell AP. Euthyroid sick syndrome. *Compr Physiol*. 2016;6(2):1071–80. Epub 2016/04/12.
184. Stathatos N, Wartofsky L. The euthyroid sick syndrome: is there a physiologic rationale for thyroid hormone treatment? *J Endocrinol Investig*. 2003;26(12):1174–9. Epub 2004/04/02.
185. Wartofsky L, Burman KD, Ringel MD. Trading one “dangerous dogma” for another? Thyroid hormone treatment of the “euthyroid sick syndrome”. *J Clin Endocrinol Metab*. 1999;84(5):1759–60. Epub 1999/05/14.
186. Fekete C, Kelly J, Mihaly E, Sarkar S, Rand WM, Legradi G, et al. Neuropeptide Y has a central inhibitory action on the hypothalamic-pituitary-thyroid axis. *Endocrinology*. 2001;142(6):2606–13. Epub 2001/05/18.
187. Fekete C, Legradi G, Mihaly E, Huang QH, Tatso JB, Rand WM, et al. Alpha-melanocyte-stimulating hormone is contained in nerve terminals innervating thyrotropin-releasing hormone-synthesizing neurons in the hypothalamic paraventricular nucleus and prevents fasting-induced suppression of prothyrotropin-releasing hormone gene expression. *J Neurosci*. 2000;20(4):1550–8. Epub 2000/02/09.
188. Fekete C, Mihaly E, Luo LG, Kelly J, Clausen JT, Mao Q, et al. Association of cocaine- and amphetamine-regulated transcript-immunoreactive elements with thyrotropin-releasing hormone-synthesizing neurons in the hypothalamic paraventricular nucleus and its role in the regulation of the hypothalamic-pituitary-thyroid axis during fasting. *J Neurosci*. 2000;20(24):9224–34. Epub 2000/01/11.
189. Fekete C, Sarkar S, Rand WM, Harney JW, Emerson CH, Bianco AC, et al. Neuropeptide Y1 and Y5 receptors mediate the effects of neuropeptide Y on the hypothalamic-pituitary-thyroid axis. *Endocrinology*. 2002;143(12):4513–9. Epub 2002/11/26.
190. Fekete C, Sarkar S, Rand WM, Harney JW, Emerson CH, Bianco AC, et al. Agouti-related protein (AGRP) has a central inhibitory action on the hypothalamic-pituitary-thyroid (HPT) axis; comparisons between the effect of AGRP and neuropeptide Y on energy homeostasis and the HPT axis. *Endocrinology*. 2002;143(10):3846–53. Epub 2002/09/20.
191. Rugarli EI, Ballabio A. Kallmann syndrome. From genetics to neurobiology. *JAMA*. 1993;270(22):2713–6. Epub 1993/12/08.
192. Ollmann MM, Wilson BD, Yang YK, Kerns JA, Chen Y, Gantz I, et al. Antagonism of central melanocortin receptors in vitro and in vivo by agouti-related protein. *Science*. 1997;278(5335):135–8. Epub 1997/10/06.
193. Ahima RS, Saper CB, Flier JS, Elmquist JK. Leptin regulation of neuroendocrine systems. *Front Neuroendocrinol*. 2000;21(3):263–307. Epub 2000/07/07.
194. Elias CF, Kelly JF, Lee CE, Ahima RS, Drucker DJ, Saper CB, et al. Chemical characterization of leptin-activated neurons in the rat brain. *J Comp Neurol*. 2000;423(2):261–81. Epub 2000/06/27.
195. Clarke IJ, Cummins JT. The temporal relationship between gonadotropin releasing hormone (GnRH) and luteinizing hormone (LH) secretion in ovariectomized ewes. *Endocrinology*. 1982;111(5):1737–9. Epub 1982/11/01.
196. Ekholm C, Clark MR, Magnusson C, Isaksson O, LeMaire WJ. Ovulation induced by a gonadotropin releasing hormone analog in hypophysectomized rats involves prostaglandins. *Endocrinology*. 1982;110(1):288–90. Epub 1982/01/01.
197. Moenter SM, Caraty A, Locatelli A, Karsch FJ. Pattern of gonadotropin-releasing hormone (GnRH) secretion leading up to ovulation in the ewe: existence of a preovulatory GnRH surge. *Endocrinology*. 1991;129(3):1175–82. Epub 1991/09/01.
198. Johnson ES, Gendrich RL, White WF. Delay of puberty and inhibition of reproductive processes in the rat by a gonadotropin-releasing hormone agonist analog. *Fertil Steril*. 1976;27(7):853–60. Epub 1976/07/01.
199. Muir AI, Chamberlain L, Elshourbagy NA, Michalovich D, Moore DJ, Calamari A, et al. AXOR12, a novel human G protein-coupled receptor, activated by the peptide KiSS-1. *J Biol Chem*. 2001;276(31):28969–75. Epub 2001/06/02.
200. Seminara SB, Messenger S, Chatzidaki EE, Thresher RR, Acierno JS Jr, Shagoury JK, et al. The GPR54 gene as a regulator of puberty. *N Engl J Med*. 2003;349(17):1614–27. Epub 2003/10/24.
201. Kotani M, Dethoux M, Vandenbogaerde A, Communi D, Vanderwinden JM, Le Poul E, et al. The metastasis suppressor gene KiSS-1 encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor GPR54. *J Biol Chem*. 2001;276(37):34631–6. Epub 2001/07/18.
202. Ohtaki T, Shintani Y, Honda S, Matsumoto H, Hori A, Kanehashi K, et al. Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-coupled receptor. *Nature*. 2001;411(6837):613–7. Epub 2001/06/01.

203. Messager S, Chatzidaki EE, Ma D, Hendrick AG, Zahn D, Dixon J, et al. Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *Proc Natl Acad Sci U S A*. 2005;102(5):1761–6. Epub 2005/01/25.
204. Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, Crowley WF, et al. A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology*. 2004;145(9):4073–7. Epub 2004/06/26.
205. Matsui H, Asami T. Effects and therapeutic potentials of kisspeptin analogs: regulation of the hypothalamic-pituitary-gonadal axis. *Neuroendocrinology*. 2014;99(1):49–60. Epub 2013/12/21.
206. Matsui H, Takatsu Y, Kumano S, Matsumoto H, Ohtaki T. Peripheral administration of metastatin induces marked gonadotropin release and ovulation in the rat. *Biochem Biophys Res Commun*. 2004;320(2):383–8. Epub 2004/06/29.
207. Bodo C, Kudwa AE, Rissman EF. Both estrogen receptor- α and - β are required for sexual differentiation of the anteroventral periventricular area in mice. *Endocrinology*. 2006;147(1):415–20. Epub 2005/10/22.
208. Decourt C, Tillet Y, Caraty A, Franceschini I, Briant C. Kisspeptin immunoreactive neurons in the equine hypothalamus interactions with GnRH neuronal system. *J Chem Neuroanat*. 2008;36(3–4):131–7. Epub 2008/09/02.
209. Franceschini I, Lomet D, Cateau M, Delsol G, Tillet Y, Caraty A. Kisspeptin immunoreactive cells of the ovine preoptic area and arcuate nucleus co-express estrogen receptor α . *Neurosci Lett*. 2006;401(3):225–30. Epub 2006/04/20.
210. Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA. Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology*. 2005;146(9):3686–92. Epub 2005/05/28.
211. Dungan HM, Clifton DK, Steiner RA. Minireview: kisspeptin neurons as central processors in the regulation of gonadotropin-releasing hormone secretion. *Endocrinology*. 2006;147(3):1154–8. Epub 2005/12/24.
212. Ramaswamy S, Guerriero KA, Gibbs RB, Plant TM. Structural interactions between kisspeptin and GnRH neurons in the mediobasal hypothalamus of the male rhesus monkey (*Macaca mulatta*) as revealed by double immunofluorescence and confocal microscopy. *Endocrinology*. 2008;149(9):4387–95. Epub 2008/05/31.
213. Luo E, Stephens SB, Chaing S, Munaganuru N, Kauffman AS, Breen KM. Corticosterone blocks ovarian cyclicity and the LH surge via decreased Kisspeptin neuron activation in female mice. *Endocrinology*. 2016;157(3):1187–99. Epub 2015/12/25.
214. Stephens SB, Tolson KP, Rouse ML Jr, Poling MC, Hashimoto-Partyka MK, Mellon PL, et al. Absent progesterone signaling in kisspeptin neurons disrupts the LH surge and impairs fertility in female mice. *Endocrinology*. 2015;156(9):3091–7. Epub 2015/06/16.
215. Homma T, Sakakibara M, Yamada S, Kinoshita M, Iwata K, Tomikawa J, et al. Significance of neonatal testicular sex steroids to defeminize anteroventral periventricular kisspeptin neurons and the GnRH/LH surge system in male rats. *Biol Reprod*. 2009;81(6):1216–25. Epub 2009/08/18.
216. Kauffman AS, Gottsch ML, Roa J, Byquist AC, Crown A, Clifton DK, et al. Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinology*. 2007;148(4):1774–83. Epub 2007/01/06.
217. Tomikawa J, Homma T, Tajima S, Shibata T, Inamoto Y, Takase K, et al. Molecular characterization and estrogen regulation of hypothalamic KISS1 gene in the pig. *Biol Reprod*. 2010;82(2):313–9. Epub 2009/10/16.
218. Billings HJ, Connors JM, Altman SN, Hileman SM, Holaskova I, Lehman MN, et al. Neurokinin B acts via the neurokinin-3 receptor in the retrochiasmatic area to stimulate luteinizing hormone secretion in sheep. *Endocrinology*. 2010;151(8):3836–46. Epub 2010/06/04.
219. Lehman MN, Coolen LM, Goodman RL. Minireview: kisspeptin/neurokinin B/dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology*. 2010;151(8):3479–89. Epub 2010/05/27.
220. Cheng G, Coolen LM, Padmanabhan V, Goodman RL, Lehman MN. The kisspeptin/neurokinin B/dynorphin (KNDy) cell population of the arcuate nucleus: sex differences and effects of prenatal testosterone in sheep. *Endocrinology*. 2010;151(1):301–11. Epub 2009/11/03.
221. Roa J, Navarro VM, Tena-Sempere M. Kisspeptins in reproductive biology: consensus knowledge and recent developments. *Biol Reprod*. 2011;85(4):650–60. Epub 2011/06/17.
222. Quennell JH, Howell CS, Roa J, Augustine RA, Grattan DR, Anderson GM. Leptin deficiency and diet-induced obesity reduce hypothalamic kisspeptin expression in mice. *Endocrinology*. 2011;152(4):1541–50. Epub 2011/02/18.
223. Freeman ME, Kanyicska B, Lerant A, Nagy G. Prolactin: structure, function, and regulation of secretion. *Physiol Rev*. 2000;80(4):1523–631. Epub 2000/10/04.
224. Li C, Chen P, Smith MS. Neural populations in the rat forebrain and brainstem activated by the suckling stimulus as demonstrated by cFos expression. *Neuroscience*. 1999;94(1):117–29. Epub 1999/12/29.
225. Grattan DR. Behavioural significance of prolactin signalling in the central nervous system during pregnancy and lactation. *Reproduction*. 2002;123(4):497–506. Epub 2002/03/27.

226. Crowley WR. Neuroendocrine regulation of lactation and milk production. *Compr Physiol*. 2015;5(1):255–91. Epub 2015/01/16.
227. John MR, Arai M, Rubin DA, Jonsson KB, Juppner H. Identification and characterization of the murine and human gene encoding the tuberoinfundibular peptide of 39 residues. *Endocrinology*. 2002;143(3):1047–57. Epub 2002/02/28.
228. Weaver RE, Mobarec JC, Wigglesworth MJ, Reynolds CA, Donnelly D. High affinity binding of the peptide agonist TIP-39 to the parathyroid hormone 2 (PTH2) receptor requires the hydroxyl group of Tyr-318 on transmembrane helix 5. *Biochem Pharmacol*. 2017;127:71–81. Epub 2016/12/26.
229. Ben-Jonathan N, Hnasko R. Dopamine as a prolactin (PRL) inhibitor. *Endocr Rev*. 2001;22(6):724–63. Epub 2001/12/12.
230. Romano N, Yip SH, Hodson DJ, Guillou A, Parnaudeau S, Kirk S, et al. Plasticity of hypothalamic dopamine neurons during lactation results in dissociation of electrical activity and release. *J Neurosci*. 2013;33(10):4424–33. Epub 2013/03/08.
231. Pi XJ, Grattan DR. Increased prolactin receptor immunoreactivity in the hypothalamus of lactating rats. *J Neuroendocrinol*. 1999;11(9):693–705. Epub 1999/08/14.
232. Walsh RJ, Slaby FJ, Posner BI. A receptor-mediated mechanism for the transport of prolactin from blood to cerebrospinal fluid. *Endocrinology*. 1987;120(5):1846–50. Epub 1987/05/01.
233. Smith MS, True C, Grove KL. The neuroendocrine basis of lactation-induced suppression of GnRH: role of kisspeptin and leptin. *Brain Res*. 2010;1364:139–52. Epub 2010/08/24.
234. Yamada S, Uenoyama Y, Kinoshita M, Iwata K, Takase K, Matsui H, et al. Inhibition of metastin (kisspeptin-54)-GPR54 signaling in the arcuate nucleus-median eminence region during lactation in rats. *Endocrinology*. 2007;148(5):2226–32. Epub 2007/02/10.
235. Smith MS. Lactation alters neuropeptide-Y and proopiomelanocortin gene expression in the arcuate nucleus of the rat. *Endocrinology*. 1993;133(3):1258–65. Epub 1993/09/01.
236. Smith MS, Grove KL. Integration of the regulation of reproductive function and energy balance: lactation as a model. *Front Neuroendocrinol*. 2002;23(3):225–56. Epub 2002/07/20.
237. Li C, Chen P, Smith MS. The acute suckling stimulus induces expression of neuropeptide Y (NPY) in cells in the dorsomedial hypothalamus and increases NPY expression in the arcuate nucleus. *Endocrinology*. 1998;139(4):1645–52. Epub 1998/04/07.
238. Chen P, Li C, Haskell-Luevano C, Cone RD, Smith MS. Altered expression of agouti-related protein and its colocalization with neuropeptide Y in the arcuate nucleus of the hypothalamus during lactation. *Endocrinology*. 1999;140(6):2645–50. Epub 1999/05/26.
239. Haskell-Luevano C, Chen P, Li C, Chang K, Smith MS, Cameron JL, et al. Characterization of the neuroanatomical distribution of agouti-related protein immunoreactivity in the rhesus monkey and the rat. *Endocrinology*. 1999;140(3):1408–15. Epub 1999/03/06.
240. Li C, Chen P, Smith MS. Morphological evidence for direct interaction between arcuate nucleus neuropeptide Y (NPY) neurons and gonadotropin-releasing hormone neurons and the possible involvement of NPY Y1 receptors. *Endocrinology*. 1999;140(11):5382–90. Epub 1999/10/28.
241. Li C, Chen P, Smith MS. Neuropeptide Y and tuberoinfundibular dopamine activities are altered during lactation: role of prolactin. *Endocrinology*. 1999;140(1):118–23. Epub 1999/01/14.
242. Li C, Chen P, Smith MS. Identification of neuronal input to the arcuate nucleus (ARH) activated during lactation: implications in the activation of neuropeptide Y neurons. *Brain Res*. 1999;824(2):267–76. Epub 1999/04/10.
243. Hetherington AW, Ranson SW. Hypothalamic lesions and adiposity in the rat. *Anat Rec*. 1940;78:149–72.
244. Reeves AG, Plum F. Hyperphagia, rage, and dementia accompanying a ventromedial hypothalamic neoplasm. *Arch Neurol*. 1969;20(6):616–24. Epub 1969/06/01.
245. Malenka RC, Nestler EJ, Hyman SE. Neural and neuroendocrine control of the internal milieu. In: Sydor A, Brown RY, editors. *Molecular neuropharmacology; a foundation for clinical neuroscience*. 2nd ed. New York: McGraw-Hill Medical; 2009. p. 263.
246. Theologides A. Anorexia-producing intermediary metabolites. *Am J Clin Nutr*. 1976;29(5):552–8. Epub 1976/05/01.
247. Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature*. 1994;372(6505):425–32. Epub 1994/12/01.
248. Barsh GS, Schwartz MW. Genetic approaches to studying energy balance: perception and integration. *Nat Rev Genet*. 2002;3(8):589–600. Epub 2002/08/03.
249. Donato J Jr, Cravo RM, Frazao R, Gautron L, Scott MM, Lachey J, et al. Leptin's effect on puberty in mice is relayed by the ventral premammillary nucleus and does not require signaling in Kiss1 neurons. *J Clin Invest*. 2011;121(1):355–68. Epub 2010/12/25.
250. Gautron L, Elmquist JK. Sixteen years and counting: an update on leptin in energy balance. *J Clin Invest*. 2011;121(6):2087–93. Epub 2011/06/03.
251. Kalra SP, Dube MG, Pu S, Xu B, Horvath TL, Kalra PS. Interacting appetite-regulating pathways in the hypothalamic regulation of body weight. *Endocr Rev*. 1999;20(1):68–100. Epub 1999/02/27.
252. Schwartz MW, Woods SC, Porte D Jr, Seeley RJ, Baskin DG. Central nervous system control of

- food intake. *Nature*. 2000;404(6778):661–71. Epub 2000/04/15.
253. Scott MM, Williams KW, Rossi J, Lee CE, Elmquist JK. Leptin receptor expression in hindbrain Glp-1 neurons regulates food intake and energy balance in mice. *J Clin Invest*. 2011;121(6):2413–21. Epub 2011/05/25.
 254. Cowley MA, Smart JL, Rubinstein M, Cerdan MG, Diano S, Horvath TL, et al. Leptin activates anorexigenic POMC neurons through a neural network in the arcuate nucleus. *Nature*. 2001;411(6836):480–4. Epub 2001/05/25.
 255. Nijenhuis WA, Oosterom J, Adan RA. AgRP(83–132) acts as an inverse agonist on the human-melanocortin-4 receptor. *Mol Endocrinol*. 2001;15(1):164–71. Epub 2001/01/06.
 256. Erickson JC, Clegg KE, Palmiter RD. Sensitivity to leptin and susceptibility to seizures of mice lacking neuropeptide Y. *Nature*. 1996;381(6581):415–21. Epub 1996/05/30.
 257. Farooqi IS, Yeo GS, Keogh JM, Aminian S, Jebb SA, Butler G, et al. Dominant and recessive inheritance of morbid obesity associated with melanocortin 4 receptor deficiency. *J Clin Invest*. 2000;106(2):271–9. Epub 2000/07/21.
 258. Huszar D, Lynch CA, Fairchild-Huntress V, Dunmore JH, Fang Q, Berkemeier LR, et al. Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*. 1997;88(1):131–41. Epub 1997/01/10.
 259. Jackson RS, Creemers JW, Ohagi S, Raffin-Sanson ML, Sanders L, Montague CT, et al. Obesity and impaired prohormone processing associated with mutations in the human prohormone convertase 1 gene. *Nat Genet*. 1997;16(3):303–6. Epub 1997/07/01.
 260. Krude H, Biebermann H, Luck W, Horn R, Brabant G, Gruters A. Severe early-onset obesity, adrenal insufficiency and red hair pigmentation caused by POMC mutations in humans. *Nat Genet*. 1998;19(2):155–7. Epub 1998/06/10.
 261. Marks DL, Ling N, Cone RD. Role of the central melanocortin system in cachexia. *Cancer Res*. 2001;61(4):1432–8. Epub 2001/03/14.
 262. Wisse BE, Frayo RS, Schwartz MW, Cummings DE. Reversal of cancer anorexia by blockade of central melanocortin receptors in rats. *Endocrinology*. 2001;142(8):3292–301. Epub 2001/07/19.
 263. Elmquist JK. Hypothalamic pathways underlying the endocrine, autonomic, and behavioral effects of leptin. *Physiol Behav*. 2001;74(4–5):703–8. Epub 2002/01/16.
 264. Saper CB. Organization of cerebral cortical afferent systems in the rat. II. Hypothalamocortical projections. *J Comp Neurol*. 1985;237(1):21–46. Epub 1985/07/01.
 265. Ahima RS, Flier JS. Leptin. *Annu Rev Physiol*. 2000;62:413–37. Epub 2000/06/09.
 266. Murphy KG, Dhillon WS, Bloom SR. Gut peptides in the regulation of food intake and energy homeostasis. *Endocr Rev*. 2006;27(7):719–27. Epub 2006/11/02.
 267. Schwartz MW, Sipols AJ, Marks JL, Sanacora G, White JD, Scheurink A, et al. Inhibition of hypothalamic neuropeptide Y gene expression by insulin. *Endocrinology*. 1992;130(6):3608–16. Epub 1992/06/01.
 268. Woods SC, Seeley RJ, Porte D Jr, Schwartz MW. Signals that regulate food intake and energy homeostasis. *Science*. 1998;280(5368):1378–83. Epub 1998/06/20.
 269. Fort P, Salvetti D, Hanriot L, Jegu S, Shimizu H, Hashimoto K, et al. The satiety molecule nesfatin-1 is co-expressed with melanin concentrating hormone in tuberal hypothalamic neurons of the rat. *Neuroscience*. 2008;155(1):174–81. Epub 2008/06/25.
 270. Oh IS, Shimizu H, Satoh T, Okada S, Adachi S, Inoue K, et al. Identification of nesfatin-1 as a satiety molecule in the hypothalamus. *Nature*. 2006;443(7112):709–12. Epub 2006/10/13.
 271. Price CJ, Samson WK, Ferguson AV. Nesfatin-1 inhibits NPY neurons in the arcuate nucleus. *Brain Res*. 2008;1230:99–106. Epub 2008/07/16.
 272. Stengel A, Goebel M, Yakubov I, Wang L, Witcher D, Coskun T, et al. Identification and characterization of nesfatin-1 immunoreactivity in endocrine cell types of the rat gastric oxyntic mucosa. *Endocrinology*. 2009;150(1):232–8. Epub 2008/09/27.
 273. Zhang JV, Ren PG, Avsian-Kretschmer O, Luo CW, Rauch R, Klein C, et al. Obestatin, a peptide encoded by the ghrelin gene, opposes ghrelin's effects on food intake. *Science*. 2005;310(5750):996–9. Epub 2005/11/15.
 274. Bauer PV, Hamr SC, Duca FA. Regulation of energy balance by a gut-brain axis and involvement of the gut microbiota. *Cell Mol Life Sci*. 2016;73(4):737–55. Epub 2015/11/07.
 275. Gavrieli A, Mantzoros CS. Novel molecules regulating energy homeostasis: physiology and regulation by macronutrient intake and weight loss. *Endocrinol Metab*. 2016;31(3):361–72. Epub 2016/07/30.
 276. Abraham G, Falcou R, Rozen R, Mandenoff A, Autissier N, Apfelbaum M. The effects of a constant T3 level and thermoneutrality in diet-induced hyperphagia. *Hormone Metab Res*. 1987;19(3):96–100. Epub 1987/03/01.
 277. Coppola A, Liu ZW, Andrews ZB, Paradis E, Roy MC, Friedman JM, et al. A central thermogenic-like mechanism in feeding regulation: an interplay between arcuate nucleus T3 and UCP2. *Cell Metab*. 2007;5(1):21–33. Epub 2006/12/26.
 278. Ishii S, Kamegai J, Tamura H, Shimizu T, Sugihara H, Oikawa S. Hypothalamic neuropeptide Y/Y1 receptor pathway activated by a reduction in circulating leptin, but not by an increase in circulating ghrelin, contributes to hyperphagia associated with triiodothyronine-induced thyrotoxicosis. *Neuroendocrinology*. 2003;78(6):321–30. Epub 2003/12/23.

279. Kong WM, Martin NM, Smith KL, Gardiner JV, Connoley IP, Stephens DA, et al. Triiodothyronine stimulates food intake via the hypothalamic ventromedial nucleus independent of changes in energy expenditure. *Endocrinology*. 2004;145(11):5252–8. Epub 2004/08/07.
280. Luo L, MacLean DB. Effects of thyroid hormone on food intake, hypothalamic Na/K ATPase activity and ATP content. *Brain Res*. 2003;973(2):233–9. Epub 2003/05/10.
281. Syed MA, Thompson MP, Pachucki J, Burmeister LA. The effect of thyroid hormone on size of fat depots accounts for most of the changes in leptin mRNA and serum levels in the rat. *Thyroid Off J Am Thy Ass*. 1999;9(5):503–12. Epub 1999/06/12.
282. Broedel O, Eravci M, Fuxius S, Smolarz T, Jeitner A, Grau H, et al. Effects of hyper- and hypothyroidism on thyroid hormone concentrations in regions of the rat brain. *Am J Physiol Endocrinol Metab*. 2003;285(3):E470–80. Epub 2003/05/09.
283. Campos-Barros A, Musa A, Flechner A, Hassenius C, Gaio U, Meinhold H, et al. Evidence for circadian variations of thyroid hormone concentrations and type II 5'-iodothyronine deiodinase activity in the rat central nervous system. *J Neurochem*. 1997;68(2):795–803. Epub 1997/02/01.
284. Kadowaki T, Yamauchi T, Kubota N. The physiological and pathophysiological role of adiponectin and adiponectin receptors in the peripheral tissues and CNS. *FEBS Lett*. 2008;582(1):74–80. Epub 2007/12/07.
285. Uchida-Kitajima S, Yamauchi T, Takashina Y, Okada-Iwabu M, Iwabu M, Ueki K, et al. 5-Hydroxytryptamine 2A receptor signaling cascade modulates adiponectin and plasminogen activator inhibitor 1 expression in adipose tissue. *FEBS Lett*. 2008;582(20):3037–44. Epub 2008/08/05.
286. Kubota N, Yano W, Kubota T, Yamauchi T, Itoh S, Kumagai H, et al. Adiponectin stimulates AMP-activated protein kinase in the hypothalamus and increases food intake. *Cell Metab*. 2007;6(1):55–68. Epub 2007/07/10.
287. Lechan RM, Fekete C. Neuroendocrine and metabolic adaptations in the central nervous system to weight loss that facilitate weight regain. In: Freemark M, editor. *Pediatric obesity: etiology, pathogenesis and treatment*. New York, USA: Humana Press/Springer, 2010.
288. Williams KW, Elmquist JK. From neuroanatomy to behavior: central integration of peripheral signals regulating feeding behavior. *Nat Neurosci*. 2012;15(10):1350–5. Epub 2012/09/26.
289. Freund TF, Katona I, Piomelli D. Role of endogenous cannabinoids in synaptic signaling. *Physiol Rev*. 2003;83(3):1017–66. Epub 2003/07/05.
290. Piomelli D. The molecular logic of endocannabinoid signalling. *Nat Rev Neurosci*. 2003;4(11):873–84. Epub 2003/11/05.
291. Wittmann G, Deli L, Kallo I, Hrabovszky E, Watanabe M, Liposits Z, et al. Distribution of type 1 cannabinoid receptor (CB1)-immunoreactive axons in the mouse hypothalamus. *J Comp Neurol*. 2007;503(2):270–9. Epub 2007/05/12.
292. Jo YH, Chen YJ, Chua SC Jr, Talmage DA, Role LW. Integration of endocannabinoid and leptin signaling in an appetite-related neural circuit. *Neuron*. 2005;48(6):1055–66. Epub 2005/12/21.
293. Kola B, Farkas I, Christ-Crain M, Wittmann G, Lolli F, Amin F, et al. The orexigenic effect of ghrelin is mediated through central activation of the endogenous cannabinoid system. *PLoS One*. 2008;3(3):e1797. Epub 2008/03/13.
294. Vong L, Ye C, Yang Z, Choi B, Chua S Jr, Lowell BB. Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron*. 2011;71(1):142–54. Epub 2011/07/13.
295. Xu Y, Berglund ED, Sohn JW, Holland WL, Chuang JC, Fukuda M, et al. 5-HT2CRs expressed by proopiomelanocortin neurons regulate insulin sensitivity in liver. *Nat Neurosci*. 2010;13(12):1457–9. Epub 2010/11/03.
296. Fulton S, Pissios P, Manchon RP, Stiles L, Frank L, Pothos EN, et al. Leptin regulation of the mesoaccumbens dopamine pathway. *Neuron*. 2006;51(6):811–22. Epub 2006/09/20.
297. Geiger BM, Haburcak M, Avena NM, Moyer MC, Hoebel BG, Pothos EN. Deficits of mesolimbic dopamine neurotransmission in rat dietary obesity. *Neuroscience*. 2009;159(4):1193–9. Epub 2009/05/05.
298. Kelley AE, Berridge KC. The neuroscience of natural rewards: relevance to addictive drugs. *J Neurosci*. 2002;22(9):3306–11. Epub 2002/04/30.
299. Reynolds SM, Berridge KC. Positive and negative motivation in nucleus accumbens shell: bivalent rostrocaudal gradients for GABA-elicited eating, taste “liking”/“disliking” reactions, place preference/avoidance, and fear. *J Neurosci*. 2002;22(16):7308–20. Epub 2002/08/15.
300. Castrellon JJ, Seaman KL, Crawford JL, Young JS, Smith CT, Dang LC, et al. Individual differences in dopamine are associated with reward discounting in clinical groups but not in healthy adults. *J Neurosci*. 2019;39(2):321–32. Epub 2018/11/18.
301. Kramer V, Juri C, Riss PJ, Pruzzo R, Soza-Ried C, Flores J, et al. Pharmacokinetic evaluation of [(18)F]PR04.MZ for PET/CT imaging and quantification of dopamine transporters in the human brain. *Eur J Nucl Med Mol Imaging*. 2019; Epub 2019/12/04.
302. Lee MR, Shin JH, Deschaine S, Daurio AM, Stangl BL, Yan J, et al. A role for the CD38 rs3796863 polymorphism in alcohol and monetary reward: evidence from CD38 knockout mice and alcohol self-administration, [11C]-raclopride binding, and functional MRI in humans. *Am J Drug Alcohol Abuse*. 2019;1–13. Epub 2019/08/01.
303. Meyer GM, Spay C, Laurencin C, Ballanger B, Sescousse G, Boulinguez P. Functional imaging studies of impulse control disorders in Parkinson's

- disease need a stronger neurocognitive footing. *Neurosci Biobehav Rev.* 2019;98:164–76. Epub 2019/01/15.
304. Stark AJ, Smith CT, Lin YC, Petersen KJ, Trujillo P, van Wouwe NC, et al. Nigrostriatal and mesolimbic D2/3 receptor expression in Parkinson's disease patients with compulsive reward-driven Behaviors. *J Neurosci.* 2018;38(13):3230–9. Epub 2018/02/28.
 305. Tanabe J, Regner M, Sakai J, Martinez D, Gowin J. Neuroimaging reward, craving, learning, and cognitive control in substance use disorders: review and implications for treatment. *Br J Radiol.* 2019;92(1101):20180942. Epub 2019/03/12.
 306. Thanarajah SE, Backes H, DiFeliceantonio AG, Albus K, Cremer AL, Hanssen R, et al. Food intake recruits Orosensory and post-ingestive dopaminergic circuits to affect eating desire in humans. *Cell Metab.* 2019;29(3):695–706. e4. Epub 2019/01/01.
 307. Ritter S, Dinh TT, Li AJ. Hindbrain catecholamine neurons control multiple glucoregulatory responses. *Physiol Behav.* 2006;89(4):490–500. Epub 2006/08/05.
 308. Blouet C, Schwartz GJ. Brainstem nutrient sensing in the nucleus of the solitary tract inhibits feeding. *Cell Metab.* 2012;16(5):579–87. Epub 2012/11/06.
 309. Kim EK, Miller I, Aja S, Landree LE, Pinn M, McFadden J, et al. C75, a fatty acid synthase inhibitor, reduces food intake via hypothalamic AMP-activated protein kinase. *J Biol Chem.* 2004;279(19):19970–6. Epub 2004/03/19.
 310. Nakamura K. Central circuitries for body temperature regulation and fever. *Am J Physiol Regul Integr Comp Physiol.* 2011;301(5):R1207–28. Epub 2011/09/09.
 311. Boulant JA. Role of the preoptic-anterior hypothalamus in thermoregulation and fever. *Clin Infect Dis.* 2000;31(Suppl 5):S157–61. Epub 2000/12/13.
 312. Chen XM, Hosono T, Yoda T, Fukuda Y, Kanosue K. Efferent projection from the preoptic area for the control of non-shivering thermogenesis in rats. *J Physiol.* 1998;512(Pt 3):883–92. Epub 1998/10/14.
 313. Farrell MJ, Trevaks D, Taylor NA, McAllen RM. Regional brain responses associated with thermogenic and psychogenic sweating events in humans. *J Neurophysiol.* 2015;114(5):2578–87. Epub 2015/08/21.
 314. Hefco E, Krulich L, Aschenbrenner JE. Effect of hypothalamic deafferentation on the secretion of thyrotropin during thyroid blockade and exposure to cold in the rat. *Endocrinology.* 1975;97(5):1234–40. Epub 1975/11/01.
 315. Hefco E, Krulich L, Aschenbrenner JE. Effect of hypothalamic deafferentation on the secretion of thyrotropin in resting conditions in the rat. *Endocrinology.* 1975;97(5):1226–33. Epub 1975/11/01.
 316. Hefco E, Krulich L, Illner P, Larsen PR. Effect of acute exposure to cold on the activity of the hypothalamic-pituitary-thyroid system. *Endocrinology.* 1975;97(5):1185–95. Epub 1975/11/01.
 317. Lu J, Zhang YH, Chou TC, Gaus SE, Elmquist JK, Shiromani P, et al. Contrasting effects of ibotenate lesions of the paraventricular nucleus and subparaventricular zone on sleep-wake cycle and temperature regulation. *J Neurosci.* 2001;21(13):4864–74. Epub 2001/06/27.
 318. Watts AG. The efferent projections of the suprachiasmatic nucleus; anatomical insights into the control of circadian rhythms. In: Klein DC, Moore RY, Reppert SM, editors. *Suprachiasmatic nucleus; the mind's clock.* New York: Oxford University Press; 1991. p. 77–106.
 319. Kolka MA, Stephenson LA. Control of sweating during the human menstrual cycle. *Eur J Appl Physiol Occup Physiol.* 1989;58(8):890–5. Epub 1989/01/01.
 320. Lindsley G, Dowse HB, Burgoon PW, Kolka MA, Stephenson LA. A persistent circadian ultradian rhythm is identified in human core temperature. *Chronobiol Int.* 1999;16(1):69–78. Epub 1999/02/19.
 321. Stephenson LA, Kolka MA. Esophageal temperature threshold for sweating decreases before ovulation in premenopausal women. *J Appl Physiol.* 1999;86(1):22–8. Epub 1999/01/14.
 322. McEwen BS. Neural gonadal steroid actions. *Science.* 1981;211(4488):1303–11. Epub 1981/03/20.
 323. Nakayama T, Suzuki M, Ishizuka N. Action of progesterone on preoptic thermosensitive neurones. *Nature.* 1975;258(5530):80. Epub 1975/11/06.
 324. Silva NL, Boulant JA. Effects of testosterone, estradiol, and temperature on neurons in preoptic tissue slices. *Am J Phys.* 1986;250(4 Pt 2):R625–32. Epub 1986/04/01.
 325. Freedman RR. Physiology of hot flashes. *Am J Hum Biol.* 2001;13(4):453–64. Epub 2001/06/16.
 326. Freedman RR, Norton D, Woodward S, Cornelissen G. Core body temperature and circadian rhythm of hot flashes in menopausal women. *J Clin Endocrinol Metab.* 1995;80(8):2354–8. Epub 1995/08/01.
 327. Freedman RR, Dinsay R. Clonidine raises the sweating threshold in symptomatic but not in asymptomatic postmenopausal women. *Fertil Steril.* 2000;74(1):20–3. Epub 2000/07/19.
 328. Cypess AM, Kahn CR. The role and importance of brown adipose tissue in energy homeostasis. *Curr Opin Pediatr.* 2010;22(4):478–84. Epub 2010/05/22.
 329. Nedergaard J, Bengtsson T, Cannon B. Unexpected evidence for active brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab.* 2007;293(2):E444–52. Epub 2007/05/03.
 330. Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, et al. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med.* 2009;360(15):1509–17. Epub 2009/04/10.
 331. Rachid B, van de Sande-Lee S, Rodovalho S, Folli F, Beltrami GC, Morari J, et al. Distinct regula-

- tion of hypothalamic and brown/beige adipose tissue activities in human obesity. *Int J Obes.* 2015;39(10):1515–22. Epub 2015/05/23.
332. Feldmann HM, Golozoubova V, Cannon B, Nedergaard J. UCP1 ablation induces obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by living at thermoneutrality. *Cell Metab.* 2009;9(2):203–9. Epub 2009/02/04.
333. Tam CS, Lecoultré V, Ravussin E. Brown adipose tissue: mechanisms and potential therapeutic targets. *Circulation.* 2012;125(22):2782–91. Epub 2012/06/06.
334. Liu J, Lin L. Small molecules for fat combustion: targeting thermosensory and satiety signals in the central nervous system. *Drug Discov Today.* 2019;24(1):300–6. Epub 2018/09/25.
335. Wang B, Li A, Li X, Ho PW, Wu D, Wang X, et al. Activation of hypothalamic RIP-Cre neurons promotes beiging of WAT via sympathetic nervous system. *EMBO Rep.* 2018;19(4). Epub 2018/02/23.
336. Contreras C, Nogueiras R, Dieguez C, Medina-Gomez G, Lopez M. Hypothalamus and thermogenesis: heating the BAT, browning the WAT. *Mol Cell Endocrinol.* 2016;438:107–15. Epub 2016/08/09.
337. Contreras C, Nogueiras R, Dieguez C, Rahmouni K, Lopez M. Traveling from the hypothalamus to the adipose tissue: the thermogenic pathway. *Redox Biol.* 2017;12:854–63. Epub 2017/04/28.
338. Lopez M, Dieguez C, Nogueiras R. Hypothalamic GLP-1: the control of BAT thermogenesis and browning of white fat. *Adipocytes.* 2015;4(2):141–5. Epub 2015/07/15.
339. Yang X, Ruan HB. Neuronal control of adaptive thermogenesis. *Front Endocrinol.* 2015;6:149. Epub 2015/10/07.
340. Duff GW. Is fever beneficial to the host: a clinical perspective. *Yale J Biol Med.* 1986;59(2):125–30. Epub 1986/03/01.
341. Kluger MJ. The adaptive value of fever. In: Mackowiak PA, editor. *Fever; basic mechanisms and management.* New York: Raven Press; 1991. p. 105–24.
342. Elmquist JK, Saper CB. Activation of neurons projecting to the paraventricular hypothalamic nucleus by intravenous lipopolysaccharide. *J Comp Neurol.* 1996;374(3):315–31. Epub 1996/10/21.
343. Elmquist JK, Scammell TE, Jacobson CD, Saper CB. Distribution of Fos-like immunoreactivity in the rat brain following intravenous lipopolysaccharide administration. *J Comp Neurol.* 1996;371(1):85–103. Epub 1996/07/15.
344. Elmquist JK, Scammell TE, Saper CB. Mechanisms of CNS response to systemic immune challenge: the febrile response. *Trends Neurosci.* 1997;20(12):565–70. Epub 1998/01/07.
345. Contreras C, Gonzalez F, Ferno J, Dieguez C, Rahmouni K, Nogueiras R, et al. The brain and brown fat. *Ann Med.* 2015;47(2):150–68. Epub 2014/06/11.
346. Erden Y, Tekin S, Sandal S, Onalan EE, Tektemur A, Kirbag S. Effects of central irisin administration on the uncoupling proteins in rat brain. *Neurosci Lett.* 2016;618:6–13. Epub 2016/03/02.
347. Ferrante C, Orlando G, Recinella L, Leone S, Chiavaroli A, Di Nisio C, et al. Central inhibitory effects on feeding induced by the adipo-myokine irisin. *Eur J Pharmacol.* 2016;791:389–94. Epub 2016/10/30.
348. Tekin S, Beytur A, Erden Y, Beytur A, Cigremis Y, Vardi N, et al. Effects of intracerebroventricular administration of irisin on the hypothalamus-pituitary-gonadal axis in male rats. *J Cell Physiol.* 2019;234(6):8815–24. Epub 2018/10/15.
349. Tekin S, Erden Y, Ozyalin F, Cigremis Y, Colak C, Sandal S. The effects of intracerebroventricular infusion of irisin on feeding behaviour in rats. *Neurosci Lett.* 2017;645:25–32. Epub 2017/03/01.
350. Tekin S, Erden Y, Ozyalin F, Onalan EE, Cigremis Y, Colak C, et al. Central irisin administration suppresses thyroid hormone production but increases energy consumption in rats. *Neurosci Lett.* 2018;674:136–41. Epub 2018/03/27.
351. Wahab F, Drummer C, Matz-Rensing K, Fuchs E, Behr R. Irisin is expressed by undifferentiated spermatogonia and modulates gene expression in organotypic primate testis cultures. *Mol Cell Endocrinol.* 2019;110670. Epub 2019/12/06.
352. Wahab F, Khan IU, Polo IR, Zubair H, Drummer C, Shahab M, et al. Irisin in the primate hypothalamus and its effect on GnRH in vitro. *J Endocrinol.* 2019;241(3):175–87. Epub 2019/03/27.
353. Kreigsfeld LJ, LeSaute J, Hamada T, Pitts SM, Silver R. Circadian rhythms in the endocrine system. In: Pfaff DW, Arnold AP, Etgen AM, Fahrback SE, Rubin RT, editors. *Hormones, brain and behavior.* Cambridge, Massachusetts: Academic Press, 2002. p. 33–91.
354. Moore RY. Circadian timing. In: Zigmond MJ, Bloom FE, Landis SC, Robetos JL, Squire LR, editors. *Fundamental neuroscience.* Cambridge, Massachusetts: Academic Press, 1999. p. 1189–1208.
355. Shirakawa T, Honma S, Honma K. Multiple oscillators in the suprachiasmatic nucleus. *Chronobiol Int.* 2001;18(3):371–87. Epub 2001/07/28.
356. Welsh DK, Logothetis DE, Meister M, Reppert SM. Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron.* 1995;14(4):697–706. Epub 1995/04/01.
357. Ibuka N, Inouye SI, Kawamura H. Analysis of sleep-wakefulness rhythms in male rats after suprachiasmatic nucleus lesions and ocular enucleation. *Brain Res.* 1977;122(1):33–47. Epub 1977/02/11.
358. Kafka MS, Wirz-Justice A, Naber D, Moore RY, Benedito MA. Circadian rhythms in rat brain neu-

- rotransmitter receptors. *Fed Proc.* 1983;42(11):2796–801. Epub 1983/08/01.
359. Moore RY. Organization and function of a central nervous system circadian oscillator: the suprachiasmatic hypothalamic nucleus. *Fed Proc.* 1983;42(11):2783–9. Epub 1983/08/01.
 360. Ralph MR, Foster RG, Davis FC, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science.* 1990;247(4945):975–8. Epub 1990/02/23.
 361. Parton LE, Ye CP, Coppari R, Enriori PJ, Choi B, Zhang CY, et al. Glucose sensing by POMC neurons regulates glucose homeostasis and is impaired in obesity. *Nature.* 2007;449(7159):228–32. Epub 2007/08/31.
 362. Mrosovsky N. Locomotor activity and non-photic influences on circadian clocks. *Biol Rev Camb Philos Soc.* 1996;71(3):343–72. Epub 1996/08/01.
 363. Zee PC, Manthena P. The brain's master circadian clock: implications and opportunities for therapy of sleep disorders. *Sleep Med Rev.* 2007;11(1):59–70. Epub 2006/09/16.
 364. Tonsfeldt KJ, Chappell PE. Clocks on top: the role of the circadian clock in the hypothalamic and pituitary regulation of endocrine physiology. *Mol Cell Endocrinol.* 2012;349(1):3–12. Epub 2011/07/27.
 365. Gillette MU, Tischkau SA. Suprachiasmatic nucleus: the brain's circadian clock. *Recent Prog Horm Res.* 1999;54:33–58. discussion -9. Epub 1999/11/05.
 366. Moore RY. Neural control of the pineal gland. *Behav Brain Res.* 1996;73(1–2):125–30. Epub 1996/01/01.
 367. Arendt J. Melatonin, circadian rhythms, and sleep. *N Engl J Med.* 2000;343(15):1114–6. Epub 2000/10/12.
 368. Azzali G, Arcari ML, Cacchioli A, Toni R. Fine structure and photoperiodical seasonal changes in pars tuberalis of hibernating bats. *Ital J Anat Embryol.* 2003;108(1):49–64. Epub 2003/05/10.
 369. Azzali G, Arcari ML, Spaggiari B, Romita G. Ultrastructural aspects of the follicular cells of the pars tuberalis in bats related to the seasonal cycle. *Anat Rec A Discov Mol Cell Evol Biol.* 2003;273(2):763–71. Epub 2003/07/08.
 370. Drew KL, Buck CL, Barnes BM, Christian SL, Rasley BT, Harris MB. Central nervous system regulation of mammalian hibernation: implications for metabolic suppression and ischemia tolerance. *J Neurochem.* 2007;102(6):1713–26. Epub 2007/06/09.
 371. Karsch FJ, Bittman EL, Robinson JE, Yellon SM, Wayne NL, Olster DH, et al. Melatonin and photorefractoriness: loss of response to the melatonin signal leads to seasonal reproductive transitions in the ewe. *Biol Reprod.* 1986;34(2):265–74. Epub 1986/03/01.
 372. Fuller PM, Gooley JJ, Saper CB. Neurobiology of the sleep-wake cycle: sleep architecture, circadian regulation, and regulatory feedback. *J Biol Rhythm.* 2006;21(6):482–93. Epub 2006/11/17.
 373. Saper CB, Chou TC, Scammell TE. The sleep switch: hypothalamic control of sleep and wakefulness. *Trends Neurosci.* 2001;24(12):726–31. Epub 2001/11/24.
 374. Sherin JE, Shiromani PJ, McCarley RW, Saper CB. Activation of ventrolateral preoptic neurons during sleep. *Science.* 1996;271(5246):216–9. Epub 1996/01/12.
 375. Lugaresi E. The thalamus and insomnia. *Neurology.* 1992;42(7 Suppl 6):28–33. Epub 1992/07/01.
 376. Steriade M. The thalamus and sleep disturbances. In: Guillemainault C, Lugaresi E, Montagna P, Gambetti P, editors. *Fatal familial insomnia; inherited prion diseases, sleep and the thalamus.* New York: Raven Press; 1994. p. 177–89.
 377. Moruzzi G, Magoun HW. Brain stem reticular formation and activation of the EEG. *Electroencephalogr Clin Neurophysiol.* 1949;1(4):455–73. Epub 1949/11/01.
 378. Canteras NS. The medial hypothalamic defensive system: hodological organization and functional implications. *Pharmacol Biochem Behav.* 2002;71(3):481–91. Epub 2002/02/07.
 379. Swanson LW. Cerebral hemisphere regulation of motivated behavior. *Brain Res.* 2000;886(1–2):113–64. Epub 2000/12/20.
 380. Swanson LW, Sawchenko PE. Paraventricular nucleus: a site for the integration of neuroendocrine and autonomic mechanisms. *Neuroendocrinology.* 1980;31(6):410–7. Epub 1980/12/01.
 381. Blanchard DC, Canteras NS, Markham CM, Pentkowski NS, Blanchard RJ. Lesions of structures showing FOS expression to cat presentation: effects on responsivity to a cat, cat odor, and nonpredator threat. *Neurosci Biobehav Rev.* 2005;29(8):1243–53. Epub 2005/08/09.
 382. McGregor IS, Adamec R, Canteras NS, Blanchard RJ, Blanchard DC. Defensive behavior. *Neurosci Biobehav Rev.* 2005;29(8):1121–2. Epub 2005/08/17.
 383. Ribeiro-Barbosa ER, Canteras NS, Cezario AF, Blanchard RJ, Blanchard DC. An alternative experimental procedure for studying predator-related defensive responses. *Neurosci Biobehav Rev.* 2005;29(8):1255–63. Epub 2005/08/27.
 384. Cezario AF, Ribeiro-Barbosa ER, Baldo MV, Canteras NS. Hypothalamic sites responding to predator threats--the role of the dorsal premammillary nucleus in unconditioned and conditioned antipredatory defensive behavior. *Eur J Neurosci.* 2008;28(5):1003–15. Epub 2008/08/12.
 385. Blanchard DC, Li CI, Hubbard D, Markham CM, Yang M, Takahashi LK, et al. Dorsal premammillary nucleus differentially modulates defensive behaviors induced by different threat stimuli in rats. *Neurosci Lett.* 2003;345(3):145–8. Epub 2003/07/05.
 386. Canteras NS, Simerly RB, Swanson LW. Projections of the ventral premammillary nucleus. *J Comp Neurol.* 1992;324(2):195–212. Epub 1992/10/08.

387. Canteras NS, Swanson LW. The dorsal premammillary nucleus: an unusual component of the mammillary body. *Proc Natl Acad Sci U S A*. 1992;89(21):10089–93. Epub 1992/11/01.
388. Behbehani MM. Functional characteristics of the midbrain periaqueductal gray. *Prog Neurobiol*. 1995;46(6):575–605. Epub 1995/08/01.
389. Lenz KM, McCarthy MM. Organized for sex - steroid hormones and the developing hypothalamus. *Eur J Neurosci*. 2010;32(12):2096–104. Epub 2010/12/15.
390. Motta SC, Goto M, Gouveia FV, Baldo MV, Canteras NS, Swanson LW. Dissecting the brain's fear system reveals the hypothalamus is critical for responding in subordinate conspecific intruders. *Proc Natl Acad Sci U S A*. 2009;106(12):4870–5. Epub 2009/03/11.
391. Lechan RM, Toni R. Functional anatomy of the hypothalamus and pituitary. In: Feingold KR, Anawalt B, Boyce A, et al., editors. *Endotext* [internet]. South Dartmouth (MA): MDText.com, Inc.; Updated; 2016.
392. Kaiser U, Ho KK. Pituitary physiology and diagnostic evaluation. In: Melmed S, Polonsky KS, Reed Larsen P, Kronenberg HM, editors. *Williams textbook of endocrinology*. 13th ed. Elsevier; 2016. p. 176–231.
393. Braak H, Braak E. Anatomy of the human hypothalamus (chiasmatic and tuberal region). *Prog Brain Res*. 1992;93:3–14; discussion -6. Epub 1992/01/01.
394. Chanson P, Daujat F, Young J, Bellucci A, Kujas M, Doyon D, et al. Normal pituitary hypertrophy as a frequent cause of pituitary incidentaloma: a follow-up study. *J Clin Endocrinol Metab*. 2001;86(7):3009–15. Epub 2001/07/10.
395. Cooke NE, Coit D, Weiner RI, Baxter JD, Martial JA. Structure of cloned DNA complementary to rat prolactin messenger RNA. *J Biol Chem*. 1980;255(13):6502–10. Epub 1980/07/10.
396. Suganuma N, Seo H, Yamamoto N, Kikkawa F, Narita O, Tomoda Y, et al. Ontogenesis of pituitary prolactin in the human fetus. *J Clin Endocrinol Metab*. 1986;63(1):156–61. Epub 1986/07/01.
397. Sinha YN. Structural variants of prolactin: occurrence and physiological significance. *Endocr Rev*. 1995;16(3):354–69. Epub 1995/06/01.
398. Lewis UJ, Singh RN, Sinha YN, Vanderlaan WP. Glycosylated human prolactin. *Endocrinology*. 1985;116(1):359–63. Epub 1985/01/01.
399. Hu ZZ, Zhuang L, Meng J, Leondires M, Dufau ML. The human prolactin receptor gene structure and alternative promoter utilization: the generic promoter hPIII and a novel human promoter hP(N). *J Clin Endocrinol Metab*. 1999;84(3):1153–6. Epub 1999/03/20.
400. Bole-Feysot C, Goffin V, Edery M, Binart N, Kelly PA. Prolactin (PRL) and its receptor: actions, signal transduction pathways and phenotypes observed in PRL receptor knockout mice. *Endocr Rev*. 1998;19(3):225–68. Epub 1998/06/17.
401. Liu JW, Ben-Jonathan N. Prolactin-releasing activity of neurohypophyseal hormones: structure-function relationship. *Endocrinology*. 1994;134(1):114–8. Epub 1994/01/01.
402. Horseman ND, Zhao W, Montecino-Rodriguez E, Tanaka M, Nakashima K, Engle SJ, et al. Defective mammapoiesis, but normal hematopoiesis, in mice with a targeted disruption of the prolactin gene. *EMBO J*. 1997;16(23):6926–35. Epub 1998/01/31.
403. Veldhuis JD, Johnson ML. Operating characteristics of the hypothalamo-pituitary-gonadal axis in men: circadian, ultradian, and pulsatile release of prolactin and its temporal coupling with luteinizing hormone. *J Clin Endocrinol Metab*. 1988;67(1):116–23. Epub 1988/07/01.
404. Katznelson L, Riskind PN, Saxe VC, Klibanski A. Prolactin pulsatile characteristics in postmenopausal women. *J Clin Endocrinol Metab*. 1998;83(3):761–4. Epub 1998/03/20.
405. Riddle O. Prolactin in vertebrate function and organization. *J Natl Cancer Inst*. 1963;31:1039–110. Epub 1963/11/01.
406. Matsuzaki T, Azuma K, Irahara M, Yasui T, Aono T. Mechanism of anovulation in hyperprolactinemic amenorrhea determined by pulsatile gonadotropin-releasing hormone injection combined with human chorionic gonadotropin. *Fertil Steril*. 1994;62(6):1143–9. Epub 1994/12/01.
407. Ho Y, Liebhaber SA, Cooke NE. Activation of the human GH gene cluster: roles for targeted chromatin modification. *Trends Endocrinol Metab*. 2004;15(1):40–5. Epub 2003/12/25.
408. Baumann G, MacCart JG, Amburn K. The molecular nature of circulating growth hormone in normal and acromegalic man: evidence for a principal and minor monomeric forms. *J Clin Endocrinol Metab*. 1983;56(5):946–52. Epub 1983/05/01.
409. Leung KC, Waters MJ, Markus I, Baumbach WR, Ho KK. Insulin and insulin-like growth factor-I acutely inhibit surface translocation of growth hormone receptors in osteoblasts: a novel mechanism of growth hormone receptor regulation. *Proc Natl Acad Sci U S A*. 1997;94(21):11381–6. Epub 1997/10/23.
410. Xu BC, Wang X, Darus CJ, Kopchick JJ. Growth hormone promotes the association of transcription factor STAT5 with the growth hormone receptor. *J Biol Chem*. 1996;271(33):19768–73. Epub 1996/08/16.
411. Le Roith D, Scavo L, Butler A. What is the role of circulating IGF-I? *Trends Endocrinol Metab*. 2001;12(2):48–52. Epub 2001/02/13.
412. Giustina A, Veldhuis JD. Pathophysiology of the neuroregulation of growth hormone secretion in experimental animals and the human. *Endocr Rev*. 1998;19(6):717–97. Epub 1998/12/23.
413. Ho KY, Veldhuis JD, Johnson ML, Furlanetto R, Evans WS, Alberti KG, et al. Fasting enhances growth hormone secretion and amplifies the com-

- plex rhythms of growth hormone secretion in man. *J Clin Invest.* 1988;81(4):968–75. Epub 1988/04/01.
414. Van Cauter E, Plat L, Copinschi G. Interrelations between sleep and the somatotrophic axis. *Sleep.* 1998;21(6):553–66. Epub 1998/10/21.
 415. Rudling M, Norstedt G, Olivecrona H, Reihner E, Gustafsson JA, Angelin B. Importance of growth hormone for the induction of hepatic low density lipoprotein receptors. *Proc Natl Acad Sci U S A.* 1992;89(15):6983–7. Epub 1992/08/01.
 416. Moller N, Jorgensen JO. Effects of growth hormone on glucose, lipid, and protein metabolism in human subjects. *Endocr Rev.* 2009;30(2):152–77. Epub 2009/02/26.
 417. Lamolet B, Pulichino AM, Lamonerie T, Gauthier Y, Brue T, Enjalbert A, et al. A pituitary cell-restricted T box factor, Tpit, activates POMC transcription in cooperation with Pitx homeoproteins. *Cell.* 2001;104(6):849–59. Epub 2001/04/06.
 418. Cochet M, Chang AC, Cohen SN. Characterization of the structural gene and putative 5'-regulatory sequences for human proopiomelanocortin. *Nature.* 1982;297(5864):335–9. Epub 1982/05/27.
 419. Clark AJL, Swords FM. Molecular pathology of corticotroph function. In: Rappaport R, Amselem S, editors. *Hypothalamic-pituitary development.* Basel: Karger; 2001.
 420. Jenks BG. Regulation of proopiomelanocortin gene expression: an overview of the signaling cascades, transcription factors, and responsive elements involved. *Ann N Y Acad Sci.* 2009;1163:17–30. Epub 2009/05/22.
 421. Jin WD, Boutillier AL, Glucksman MJ, Salton SR, Loeffler JP, Roberts JL. Characterization of a corticotropin-releasing hormone-responsive element in the rat proopiomelanocortin gene promoter and molecular cloning of its binding protein. *Mol Endocrinol.* 1994;8(10):1377–88. Epub 1994/10/01.
 422. Seidah NG, Chretien M. Complete amino acid sequence of a human pituitary glycopeptide: an important maturation product of proopiomelanocortin. *Proc Natl Acad Sci U S A.* 1981;78(7):4236–40. Epub 1981/07/01.
 423. Keeney DS, Waterman MR. Regulation of steroid hydroxylase gene expression: importance to physiology and disease. *Pharmacol Ther.* 1993;58(3):301–17. Epub 1993/06/01.
 424. Veldhuis JD, Iranmanesh A, Johnson ML, Lizarralde G. Twenty-four-hour rhythms in plasma concentrations of adenohypophyseal hormones are generated by distinct amplitude and/or frequency modulation of underlying pituitary secretory bursts. *J Clin Endocrinol Metab.* 1990;71(6):1616–23. Epub 1990/12/01.
 425. Dorin RI, Ferries LM, Roberts B, Qualls CR, Veldhuis JD, Lisansky EJ. Assessment of stimulated and spontaneous adrenocorticotropin secretory dynamics identifies distinct components of cortisol feedback inhibition in healthy humans. *J Clin Endocrinol Metab.* 1996;81(11):3883–91. Epub 1996/11/01.
 426. Gharib SD, Wierman ME, Shupnik MA, Chin WW. Molecular biology of the pituitary gonadotropins. *Endocr Rev.* 1990;11(1):177–99. Epub 1990/02/01.
 427. Abreu AP, Dauber A, Macedo DB, Noel SD, Brito VN, Gill JC, et al. Central precocious puberty caused by mutations in the imprinted gene MKRN3. *N Engl J Med.* 2013;368(26):2467–75. Epub 2013/06/07.
 428. Ying SY. Inhibins, activins, and follistatins: gonadal proteins modulating the secretion of follicle-stimulating hormone. *Endocr Rev.* 1988;9(2):267–93. Epub 1988/05/01.
 429. Kristrom B, Zdunek AM, Rydh A, Jonsson H, Sehlin P, Escher SA. A novel mutation in the LIM homeobox 3 gene is responsible for combined pituitary hormone deficiency, hearing impairment, and vertebral malformations. *J Clin Endocrinol Metab.* 2009;94(4):1154–61. Epub 2009/01/08.
 430. Moschos S, Chan JL, Mantzoros CS. Leptin and reproduction: a review. *Fertil Steril.* 2002;77(3):433–44. Epub 2002/03/02.
 431. Naor Z. Signaling by G-protein-coupled receptor (GPCR): studies on the GnRH receptor. *Front Neuroendocrinol.* 2009;30(1):10–29. Epub 2008/08/19.
 432. Santen RJ, Bardin CW. Episodic luteinizing hormone secretion in man. Pulse analysis, clinical interpretation, physiologic mechanisms. *J Clin Invest.* 1973;52(10):2617–28. Epub 1973/10/01.
 433. Richards JS, Pangas SA. The ovary: basic biology and clinical implications. *J Clin Invest.* 2010;120(4):963–72. Epub 2010/04/07.
 434. Themmen APN, Huhtaniemi IT. Mutations of gonadotropins and gonadotropin receptors: elucidating the physiology and pathophysiology of pituitary-gonadal function. *Endocr Rev.* 2000;21(5):551–83. Epub 2000/10/21.
 435. Ruwanpura SM, McLachlan RI, Meachem SJ. Hormonal regulation of male germ cell development. *J Endocrinol.* 2010;205(2):117–31. Epub 2010/02/11.
 436. Pierce JG, Parsons TF. Glycoprotein hormones: structure and function. *Annu Rev Biochem.* 1981;50:465–95. Epub 1981/01/01.
 437. Grossmann M, Weintraub BD, Szkudlinski MW. Novel insights into the molecular mechanisms of human thyrotropin action: structural, physiological, and therapeutic implications for the glycoprotein hormone family. *Endocr Rev.* 1997;18(4):476–501. Epub 1997/08/01.
 438. Lania A, Persani L, Ballare E, Mantovani S, Losa M, Spada A. Constitutively active Gs alpha is associated with an increased phosphodiesterase activity in human growth hormone-secreting adenomas. *J*

- Clin Endocrinol Metab. 1998;83(5):1624–8. Epub 1998/05/20.
439. Chiamolera MI, Wondisford FE. Minireview: Thyrotropin-releasing hormone and the thyroid hormone feedback mechanism. *Endocrinology*. 2009;150(3):1091–6. Epub 2009/01/31.
 440. Lechan RM, Fekete C. The TRH neuron: a hypothalamic integrator of energy metabolism. *Prog Brain Res*. 2006;153:209–35. Epub 2006/08/01.
 441. Abel ED, Kaulbach HC, Campos-Barros A, Ahima RS, Boers ME, Hashimoto K, et al. Novel insight from transgenic mice into thyroid hormone resistance and the regulation of thyrotropin. *J Clin Invest*. 1999;103(2):271–9. Epub 1999/01/23.
 442. Samuels MH, Henry P, Ridgway EC. Effects of dopamine and somatostatin on pulsatile pituitary glycoprotein secretion. *J Clin Endocrinol Metab*. 1992;74(1):217–22. Epub 1992/01/01.
 443. Cooper DS, Klibanski A, Ridgway EC. Dopaminergic modulation of TSH and its subunits: in vivo and in vitro studies. *Clin Endocrinol*. 1983;18(3):265–75. Epub 1983/03/01.
 444. Wang R, Nelson JC, Wilcox RB. Salsalate administration—a potential pharmacological model of the sick euthyroid syndrome. *J Clin Endocrinol Metab*. 1998;83(9):3095–9. Epub 1998/09/24.
 445. Ridgway EC, Weintraub BD, Maloof F. Metabolic clearance and production rates of human thyrotropin. *J Clin Invest*. 1974;53(3):895–903. Epub 1974/03/01.
 446. Van den Berghe G, de Zegher F, Veldhuis JD, Wouters P, Gouwy S, Stockman W, et al. Thyrotrophin and prolactin release in prolonged critical illness: dynamics of spontaneous secretion and effects of growth hormone-secretagogues. *Clin Endocrinol*. 1997;47(5):599–612. Epub 1998/01/13.
 447. Goichot B, Weibel L, Chapotot F, Gronfier C, Piquard F, Brandenberger G. Effect of the shift of the sleep-wake cycle on three robust endocrine markers of the circadian clock. *Am J Phys*. 1998;275(2):E243–8. Epub 1998/08/04.
 448. Roelfsema F, Pijl H, Kok P, Endert E, Fliers E, Biermasz NR, et al. Thyrotropin secretion in healthy subjects is robust and independent of age and gender, and only weakly dependent on body mass index. *J Clin Endocrinol Metab*. 2014;99(2):570–8. Epub 2013/11/28.
 449. Makarenko IG, Ugrumov MV, Derer P, Calas A. Projections from the hypothalamus to the posterior lobe in rats during ontogenesis: 1,1'-dioctadecyl-3,3,3', 3'-tetramethylindocarbocyanine perchlorate tracing study. *J Comp Neurol*. 2000;422(3):327–37. Epub 2000/06/22.
 450. Robinson AG. The posterior pituitary. In: Gardner DG, Shoback D, editors. *Greenspan's basic and clinical endocrinology*. 10th ed. New York: McGraw-Hill; 2017. p. 121–36.
 451. Sawchenko PE, Swanson LW. The organization and biochemical specificity of afferent projections to the paraventricular and supraoptic nuclei. *Prog Brain Res*. 1983;60:19–29. Epub 1983/01/01.
 452. Sofroniew MV. Morphology of vasopressin and oxytocin neurones and their central and vascular projections. *Prog Brain Res*. 1983;60:101–14. Epub 1983/01/01.
 453. Treier M, Rosenfeld MG. The hypothalamic-pituitary axis: co-development of two organs. *Curr Opin Cell Biol*. 1996;8(6):833–43. Epub 1996/12/01.
 454. Robinson AG, Verbalis JG. Posterior Pituitary. In: Melmed S, Polonsky KS, Reed Larsen P, Kronenberg HM, editors. *Williams textbook of endocrinology*. 13th ed: Elsevier; 2015. p. 300–33.
 455. Roberts MM, Robinson AG, Hoffman GE, Fitzsimmons MD. Vasopressin transport regulation is coupled to the synthesis rate. *Neuroendocrinology*. 1991;53(4):416–22. Epub 1991/04/01.
 456. Barberis C, Mouillac B, Durroux T. Structural bases of vasopressin/oxytocin receptor function. *J Endocrinol*. 1998;156(2):223–9. Epub 1998/03/31.
 457. Thrasher TN. Baroreceptor regulation of vasopressin and renin secretion: low-pressure versus high-pressure receptors. *Front Neuroendocrinol*. 1994;15(2):157–96. Epub 1994/06/01.
 458. Pump B, Gabrielsen A, Christensen NJ, Bie P, Bestle M, Norsk P. Mechanisms of inhibition of vasopressin release during moderate antithrostatic posture change in humans. *Am J Phys*. 1999;277(1):R229–35. Epub 1999/07/17.
 459. Johnson AK, Thunhorst RL. The neuroendocrinology of thirst and salt appetite: visceral sensory signals and mechanisms of central integration. *Front Neuroendocrinol*. 1997;18(3):292–353. Epub 1997/07/01.
 460. Robertson GL. The regulation of vasopressin function in health and disease. *Recent Prog Horm Res*. 1976;33:333–85. Epub 1976/01/01.
 461. Eto K, Noda Y, Horikawa S, Uchida S, Sasaki S. Phosphorylation of aquaporin-2 regulates its water permeability. *J Biol Chem*. 2010;285(52):40777–84. Epub 2010/10/26.
 462. Brown D. The ins and outs of aquaporin-2 trafficking. *Am J Physiol Renal Physiol*. 2003;284(5):F893–901. Epub 2003/04/05.
 463. Nielsen S, Frokiaer J, Marples D, Kwon TH, Agre P, Knepper MA. Aquaporins in the kidney: from molecules to medicine. *Physiol Rev*. 2002;82(1):205–44. Epub 2002/01/05.
 464. Glasier A, McNeilly AS. Physiology of lactation. *Baillieres Clin Endocrinol Metab*. 1990;4(2):379–95. Epub 1990/06/01.
 465. Terzidou V, Blanks AM, Kim SH, Thornton S, Bennett PR. Labor and inflammation increase the expression of oxytocin receptor in human amnion. *Biol Reprod*. 2011;84(3):546–52. Epub 2010/10/12.

466. Vrachnis N, Malamas FM, Sifakis S, Deligeoroglou E, Iliodromiti Z. The oxytocin-oxytocin receptor system and its antagonists as tocolytic agents. *Int J Endocrinol*. 2011;2011:350546. Epub 2011/12/23.
467. Boyd WH. Morphological features of the hypophyseal intermediate lobe directly related to its activity. *Arch Histol Jpn*. 1972;34(1):1–17. Epub 1972/01/01.
468. Brander J. The Intraglandular cleft of the pituitary body and its connections. *J Anat*. 1932;66(Pt 2):202–9. Epub 1932/01/01.
469. Howe A. The mammalian pars intermedia: a review of its structure and function. *J Endocrinol*. 1973;59(2):385–409. Epub 1973/11/01.
470. Wood S, Loudon A. The pars tuberalis: the site of the circannual clock in mammals? *Gen Comp Endocrinol*. 2018;258:222–35. Epub 2017/07/04.
471. Korf HW. Leopoldina symposium “seasonal rhythms”, Leuven Belgium, 25. 8. 2016. *Gen Comp Endocrinol*. 2018;258:213–4. Epub 2017/12/06.
472. Korf HW. Signaling pathways to and from the hypophysial pars tuberalis, an important center for the control of seasonal rhythms. *Gen Comp Endocrinol*. 2018;258:236–43. Epub 2017/05/18.
473. Pfeffer M, Korf HW, Wicht H. Synchronizing effects of melatonin on diurnal and circadian rhythms. *Gen Comp Endocrinol*. 2018;258:215–21. Epub 2017/05/24.

Part II

Pathobiology and Dysfunction of the Hypothalamus

Imaging Aspects of Pathologies of the Sella, the Pituitary Gland, and the Hypothalamus

Manuel Schmidt and Arnd Doerfler

Introduction

Imaging of the sellar region, the pituitary gland, and the hypothalamus is quite challenging since there are large vessels, subarachnoid space, bone, and air-filled cavities in close neighborhood. Varieties of pathologies arise in the central skull base due to adjacent neuronal, vascular, and endocrine structures in this anatomical region. The pituitary gland itself is a very small-volume organ. High-quality imaging necessitates high contrast and spatial resolution to depict very subtle pathologies. Additionally, anatomic variations can hamper differential diagnosis. Magnetic resonance imaging is the modality of choice providing multiplanar high-contrast images of the pituitary gland, the hypothalamus, and the adjacent structures. Computed tomography is used only as supplementary, i.e., when information about the osseous anatomy needs to be obtained or to exclude or visualize calcifications.

Located in the pituitary fossa (Fig. 6.1), the pituitary gland can be morphologically and functionally divided into two parts – the anterior (adenohypophysis) and posterior (neurohypophysis) lobes. Embryologically, the distal part of the

adenohypophysis arises from the epithelium of Rathke's pouch, an invagination of the roof of the oropharyngeal membrane. As part of the brain, the neurohypophysis is composed of the stalk (infundibulum) and the neural lobe (infundibular process). The pars intermedia, derived from the posterior wall of Rathke's pouch, is located in-between the anterior and posterior lobes and is usually not seen in MRI. The posterior lobe of the pituitary gland and the pituitary stalk receive their blood supply from the superior and inferior hypophyseal branches of the internal carotid artery, whereas the anterior lobe receives its blood supply from penetrating capillary loops from the portal vessels of the hypophyseal-portal circulation, respectively.

The adenohypophysis is producing a variety of hormones, i.e., prolactin, growth hormone (GH), thyroid-stimulating hormone (TSH), follicle-stimulating hormone (FSH), and luteinizing hormone (LH). In addition, prohormone precursors of corticotropin (ACTH) and melanocyte-stimulating hormone are secreted, respectively. Thus, lesions of the adenohypophysis may cause hormonal deficiency resulting in a variety of clinical symptoms. The posterior pituitary lobe has no independent secretory function and receives vasopressin (ADH) and oxytocin from the hypothalamic neurons through capillaries for storage. In newborns up to 3 months of age, both anterior and posterior pituitary lobes exhibit hyperintensity on T1-weighted imaging [1, 2]. With further age, the adenohy-

M. Schmidt (✉) · A. Doerfler
Department of Neuroradiology, University Hospital
of Erlangen Medical School, Friedrich-Alexander-
University Erlangen-Nürnberg, Erlangen, Germany
e-mail: manuel.schmidt@uk-erlangen.de; arnd.doerfler@uk-erlangen.de




Fig. 6.1 Normal hypophysis. Coronal and sagittal T2w TSE (**a + b**). Sagittal T1w native (**b**). Note the bright signal of the normal neurohypophysis in T1w (**c**, arrow)




Fig. 6.2 Normal hypothalamus (arrows), hypointense to the adjacent structures on T1w. Ultrahigh resolution T1w MP2RAGE (0.6 mm isotropic voxels). The anterior commissure can be clearly delineated as superior border (**a**,

coronal reformation, empty arrow). Medial border is the third ventricle, while there is no distinct lateral border. Superior and posterior border is the thalamus (**b** and **c**, sagittal and axial reformations)

pophysis loses its hyperintensity gradually, whereas the neurohypophysis remains hyperintense [3] (Fig. 6.1b). Experimental studies have shown that the high signal intensity of the posterior lobe is caused by accumulated neurosecretory granules containing ADH. Thus, in patients with a central diabetes insipidus, the high signal of the posterior lobe is absent, returning after appropriate medical substitution [4].

As the name suggests, the hypothalamus is anatomically located below the thalamus (Fig. 6.2). It is intimately associated with both the limbic system and the pituitary gland. Its boundaries are in part not clearly defined:

- Superior: between the anterior and posterior commissures; thalamus
- Anterior: lamina terminalis, with optic chiasm at the lower border and anterior commissure above
- Medial: third ventricle
- Posterior: a line running antero-inferiorly from the posterior commissure to the mammillary bodies; thalamus
- Lateral: no distinct border
- Inferior: infundibular stalk, tuber cinereum, and mammillary bodies

Imaging Techniques

A standard MRI protocol for examination of the pituitary and parasellar region consists of thin sliced (max. 2 mm) sagittal and coronal T1-weighted images with and without application of contrast medium (Table 6.1). Angulation of the coronal images should be parallel to the pituitary stalk. Thin sliced, sagittal T2-weighted images can be added for evaluation of cystic

Table 6.1 Possible protocol for MR imaging of the sella

	Description
MRI characteristics	Field strength of 1.5 T or 3 T (Tesla), slice thickness of 1.5–2 mm, small field of view (FoV), high matrix and in-plane resolution, ideally 0.5×0.5 mm
	T2-weighting (T2w) in coronal orientation, parallel to pituitary stalk (sagittal T2w localizer); additionally at least one plane in axial or sagittal orientation
	T1-weighting (T1w) native in coronal and sagittal orientation
	T1w coronal and sagittal after administration of i.v. contrast agent (e.g., 0.05 mmol/kg bodyweight (microadenoma) and 0.1 mmol/kg bodyweight (macroadenoma) of gadobutrol, respectively)
	3D T1 MPRAGE or 3D T1 SPACE+Gd volume dataset with 1 mm isotropic voxels
Dynamic hypophysis (T1w + Gd)	Flow rate: 2 ml/s
	Time of injection: start of the second dynamic measurement
	Duration of dynamic measurement: max. 30 s
	Number of measurements: 6–8
	Sequence: coronal T1w turbo-spin-echo or T1w VIBE (GRASP or TWIST)
	Voxel size: $0.5 \times 0.5 \times 2$ mm
	Flush: 20 ml NaCl-Flush with a flow rate of 2.0 ml/s

lesions. We administer half of the standard dose of Gd-DTPA (0,05 mmol/kg) to search for microadenomas. In dynamic sequences after rapid injection of contrast medium, a diminished enhancement of microadenomas may be seen. Studies of normal pituitary tissue have also shown an earlier enhancement of the posterior lobe, due to the direct blood supply of the neurohypophysis via hypophyseal branches of the internal carotid artery. Adenomas also have direct arterial blood supply similar to the neurohypophysis. Exploiting this pattern of blood supply, some microadenomas can only be diagnosed using dynamic T1-weighted images. Additionally, there should be one scan covering the whole brain (T2w or FLAIR).

Computed tomography (CT) may become important when supportive information concerning bony structures or calcifications is required. CT is used in extensively growing pituitary adenomas, invading the sphenoid sinus, nasal cavity, or the skull base. Additionally, the anatomy of the sphenoid sinus can be evaluated as a precursor to transsphenoidal surgery. To exclude acute pituitary hemorrhage, CT may still be helpful in the emergency situation.

Nowadays, conventional radiography is no longer important in the diagnostic workup of pituitary adenomas. Adenomas are usually very

slow-growing lesions, resulting in a vertical and horizontal enlargement of the bony pituitary fossa (balloon sella) with no demineralization. Asymmetric growth of pituitary adenomas may lead to a double outline of the sellar fundus in the lateral view. Thinning, destruction, or dorsal shift of the dorsum sellae might also occur. In patients with central Cushing's disease, if no adenoma is visible in MRI, selective inferior petrous sinus venous blood sampling is a highly specific technique that might be helpful in the diagnosis and especially lateralization of the microadenoma and might guide selected surgical resection.

Developmental Lesions of the Sella and the Hypothalamus

Rathke's Cleft Cyst

Most sellar epithelial cysts are remnants of derivatives of the Rathke's cleft and arise in the region of the pars intermedia. They are relatively common incidental findings at autopsy (up to 30%) and usually remain asymptomatic [5]. Location is usually intrasellar, between the anterior and posterior pituitary lobes (Fig. 6.4c). Occasionally, they can occur in the suprasellar region, anterior to the infundibulum. Symptoms may result from

mass effect resulting in headache, endocrine dysfunction, or visual impairment due to compression of the optic chiasm.

Differentiation from craniopharyngioma is very important. Histologically, these lesions can be distinguished by the composition of their walls. Rathke's cleft cysts can contain mucinous or serous fluid and thus return variable signals in MRI. Cysts containing serous fluid are typically hypointense, whereas mucoid cysts show hyperintensity on T1-weighted images [5, 6]. Sometimes differentiation from acute hemorrhage can be difficult. Clinical symptoms and follow-up imaging are helpful in this setting.

Arachnoidal Cysts

While suprasellar arachnoidal cysts usually present with symptoms due to the local mass effect in children, the rare intrasellar arachnoidal cysts are regarded as acquired and may become symptomatic later in life [7–9]. Clinical symptoms may include increased intracranial pressure, hormone deficiency, gait disturbance, and visual impairment. Arachnoidal cysts arise from herniation of an arachnoidal diverticulum through an incomplete diaphragma sellae. Although usually indistinguishable from Rathke's cleft cysts, they typically displace the anterior lobe and the infundibulum posteriorly [8]. On MRI, a focal mass with CSF signal intensity might be seen.

Ectopic Posterior Lobe

An ectopic posterior lobe is usually found in diagnostic imaging studies for growth hormone deficiency [10–13]. It is thought to be a developmental anomaly, rather than a residuum of traumatic (birth) incidents. Patients become conspicuous with an isolated growth hormone deficiency or multiple anterior pituitary lobe hormone deficiencies. On T1-weighted images, the ectopic posterior lobe is typically seen as a small nodule with characteristic high signal at the median eminence in the floor of the third ventricle [14–16] (Fig. 6.3). Sometimes, a tender pituitary stalk can only be seen after contrast enhancement.

Dermoid and Epidermoid Tumors

Both dermoid and epidermoid tumors are benign, slow-growing congenital lesions which result from inclusion of epithelial elements during embryogenesis. They may cause mass effect in the sellar, parasellar, or suprasellar region resulting in visual disturbance or endocrine dysfunction and consist of less than 2% of all intracranial neoplasms. Both lesions are often hypointense in T1- and hyperintense in T2-weighted images. Depending upon fat and calcium content, dermoid tumors can also show a hyperintense signal in T1 [17, 18]. For epidermoid tumors, diffusion-




Fig. 6.3 Ectopic posterior lobe of the pituitary gland with bright signal in T1w. Sagittal (a) and coronal T1w (b)




Fig. 6.4 Typical hamartoma of the tuber cinereum. Note the iso- to slight hyperintense signal with regard to the cortex in T2w (a, b). No enhancement in T1w + Gd (c). Secondary finding: Rathke's cleft cyst (empty arrow)

weighted images can be helpful, typically showing a markedly increased signal.

Hypothalamic Hamartomas

Hypothalamic hamartomas are of neuronal origin and represent congenital heterotopias usually located within the tuber cinereum. They usually affect children, who present with precocious puberty and epileptic seizures (gelastic seizures) [19]. Characteristically, MRI and CT show a rounded expansion of the tuber cinereum, best seen in coronal and sagittal images. Hamartomas are iso- to hyperintense to the cerebral cortex in T2-weighted images and isointense to the cortex in T1w images (Fig. 6.4). Because they are rarely larger than 1–2 cm in diameter, little mass effect is seen [20, 21].

Neoplasms of the Sella and the Hypothalamus

Pituitary Adenomas

Pituitary adenomas are benign epithelial lesions and account for about 10–15% of all intracranial tumors and thus represent the most common intrasellar pathology. In unselected autopsy series, the estimated incidence is up to 27%. Using the size as a criteria, adenomas larger than

10 mm are classified as macroadenomas, whereas tumors smaller than 10 mm are referred to as microadenomas (Fig. 6.5.). Classification concerning endocrine function distinguish hormone secreting from nonsecreting (nonfunctional) adenomas.

Prolactin-secreting adenomas are the most common secretive tumors and account for about 30% of all pituitary adenomas. Clinical manifestations in women are secondary amenorrhea, galactorrhea, and infertility [22]. In men, loss of libido and impotence can occur [23, 24]. Because of the varying symptoms, prolactinomas in men are often diagnosed at a later stage. Acromegaly in adults and gigantism in children are the cardinal symptoms of growth hormone-producing adenomas [25]. Incidence peak lies in the fifth life decade, causing growth of the feet and hands as well as coarsening of facial features like the nose and chin. Retrospective studies indicate that mortality is approximately doubled relative to the general population, mostly due to cardiovascular events [26, 27]. Their insidious onset often leads to significant delays in diagnosis.

About 5–10% of pituitary tumors cause elevated glucocorticoid levels (ACTH-producing adenomas). Overproduction leads to the stigmata of Cushing's disease, including diabetes, hypertension, osteoporosis, easy skin bruisability and striae rubrae, truncal obesity, moon facies, amenorrhea, impotence, and a generalized weakness. In children, growth retardation is a common




Fig. 6.5 Typical microadenoma (histologically prolactinoma), hyperintense in T2w, in the left aspect of the pituitary gland. Coronal T2w (a) and T1w + Gd (b)

manifestation [28]. Treatment can either be surgically or conservative with neuromodulatory drugs (i.e., bromocriptine) [29]. Patients with Cushing's syndrome that are treated surgically with bilateral adrenalectomy can develop an aggressive ACTH-producing adenoma (Nelson tumor) in about 15% [30, 31] due to the lack of the negative feedback loop.

Thyroid-stimulating hormone-producing adenomas occur in <1%, leading to hyperthyroidism [32]. Aberrant secretion of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) is very rare, causing symptoms such as hypogonadism. About 10% of all adenomas are mixed tumors, out of which 75% are macroadenomas.

Nonfunctioning adenomas are the second most common tumors, comprising 25–30% of pituitary adenomas. The majority are macroadenomas that often grow to a significant size, causing visual field defects, hydrocephalus, or other mass effects. Parasellar infiltration into the cavernous sinus occurs in 40%, rarely leading to cranial nerve palsies [33, 34]. Cranial nerve palsy in combination with a sellar mass rather points towards metastasis or ophthalmoplegic aneurysm [35, 36].

Microadenomas are best evaluated in coronal images and usually appear hypo- or isointense relative to normal pituitary tissue on unenhanced T1-weighted images [3, 37]. On T2w images

they often appear hyperintense. After contrast administration, due to an earlier and more intense enhancement of normal pituitary tissue, the microadenoma usually remains hypointense. Indirect radiologic features of microadenomas may be a one-sided elevation of the diaphragm or lapse of the pituitary fossa. In dynamic sequences after rapid injection of contrast medium, a diminished enhancement of the adenoma may be seen [38–41]. Studies of normal pituitary tissue have also shown an earlier enhancement of the posterior lobe, due to the direct blood supply of the neurohypophysis via hypophyseal branches of the internal carotid artery. Using fast T1-weighted images, some microadenomas demonstrate an early arterial enhancement, occurring simultaneously as enhancement of the posterior lobe [42]. However, compared with normal pituitary tissue, most microadenomas display a slightly lower signal intensity.

Macroadenomas are also best evaluated with MR imaging. These tumors usually extend beyond the sella and often show infiltration of the cavernous sinus, the sphenoid sinus, or the clivus. Compression of the optic chiasm and encasement of the internal carotid artery can be present [43]. Cavernous sinus invasion often restricts complete surgical tumor resection. In contrast to parasellar meningiomas, encasement of the internal carotid artery rarely causes luminal compromise in pitu-




Fig. 6.6 Typical left sided macroadenoma with infiltration of the cavernous sinus und compression of the oculomotor nerve. Coronal T2w (a) and T1w + Gd (b)

itary adenomas. In about 94% macroadenomas lead to enlargement of the bony pituitary fossa (Fig. 6.6).

Intratumoral hemorrhage of macroadenomas occurs in up to 10–15% of incidental pituitary adenomas [44]. Medical treatment with bromocriptine and pregnancy are typical predisposing factors associated with pituitary apoplexy and hemorrhage [45, 46]. Primary pituitary hemorrhage may also occur traumatically, after surgery, after viral diseases or radiation, or during delivery. Acute hemorrhagic infarction of a pituitary adenoma is referred to as pituitary apoplexy with an impressive clinical presentation, i.e., acute onset of headache, vomiting, ophthalmoplegia, and visual loss [47, 48]. Rarely, meningism, loss of consciousness, or seizures can occur.

Other Pituitary Neoplasms

Primary tumors of the posterior lobe of the pituitary, e.g., pituicytomas or gangliogliomas, are very rare lesions.

Pituitary metastases most commonly arise from the common cancers such as breast, thyroid, or bronchogenic cancer and occur in approximately 3% of terminally ill patients [49]. Only about 5–15% of these patients become symptomatic [36]. Metastases tend to arise within the posterior lobe of the pituitary or the pituitary stalk and spread to the anterior lobe of the pituitary. Typical MRI findings include a relatively small

enhancing pituitary lesion and lack of sella enlargement. Bony destruction may occur. Third nerve palsy in combination with a sellar mass rather points towards metastasis or ophthalmoplegic aneurysm and is unlikely in the case of adenoma. Rarely, pituitary adenomas can transform to pituitary carcinomas (0.1–0.5% of all pituitary tumors) [50, 51]. They are locally destructive and can metastasize to intracranial and intraspinal sites via CSF pathways. Hepatic, bronchial, bone, and lymphatic spread are also reported.

Hypothalamic Glioma/Optic Pathway Glioma

Sometimes the distinction between optic pathway gliomas and hypothalamic gliomas is not clear. In cases where a tumor is confined to the optic nerve, it can be referred to as optic nerve glioma. Often, however, they are either centered on or extend to involve the chiasm and optic radiations. In such cases, they are difficult to distinguish from hypothalamic gliomas, and such a distinction is arbitrary in most cases. In these cases, the term hypothalamic-optochiasmatic glioma should be used. Generally, the term optic pathway glioma is favored, recognizing that there may be involvement of the hypothalamus.

Optic pathway gliomas are relatively uncommon tumors, with a variable clinical course. They are often seen in the setting of neurofibromatosis type I (NF1) and are frequently bilateral.




Fig. 6.7 Vividly enhancing mass of the right hypothalamus, extending along the optic tract (**a** and **b**, coronal and sagittal T1w + Gd). Hyperintense signal in T2w (**c**, sagit-

tal). Note the small enhancing secondary manifestation in the contralateral hypothalamus (empty arrow)

Histologically, the majority are pilocytic astrocytomas. They are characterized on imaging by an enlarged optic nerve. These tumors usually return low signal on T1w and high signal on T2w images, enhancement of contrast agent is often vivid but variable (Fig. 6.7). A widely accepted classification of optic pathway gliomas is the Dodge classification [52]. This simple classification divides the tumors into three groups based on anatomical localization: stage 1, optic nerves only; stage 2, chiasm involved (with or without optic nerve involvement); and stage 3, hypothalamic involvement and/or other adjacent structures.

Optic pathway gliomas demonstrate variable clinical and radiological progression. In patients with NF1, it is not unusual for these tumors to be clinically quiescent, with little progression over some years. In others, the tumors are more aggressive (Fig. 6.8) and extend along the optic pathways; therefore, treatment depends on the clinical situation.

Meningiomas

Meningiomas of the sellar region (cavernous sinus, planum sphenoidale, diaphragma sellae, clinoid process) account for 20–30% of all intracranial meningiomas. They are benign, slow-growing tumors that can reach considerable size at the time of diagnosis.

To distinguish meningiomas from neurinomas, dynamic contrast enhancing MRI sequences are useful, since meningiomas usually show early enhancement, while neurinomas present gradual enhancement [53, 54].

Due to an en plaque growth pattern often associated with a “dural-tail,” demarcation from pituitary macroadenomas is usually possible. Pure intrasellar meningiomas are very rare and may be hard to distinguish from adenomas [55, 56]. They usually originate from the dorsum sellae. When invading into the cavernous sinus, meningiomas tend to constrict the carotid lumen, unlike adenomas, which are thus mostly easily distinguishable [43, 57–59]. They often cause hyperostosis at the sites of bony attachment.

Craniopharyngioma

Craniopharyngioma is an important differential accounting for approximately 3% of all intracranial tumors. Usually slow growing, craniopharyngiomas arise from squamous epithelial cell rests of Rathke’ pouch [60]. Craniopharyngiomas may be divided into adamantinomatous or squamous-papillary histological types [61]. In children they make up about 5–10% of all intracranial neoplasms, being the third frequent tumor after medulloblastoma and astrocytoma. Patients usually present with headache, visual impairment, hydrocephalus, or hypopituitarism. Although suprasellar




Fig. 6.8 T1w + Gd (a + b + c). T2w fat-sat (d + e + f). Partly enhancing, T2w hyperintense mass of the optic chiasm, extending along the left optic nerve (a). Diffuse infiltration of the hypothalamus. Patient with neurofibromatosis I

in origin, about 50% extend into the sella. The suprasellar component of the tumor may induce edema to spread along the optic tract [62, 63]. In CT and MRI craniopharyngiomas have a typical heterogeneous appearance with cystic and solid components (Fig. 6.9) and frequently (approximately 93%) abundant calcifications [61, 64] (Table 6.2). The solid tumor portions as well as the cyst wall enhance usually after contrast administration.

Germ Cell Tumors

Intracranial germinomas occur most frequently in the pineal and suprasellar region but can also have a primary intrasellar origin [65, 66]. Occasionally, these tumors can extend and involve the sella turcica and mimic pituitary macroadenoma. Germinomas usually show prominent contrast

enhancement and present well-defined margins. Individual case reports describe sellar lesions that radiologically cannot be differentiated from pituitary adenomas, such as chondrosarcomas, granular cell tumors [67, 68], gangliocytomas [69, 70], fibrosarcomas [71, 72], hemangiopericytomas [73], esthesioneuroblastomas [74], melanomas [75], ependymomas [76], or lymphomas [77].

Schwannoma/Neurinoma

Nerve sheath tumors in the parasellar region are very rare lesions and usually arise from the trigeminal nerve (V1/V2) or the third, fourth, and sixth cranial nerve [78]. Neurinomas are slowly growing tumors. Expansion through the superior orbital fissure or the oval and round foramen has been described. On MRI they show an intense (usually heterogeneous) contrast enhancement [79, 80].




Fig. 6.9 Large cystic mass with hyperintense signal in T2w (**a**, coronal) and suprasellar epicenter. Thin, enhancing cyst wall (**b** and **c**, coronal and sagittal, T1w + Gd). Peripheral calcifications (**d**, axial CT). Obstructive hydrocephalus is present due to compression of the third ventricle and the mid-brain/aqueduct (**c**). Neuropathology revealed adamantinomatous craniopharyngioma

Neoplasms Involving the Clivus

Neoplasms in the clival region can include chordoma, chondrosarcoma, hemangiopericytoma [81], meningioma, lymphoma [82], plasmacytoma, paraganglioma [83], or metastasis. With an incidence below 1%, chordoma and chondrosarcoma are even rarer than malformative tumors [84]. Clival chordomas are slow-growing tumors arising from the remnants of the primitive notochord. It shows no age or sex predilection and usually causes bony destruction and can reach considerable size at the time of diagnosis. This extracranial tumor is heterogeneously hyperintense on T2-weighted images and shows a marked contrast enhancement. Chondroma is another bone-destructing, nodular/lobular tumor that tends to undergo mucinous, cystic regression and calcification. It most commonly arises from cartilaginous remnants in the area of the foramen lacerum. Imaging findings are similar to those in chordoma. Clival tumors tend to extend posteriorly into the prepontine cistern [85–88].

Neoplasms Involving the Sphenoidal Sinus

Mucoceleles are the most common space-occupying lesions of the sphenoid sinus. The pathogenesis relates to obstruction of the sphenoid ostium (secondary mucocele) or mucous retention cyst expansion (primary mucocele). In CT, a nondestructive mass, causing a thinning and bulging of the bony sinus walls, may be seen. The sellar part can mimic a para- or suprasellar mass. MRI better demonstrates the wall-enhancing fluid-filled lesion [89–91]. Rarely, inverted papillomas arising from the ethmoidal sinus can extend to the sphenoid sinus.

Table 6.2 Imaging features of craniopharyngiomas and macroadenomas

	Craniopharyngioma	Macroadenoma
Calcifications	90%	1–2%
Localization	Suprasellar (rarely purely intrasellar)	Intrasellar epicenter (may extend supra-/parasellar)

Vascular and Inflammatory Lesions of the Sella and the Hypothalamus

Aneurysms

Aneurysms of the sellar region usually originate from the cavernous or supraclinoid portions of the internal carotid artery and account for about up to 10% of all cerebral aneurysms. In selected cases they can mimic other supra-, para-, or intrasellar lesions [92]. MR angiography (time-of-flight or contrast enhanced) is useful in differentiating tumor from aneurysms (Fig. 6.10). Usually, the aneurysm itself can be sufficiently visualized by noninvasive imaging. These aneurysms may cause hypopituitarism or, in case of rupture, may also lead to (subarachnoid) hemorrhage in up to 15% [93].

Cavernous Sinus Thrombosis

Thrombosis of the cavernous sinus is a rare condition and often secondary to iatrogenic or septic etiologies [94, 95]. On MRI, enlargement of the cavernous sinus with internal filling defects and incomplete enhancement of the sinus may be noted, along with a diffusion restriction in the case of a (sub-)acute thrombus [96] (Fig. 6.11). Additionally, periorbital edema, exophthalmos, or dilatation of the superior ophthalmic vein can occur.

Inflammatory and Infectious Lesions

Sarcoidosis, a systemic disease featuring multiple noncaseating granulomas, may involve the

central nervous system in about 5% of cases. Predilection sites are the basal leptomeninges especially the sellar and suprasellar region, such as the pituitary stalk, optic chiasm, and the hypothalamus, respectively [97] (Fig. 6.12). Since radiological differentiation from other disease such as lymphocytic hypophysitis is often impossible, clinical history and time course are of great importance.

Lymphocytic hypophysitis is a rare autoimmune inflammatory disease most often seen in women in the peri- or postpartum period. Clinical symptoms usually include diabetes insipidus, amenorrhea, hypopituitarism, headache, and visual impairment favorably responding to steroid therapy [98–100]. MRI usually demonstrates thickening of the pituitary stalk in combination with an intense contrast enhancement [101].

Granulomatous hypophysitis can occur with fungal infections, tuberculosis [102], sarcoidosis [103], Langerhans cell histiocytosis, and Wegner's granulomatosis [104] and accounts for approximately 1% of sellar masses. The typical radiological appearance is similar to that of lymphocytic hypophysitis [105, 106].

Pituitary abscess can be primary or secondary due to an adenoma or to surgical procedure [107]. Spread of gram-positive bacteria from the sphenoid sinus is the usual source of infection, but rare infectious disease such as toxoplasmosis [108] or cryptococcosis has been described. Typical MRI findings are those of a round sellar mass with ring enhancement. Furthermore, meningeal enhancement due to concurrent meningitis may help distinguishing pituitary abscess from pituitary adenoma [109–111].




Fig. 6.10 Left infraorbital, saccular aneurysm of the ICA. Medially, the aneurysm reaches the pituitary body; cranially it slightly elevates the diaphragma sellae.

Coronal T2w (a) and fusion of ToF-angiography with 3D T2w CISS (b). VRT reconstruction of ToF-angiography showing the saccular aneurysm (c)




Fig. 6.11 Filling defect in the left cavernous sinus (a) and corresponding diffusion restriction of the acute thrombus (b). Thrombosis was presumably secondary due to sinusitis sphenoidalis (c)




Fig. 6.12 Neurosarcoidosis. Vivid contrast enhancement of the tuber cinereum (a and b, T1w + Gd coronal and sagittal). Distant leptomeningeal enhancement frontal (not shown). Skin biopsy revealed sarcoidosis




Fig. 6.13 Contrast enhancement extending to the right cavernous sinus with thickening of the adjacent lateral and medial rectus muscle (a and b, T1w + Gd). The inflamma-

tory changes have fully resolved after 6 months and therapy with steroids (c)

Tolosa-Hunt/Orbital Apex Syndrome

Tolosa-Hunt syndrome is a painful, recurrent ophthalmoplegia that usually responds favorably to steroid medication. It is essentially a diagnosis of exclusion. Associated with granulomatous inflammatory changes at the cavernous sinus and the superior orbital fissure, it can frequently strike various cerebral nerves located within this area [112]. Contrast-enhanced MRI shows an asymmetrical expansion and enhancement of the orbital apex extending to the cavernous sinus [113–115] (Fig. 6.13) and may also lead to thickening of the orbital muscles as well as cause secondary thrombosis of the cavernous sinus.

References

- Cox TD, Elster AD. Normal pituitary gland: changes in shape, size, and signal intensity during the 1st year of life at MR imaging. *Radiology*. 1991;179(3):721–4.
- Dietrich RB, Lis LE, Greensite FS, Pitt D. Normal MR appearance of the pituitary gland in the first 2 years of life. *AJNR Am J Neuroradiol*. 1995;16(7):1413–9.
- Elster AD. Modern imaging of the pituitary. *Radiology*. 1993;187(1):1–14.
- Fujisawa I, Nishimura K, Asato R, Togashi K, Itoh K, Noma S, et al. Posterior lobe of the pituitary in diabetes insipidus: MR findings. *J Comput Assist Tomogr*. 1987;11(2):221–5.
- Naylor MF, Scheithauer BW, Forbes GS, Tomlinson FH, Young WF. Rathke cleft cyst: CT, MR, and pathology of 23 cases. *J Comput Assist Tomogr*. 1995;19(6):853–9.
- Brassier G, Morandi X, Tayiar E, Riffaud L, Chabert E, Heresbach N, et al. Rathke's cleft cysts: surgical-MRI correlation in 16 symptomatic cases. *J Neuroradiol*. 1999;26(3):162–71.
- Harter LP, Silverberg GD, Brant-Zawadzki M. Intracellar arachnoid cyst: case report. *Neurosurgery*. 1980;7(4):387–90.
- Nomura M, Tachibana O, Hasegawa M, Kohda Y, Nakada M, Yamashita T, et al. Contrast-enhanced MRI of intrasellar arachnoid cysts: relationship between the pituitary gland and cyst. *Neuroradiology*. 1996;38(6):566–8.
- Starzyk J, Kwiatkowski S, Urbanowicz W, Starzyk B, Harasiewicz M, Kalicka-Kasperczyk A, et al. Suprasellar arachnoid cyst as a cause of precocious puberty—report of three patients and literature overview. *J Pediatr Endocrinol Metab*. 2003;16(3):447–55.
- Abrahams JJ, Trefelner E, Boulware SD. Idiopathic growth hormone deficiency: MR findings in 35 patients. *AJNR Am J Neuroradiol*. 1991;12(1):155–60.
- Argyropoulou M, Perignon F, Brauner R, Brunelle F. Magnetic resonance imaging in the diagnosis of growth hormone deficiency. *J Pediatr*. 1992;120(6):886–91.
- Hamilton J, Blaser S, Daneman D. MR imaging in idiopathic growth hormone deficiency. *AJNR Am J Neuroradiol*. 1998;19(9):1609–15.
- Kornreich L, Horev G, Lazar L, Schwarz M, Sulkes J, Pertzalan A. MR findings in growth hormone deficiency: correlation with severity of hypopituitarism. *AJNR Am J Neuroradiol*. 1998;19(8):1495–9.
- Fujisawa I, Kikuchi K, Nishimura K, Togashi K, Itoh K, Noma S, et al. Transection of the pituitary stalk: development of an ectopic posterior lobe assessed with MR imaging. *Radiology*. 1987;165(2):487–9.
- Kelly WM, Kucharczyk W, Kucharczyk J, Kjos B, Peck WW, Norman D, et al. Posterior pituitary ectopia: an MR feature of pituitary dwarfism. *AJNR Am J Neuroradiol*. 1988;9(3):453–60.
- Saeki N, Tokunaga H, Wagai N, Sunami K, Murai H, Kubota M, et al. MRI of ectopic posterior pituitary bright spot with large adenomas: appearances and relationship to transient postoperative diabetes insipidus. *Neuroradiology*. 2003;45(10):713–6.
- Civit T, Marchal JC, Pinelli C, Auque J, Hepner H. Intracellar epidermoid cysts. *Neurochirurgie*. 1999;45(2):150–4.

18. Yasargil MG, Abernathy CD, Sarioglu AC. Microneurosurgical treatment of intracranial dermoid and epidermoid tumors. *Neurosurgery*. 1989;24(4):561–7.
19. Judge DM, Kulin HE, Page R, Santen R, Trapukdi S. Hypothalamic hamartoma: a source of luteinizing-hormone-releasing factor in precocious puberty. *N Engl J Med*. 1977;296(1):7–10.
20. Amstutz DR, Coons SW, Kerrigan JF, Reke HL, Heiserman JE. Hypothalamic hamartomas: correlation of MR imaging and spectroscopic findings with tumor glial content. *AJNR Am J Neuroradiol*. 2006;27(4):794–8.
21. Freeman JL, Coleman LT, Wellard RM, Kean MJ, Rosenfeld JV, Jackson GD, et al. MR imaging and spectroscopic study of epileptogenic hypothalamic hamartomas: analysis of 72 cases. *AJNR Am J Neuroradiol*. 2004;25(3):450–62.
22. Colao A, Lombardi G. Growth-hormone and prolactin excess. *Lancet*. 1998;352(9138):1455–61.
23. Calle-Rodrigue RD, Giannini C, Scheithauer BW, Lloyd RV, Wollan PC, Kovacs KT, et al. Prolactinomas in male and female patients: a comparative clinicopathologic study. *Mayo Clin Proc*. 1998;73(11):1046–52.
24. Carter JN, Tyson JE, Tolis G, Van Vliet S, Faiman C, Friesen HG. Prolactin-screening tumors and hypogonadism in 22 men. *N Engl J Med*. 1978;299(16):847–52.
25. Melmed S. Acromegaly. *N Engl J Med*. 1990;322(14):966–77.
26. Holdaway IM, Rajasoorya CR, Gamble GD, Stewart AW. Long-term treatment outcome in acromegaly. *Growth Hormon IGF Res*. 2003;13(4):185–92.
27. Orme SM, McNally RJ, Cartwright RA, Belchetz PE. Mortality and cancer incidence in acromegaly: a retrospective cohort study. United Kingdom Acromegaly Study Group. *J Clin Endocrinol Metab*. 1998;83(8):2730–4.
28. Magiakou MA, Mastorakos G, Oldfield EH, Gomez MT, Doppman JL, Cutler GB Jr, et al. Cushing's syndrome in children and adolescents. Presentation, diagnosis, and therapy. *N Engl J Med*. 1994;331(10):629–36.
29. Sonino N, Boscaro M. Medical therapy for Cushing's disease. *Endocrinol Metab Clin N Am*. 1999;28(1):211–22.
30. Nelson DH. Cushing's syndrome--pituitary or adrenal origin? *J Chronic Dis*. 1960;12:499–503.
31. Nelson DH, Meakin JW, Thorn GW. ACTH-producing pituitary tumors following adrenalectomy for Cushing's syndrome. *Ann Intern Med*. 1960;52:560–9.
32. Beck-Peccoz P, Brucker-Davis F, Persani L, Smallridge RC, Weintraub BD. Thyrotropin-secreting pituitary tumors. *Endocr Rev*. 1996;17(6):610–38.
33. Ebersold MJ, Quast LM, Laws ER Jr, Scheithauer B, Randall RV. Long-term results in transsphenoidal removal of nonfunctioning pituitary adenomas. *J Neurosurg*. 1986;64(5):713–9.
34. Snyder PJ. Clinically nonfunctioning pituitary adenomas. *Endocrinol Metab Clin N Am*. 1993;22(1):163–75.
35. Aaberg TM Jr, Kay M, Sternau L. Metastatic tumors to the pituitary. *Am J Ophthalmol*. 1995;119(6):779–85.
36. Mayr NA, Yuh WT, Muhonen MG, Koci TM, Tali ET, Nguyen HD, et al. Pituitary metastases: MR findings. *J Comput Assist Tomogr*. 1993;17(3):432–7.
37. Kulkarni MV, Lee KF, McArdle CB, Yeakley JW, Haar FL. 1.5-T MR imaging of pituitary microadenomas: technical considerations and CT correlation. *AJNR Am J Neuroradiol*. 1988;9(1):5–11.
38. Bartynski WS, Lin L. Dynamic and conventional spin-echo MR of pituitary microlesions. *AJNR Am J Neuroradiol*. 1997;18(5):965–72.
39. Rand T, Lippitz P, Kink E, Huber H, Schneider B, Imhof H, et al. Evaluation of pituitary microadenomas with dynamic MR imaging. *Eur J Radiol*. 2002;41(2):131–5.
40. Stadnik T, Spruyt D, van Binst A, Luybaert R, d'Haens J, Osteaux M. Pituitary microadenomas: diagnosis with dynamic serial CT, conventional CT and T1-weighted MR imaging before and after injection of gadolinium. *Eur J Radiol*. 1994;18(3):191–8.
41. Tabarin A, Laurent F, Catargi B, Olivier-Puel F, Lescene R, Berge J, et al. Comparative evaluation of conventional and dynamic magnetic resonance imaging of the pituitary gland for the diagnosis of Cushing's disease. *Clin Endocrinol*. 1998;49(3):293–300.
42. Yuh WT, Fisher DJ, Nguyen HD, Tali ET, Gao F, Simonson TM, et al. Sequential MR enhancement pattern in normal pituitary gland and in pituitary adenoma. *AJNR Am J Neuroradiol*. 1994;15(1):101–8.
43. Donovan JL, Nesbit GM. Distinction of masses involving the sella and suprasellar space: specificity of imaging features. *AJR Am J Roentgenol*. 1996;167(3):597–603.
44. Kovacs K. Adenohypophysial necrosis in routine autopsies. *Endokrinologie*. 1972;60(3):309–16.
45. Rolih CA, Ober KP. Pituitary apoplexy. *Endocrinol Metab Clin N Am*. 1993;22(2):291–302.
46. Vidal E, Cevallos R, Vidal J, Ravon R, Moreau JJ, Rogues AM, et al. Twelve cases of pituitary apoplexy. *Arch Intern Med*. 1992;152(9):1893–9.
47. Bonicki W, Kasperlik-Zaluska A, Koszewski W, Zgliczynski W, Wislawski J. Pituitary apoplexy: endocrine, surgical and oncological emergency. Incidence, clinical course and treatment with reference to 799 cases of pituitary adenomas. *Acta Neurochir*. 1993;120(3–4):118–22.
48. Randeve HS, Schoebel J, Byrne J, Esiri M, Adams CB, Wass JA. Classical pituitary apoplexy: clinical features, management and outcome. *Clin Endocrinol*. 1999;51(2):181–8.
49. Abrams HL, Spiro R, Goldstein N. Metastases in carcinoma; analysis of 1000 autopsied cases. *Cancer*. 1950;3(1):74–85.
50. Pernicone PJ, Scheithauer BW, Sebo TJ, Kovacs KT, Horvath E, Young WF Jr, et al. Pituitary carci-

- noma: a clinicopathologic study of 15 cases. *Cancer*. 1997;79(4):804–12.
51. Scheithauer BW, Kurtkaya-Yapici O, Kovacs KT, Young WF Jr, Lloyd RV. Pituitary carcinoma: a clinicopathological review. *Neurosurgery*. 2005;56(5):1066–74; discussion -74.
 52. Dodge HW Jr, Love JG, Craig WM, Dockerty MB, Kearns TP, Holman CB, Hayles AB. Gliomas of the optic nerves. *AMA Arch Neurol Psychiatry*. 1958;79(6):607–21. <https://doi.org/10.1001/archneurpsyc.958.02340060003001>.
 53. Fujii K, Fujita N, Hirabuki N, Hashimoto T, Miura T, Kozuka T. Neuromas and meningiomas: evaluation of early enhancement with dynamic MR imaging. *AJNR Am J Neuroradiol*. 1992;13(4):1215–20.
 54. Joo YG, Korogi Y, Hirai T, Sakamoto Y, Sumi M, Takahashi M, et al. Differential diagnosis of extra-axial intracranial tumours by dynamic spin-echo MRI. *Neuroradiology*. 1995;37(7):522–5.
 55. Cappabianca P, Cirillo S, Alfieri A, D'Amico A, Maiuri F, Mariniello G, et al. Pituitary macroadenoma and diaphragma sellae meningioma: differential diagnosis on MRI. *Neuroradiology*. 1999;41(1):22–6.
 56. Pinzer T, Krishnan KG, Schackert G. The diaphragma sellae meningioma -- a rare differential diagnosis of non-functioning pituitary adenoma. *Zentralbl Neurochir*. 2004;65(4):195–7.
 57. Michael AS, Paige ML. MR imaging of intrasellar meningiomas simulating pituitary adenomas. *J Comput Assist Tomogr*. 1988;12(6):944–6.
 58. Nozaki K, Nagata I, Yoshida K, Kikuchi H. Intrasellar meningioma: case report and review of the literature. *Surg Neurol*. 1997;47(5):447–52; discussion 52–4.
 59. Yeakley JW, Kulkarni MV, McArdle CB, Haar FL, Tang RA. High-resolution MR imaging of juxtaseellar meningiomas with CT and angiographic correlation. *AJNR Am J Neuroradiol*. 1988;9(2):279–85.
 60. Hoffman HJ. Surgical management of craniopharyngioma. *Pediatr Neurosurg*. 1994;21(Suppl 1):44–9.
 61. Sartoretti-Schefer S, Wichmann W, Aguzzi A, Valavanis A. MR differentiation of adamantinous and squamous-papillary craniopharyngiomas. *AJNR Am J Neuroradiol*. 1997;18(1):77–87.
 62. Nagahata M, Hosoya T, Kayama T, Yamaguchi K. Edema along the optic tract: a useful MR finding for the diagnosis of craniopharyngiomas. *AJNR Am J Neuroradiol*. 1998;19(9):1753–7.
 63. Saeki N, Uchino Y, Murai H, Kubota M, Isobe K, Uno T, et al. MR imaging study of edema-like change along the optic tract in patients with pituitary region tumors. *AJNR Am J Neuroradiol*. 2003;24(3):336–42.
 64. Eldevik OP, Blaiwas M, Gabrielsen TO, Hald JK, Chandler WF. Craniopharyngioma: radiologic and histologic findings and recurrence. *AJNR Am J Neuroradiol*. 1996;17(8):1427–39.
 65. Cho DY, Wang YC, Ho WL. Primary intrasellar mixed germ-cell tumor with precocious puberty and diabetes insipidus. *Childs Nerv Syst*. 1997;13(1):42–6.
 66. Jennings MT, Gelman R, Hochberg F. Intracranial germ-cell tumors: natural history and pathogenesis. *J Neurosurg*. 1985;63(2):155–67.
 67. Bubl R, Hugo HH, Hempelmann RG, Barth H, Mehdorn HM. Granular-cell tumour: a rare suprasellar mass. *Neuroradiology*. 2001;43(4):309–12.
 68. Wilkinson MD, Fulham MJ, Besser M. Neuroimaging findings in a suprasellar granular cell tumor. *J Comput Assist Tomogr*. 2003;27(1):26–9.
 69. Isidro ML, Iglesias Diaz P, Matias-Guiu X, Cordido F. Acromegaly due to a growth hormone-releasing hormone-secreting intracranial gangliocytoma. *J Endocrinol Investig*. 2005;28(2):162–5.
 70. Song JK, Weil RJ. Pathologic quiz case: an unusual cause of acromegaly. Intrasellar gangliocytoma. *Arch Pathol Lab Med*. 2005;129(3):415–6.
 71. Lopes MB, Lanzino G, Cloft HJ, Winston DC, Vance ML, Laws ER Jr. Primary fibrosarcoma of the sella unrelated to previous radiation therapy. *Mod Pathol*. 1998;11(6):579–84.
 72. Moro M, Giannini C, Scheithauer BW, Lloyd RV, Restall P, Eagleton C, et al. Combined sellar fibrosarcoma and prolactinoma with neuronal metaplasia: report of a case unassociated with radiotherapy. *Endocr Pathol*. 2004;15(2):149–58.
 73. Gharbi A, Ousehal A, Kissani N, Lakhdar A, Azhari A, Kadir R. Sellar hemangiopericytoma. Report of a case. *J Neurosurg*. 2001;28(3):195–9.
 74. Sarwar M. Primary sellar-parasellar esthesioneuroblastoma. *AJR Am J Roentgenol*. 1979;133(1):140–1.
 75. Jacob S, Pye E, Hbabih M, Messios N, Rajabally YA. Rapidly progressive bilateral ophthalmoplegia and enlarging sellar mass caused by amelanotic melanoma. *J Neuroophthalmol*. 2006;26(1):49–50.
 76. Kimura M, Takayasu M, Suzuki Y, Negoro M, Nagasaka T, Nakashima N, et al. Primary choroid plexus papilloma located in the suprasellar region: case report. *Neurosurgery*. 1992;31(3):563–6.
 77. Huang YY, Lin SF, Dunn P, Wai YY, Hsueh C, Tsai JS. Primary pituitary lymphoma presenting as hypophysitis. *Endocr J*. 2005;52(5):543–9.
 78. Wilberger JE Jr. Primary intrasellar schwannoma: case report. *Surg Neurol*. 1989;32(2):156–8.
 79. Johnsen DE, Woodruff WW, Allen IS, Cera PJ, Funkhouser GR, Coleman LL. MR imaging of the sellar and juxtaseellar regions. *Radiographics*. 1991;11(5):727–58.
 80. Whee SM, Lee JI, Kim JH. Intrasellar schwannoma mimicking pituitary adenoma: a case report. *J Korean Med Sci*. 2002;17(1):147–50.
 81. Morrison DA, Bibby K. Sellar and suprasellar hemangiopericytoma mimicking pituitary adenoma. *Arch Ophthalmol*. 1997;115(9):1201–3.
 82. Landman RE, Wardlaw SL, McConnell RJ, Khandji AG, Bruce JN, Freda PU. Pituitary lymphoma presenting as fever of unknown origin. *J Clin Endocrinol Metab*. 2001;86(4):1470–6.

83. Naggara O, Varlet P, Page P, Oppenheim C, Meder JF. Suprasellar paraganglioma: a case report and review of the literature. *Neuroradiology*. 2005;47(10):753–7.
84. Freda PU, Post KD. Differential diagnosis of sellar masses. *Endocrinol Metab Clin N Am*. 1999;28(1):81–117, vi.
85. Allan CA, Kaltsas G, Evanson J, Geddes J, Lowe DG, Plowman PN, et al. Pituitary chondrosarcoma: an unusual cause of a sellar mass presenting as a pituitary adenoma. *J Clin Endocrinol Metab*. 2001;86(1):386–91.
86. Gay E, Sekhar LN, Rubinstein E, Wright DC, Sen C, Janecka IP, et al. Chordomas and chondrosarcomas of the cranial base: results and follow-up of 60 patients. *Neurosurgery*. 1995;36(5):887–96; discussion 96–7.
87. Lee HJ, Kalnin AJ, Holodny AI, Schulder M, Grigorian A, Sharer LR. Hemorrhagic chondroid chordoma mimicking pituitary apoplexy. *Neuroradiology*. 1998;40(11):720–3.
88. Rosenberg AE, Nielsen GP, Keel SB, Renard LG, Fitzek MM, Munzenrider JE, et al. Chondrosarcoma of the base of the skull: a clinicopathologic study of 200 cases with emphasis on its distinction from chordoma. *Am J Surg Pathol*. 1999;23(11):1370–8.
89. Akan H, Cihan B, Celenk C. Sphenoid sinus mucocele causing third nerve paralysis: CT and MR findings. *Dentomaxillofac Radiol*. 2004;33(5):342–4.
90. Di Girolamo S, Cannizzaro P, Picciotti P, Nardi C. Ophthalmoplegia and ptosis as onset symptoms of an isolated primary mucocele of the sphenoid sinus. *J Oral Maxillofac Surg*. 2002;60(12):1500–2.
91. Som PM. CT of the paranasal sinuses. *Neuroradiology*. 1985;27(3):189–201.
92. Atlas SW, Grossman RI, Goldberg HI, Hackney DB, Bilaniuk LT, Zimmerman RA. Partially thrombosed giant intracranial aneurysms: correlation of MR and pathologic findings. *Radiology*. 1987;162(1 Pt 1):11–4.
93. Hanak BW, Zada G, Nayar VV, Thies R, Du R, Day AL, Laws ER. Cerebral aneurysms with intrasellar extension: a systematic review of clinical, anatomical, and treatment characteristics. *J Neurosurg*. 2012;116(1):164–78. <https://doi.org/10.3171/2011.9.JNS.11380>. Epub 2011 Nov 4.
94. DiNubile MJ. Septic thrombosis of the cavernous sinuses. *Arch Neurol*. 1988;45(5):567–72.
95. Ebright JR, Pace MT, Niazi AF. Septic thrombosis of the cavernous sinuses. *Arch Intern Med*. 2001;161(22):2671–6.
96. Schuknecht B, Simmen D, Yuksel C, Valavanis A. Tributary venous occlusion and septic cavernous sinus thrombosis: CT and MR findings. *AJNR Am J Neuroradiol*. 1998;19(4):617–26.
97. Guoth MS, Kim J, de Lotbiniere AC, Brines ML. Neurosarcoidosis presenting as hypopituitarism and a cystic pituitary mass. *Am J Med Sci*. 1998;315(3):220–4.
98. Bellastella A, Bizzarro A, Coronella C, Bellastella G, Sinisi AA, De Bellis A. Lymphocytic hypophysitis: a rare or underestimated disease? *Eur J Endocrinol*. 2003;149(5):363–76.
99. Leung GK, Lopes MB, Thorner MO, Vance ML, Laws ER Jr. Primary hypophysitis: a single-center experience in 16 cases. *J Neurosurg*. 2004;101(2):262–71.
100. Thodou E, Asa SL, Kontogeorgos G, Kovacs K, Horvath E, Ezzat S. Clinical case seminar: lymphocytic hypophysitis: clinicopathological findings. *J Clin Endocrinol Metab*. 1995;80(8):2302–11.
101. Saiwai S, Inoue Y, Ishihara T, Matsumoto S, Nemoto Y, Tashiro T, et al. Lymphocytic adenohypophysitis: skull radiographs and MRI. *Neuroradiology*. 1998;40(2):114–20.
102. Basaria S, Ayala AR, Guerin C, Dobs AS. A rare pituitary lesion. *J Endocrinol Investig*. 2000;23(3):189–92.
103. Bullmann C, Faust M, Hoffmann A, Heppner C, Jockenhovel F, Muller-Wieland D, et al. Five cases with central diabetes insipidus and hypogonadism as first presentation of neurosarcoidosis. *Eur J Endocrinol*. 2000;142(4):365–72.
104. Goyal M, Kucharczyk W, Keystone E. Granulomatous hypophysitis due to Wegener's granulomatosis. *AJNR Am J Neuroradiol*. 2000;21(8):1466–9.
105. Honegger J, Fahlbusch R, Bornemann A, Hensen J, Buchfelder M, Muller M, et al. Lymphocytic and granulomatous hypophysitis: experience with nine cases. *Neurosurgery*. 1997;40(4):713–22; discussion 22–3.
106. Sato N, Sze G, Endo K. Hypophysitis: endocrinologic and dynamic MR findings. *AJNR Am J Neuroradiol*. 1998;19(3):439–44.
107. Jadhav RN, Dahiwadkar HV, Palande DA. Abscess formation in invasive pituitary adenoma: case report. *Neurosurgery*. 1998;43(3):616–9.
108. Milligan SA, Katz MS, Craven PC, Strandberg DA, Russell IJ, Becker RA. Toxoplasmosis presenting as panhypopituitarism in a patient with the acquired immune deficiency syndrome. *Am J Med*. 1984;77(4):760–4.
109. Maartens NF, Ellegala DB, Lopes MB. Pituitary abscess. *J Neurosurg*. 2001;95(6):1110–2.
110. Sabbah P, Bonardel G, Herve R, Marjou F, Hor F, Pharaboz C, et al. CT and MRI findings in primitive pituitary abscess: a case report and review of literature. *J Neuroradiol*. 1999;26(3):196–9.
111. Vates GE, Berger MS, Wilson CB. Diagnosis and management of pituitary abscess: a review of twenty-four cases. *J Neurosurg*. 2001;95(2):233–41.
112. Kline LB, Hoyt WF. The Tolosa-Hunt syndrome. *J Neurol Neurosurg Psychiatry*. 2001;71(5):577–82.
113. de Arcaya AA, Cerezal L, Canga A, Polo JM, Berciano J, Pascual J. Neuroimaging diagnosis of Tolosa-Hunt syndrome: MRI contribution. *Headache*. 1999;39(5):321–5.
114. Haque TL, Miki Y, Kashii S, Yamamoto A, Kanagaki M, Takahashi T, et al. Dynamic MR imaging in Tolosa-Hunt syndrome. *Eur J Radiol*. 2004;51(3):209–17.
115. Odabasi Z, Gokcil Z, Atilla S, Pabuscu Y, Vural O, Yardim M. The value of MRI in a case of Tolosa-Hunt syndrome. *Clin Neurol Neurosurg*. 1997;99(2):151–4.

Neurological Syndromes of the Hypothalamus

7

Christopher Morgan Smith, Rima El-Abassi,
and David Chachkhiani

Introduction

The hypothalamus is a small, but crucial and unique, structure of the brain that has many functions, including homeostatic mechanisms (i.e., hunger, thirst, circadian rhythms), endocrine hormone production, autonomic control, and limbic regulation. One can recall its function with the mnemonic “HEAL” (Homeostasis, Endocrine, Autonomic, Limbic) [1]. However, in this chapter, rather than its function, we will be discussing the *dysfunction* of the hypothalamus and many of the consequential clinical syndromes of hypothalamic dysregulation.

Recall from Part I, Chap. 4, that there are three major regions of the hypothalamus – supraoptic, tuberal, and mamillary regions. Despite the very small size of the hypothalamus, these regions encompass many key nuclei including the paraventricular nucleus, supraoptic nucleus, dorsomedial nucleus, posterior nucleus, anterior nucleus, lateral nucleus, mammillary body, arcuate nucleus, suprachiasmatic nucleus, ventromedial nucleus, and preoptic nucleus. It is important to take note of key hypothalamic nuclei responsibilities to best understand their dysfunction. As we will see, any insult to these cramped regions

and nuclei can produce profound neurological changes.

Neuro-structural Syndromes – *Neurovascular Ischemia/Infarct/ Trauma*

The etiology of structural insults to the CNS can vary including neurovascular lesions, vasculopathies, trauma, inflammatory processes, infectious processes, tumors/malignancy, electrolyte/metabolic disturbances, congenital defects, and neurodegenerative disorders. One of the most common neurological insults is neurovascular lesions, or stroke.

The arterial blood supply of the hypothalamus largely comes from the branches of the circle of Willis. Almost every nuclear group within the hypothalamus receives its arterial supply from multiple blood vessels and/or branches of local vessels. The ventral supraoptic region obtains its blood supply from the tubero-optic branch of the internal carotid artery, the dorsal aspect of the supraoptic region gets its blood supply from the branches of the anterior cerebral artery, while the preoptic areas receive its blood supply via the anterior communicating artery. The paraventricular, lateral, and dorsomedial nuclear regions obtain its supply from the posterior communicating artery. The lateral and median tuberal nuclei are supplied by the cir-

C. M. Smith (✉) · R. El-Abassi · D. Chachkhiani
Department of Neurology, Louisiana State University,
Health Sciences Center, New Orleans, LA, USA
e-mail: Csmi43@lsuhsc.edu; dchach@lsuhsc.edu

Table 7.1 Hypothalamic nuclei and correlating functions

Hypothalamic nuclei	Function
Preoptic n.	Thermoregulation
Suprachiasmatic n.	Circadian rhythms; sleep
Anterior n.	Thermoregulation; reduction in temperature
Paraventricular n.	Oxytocin release, ADH release
Supraoptic n.	Thirst, ADH release
Periventricular n.	Autonomic regulation, metabolism, growth, reproduction
Arcuate n.	Appetite regulation
Ventromedial n.	Appetite regulation, reproduction
Posterior hypothalamic n.	Autonomic regulation, thermoregulation
Mammillary n.	Memory
Lateral hypothalamic n.	Hunger, thirst, arousal

cuminfundibular plexus. Finally, the mammillary bodies are supplied by the posterior cerebral arteries distal to the bifurcation of the basilar artery as well as the posterior communicating arteries. Interestingly, the vessels that supply the deepest nuclei enter more superficially [2].

Figure 7.1 demonstrates many of the arteries responsible for supplying the hypothalamus [3]. Any insult to the aforementioned arteries can give rise to hypoperfusion and ischemia to its area of blood supply. As many of these vessels are notably small vessels, a single lesion may not be enough to infarct the correlating hypothalamic tissues; however, a more proximal lesion can produce devastating effects.

In review of some of the important hypothalamic regions and nuclei (Table 7.1 demonstrates major hypothalamic nuclei and correlating function), remember that the anterior and posterior hypothalami are important in autonomic functions such as temperature regulation. The posterior hypothalamus is responsible in increasing body temperature when the body temperature is reduced. As the anterior region (preoptic n.) aids in reducing an overheated body temperature and the posterior hypothalamus (posterior n.) is useful in the hyperthermic effects, an imbalance between the two opposing forces can yield an abnormal baseline body temperature. Considering the anterior communicating artery supplies the




Fig. 7.1 Circle of Willis. Labeled are some of the arteries responsible for supplying the hypothalamus. Anterior cerebral arteries (A1)(A2); posterior cerebral arteries (P1) (P2) near their origin; posterior communicating artery (PcoA); and anterior communicating artery (AcoA) [3]

preoptic area, obstruction in blood flow through the anterior communicating artery can lead to infarct, which would produce an overall hyperthermic effect.

Demyelinating Disorders: Multiple Sclerosis and Neuromyelitis Optica

One of the more well-known neurological demyelinating disorders is multiple sclerosis. Multiple sclerosis is defined by the American Academy of Neurology as a chronic disorder characterized by inflammation and demyelination of the central nervous system associated with variable degrees of axonal and neuronal damage [4]. Multiple sclerosis classically affects CNS white matter, or neuronal fiber bundles. Demyelination of these fibers within or near the hypothalamus can produce autonomic and endocrine abnormalities and

mood disorders in individuals living with multiple sclerosis. The hypothalamus, however, frequently appears normal on neuroimaging, despite clinical evidence of hypothalamic involvement and confirmation on necropsy studies [5].

Clinically, hypothalamic dysfunction in multiple sclerosis can present similarly to those with neurovascular or structural lesions. Thus, it is important to note that autonomic features and sleep disturbances are common among multiple sclerosis patients. Parenteral glucocorticoids (i.e., methylprednisolone) may be considered the drug of choice in patients presenting with acute demyelinating lesions in multiple sclerosis, and the hypothalamo-pituitary-adrenal axis (HPA) is crucial in the production of intrinsic glucocorticoids. Demyelinating lesions to the HPA may impede innate steroidal production and theoretically increase the susceptibility and severity of future multiple sclerosis flares. When compared to controls, few studies have shown that patients living with multiple sclerosis showed significantly higher neuronal activation in production of corticotropin-releasing hormone (CRH) [6].

Neuromyelitis optica spectrum disorder (NMOSD), previously known as Devic’s disease, is a similar process to multiple sclerosis in that it is an inflammatory demyelinating disease of the central nervous system and can easily be mistaken for multiple sclerosis (Table 7.2 demonstrates major differences between NMOSD and MS). NMOSD is an aggressive demyelinating process associated with episodic optic neuritis, transverse myelitis, and serum biomarkers: aquaporin-4 (AQP4)-IgG and myelin oligodendrocyte glycoprotein (MOG)-IgG. The transverse myelitis seen in NMOSD is typically more longitudinally extensive involving 3 or more vertebral segments of the spinal cord. Episodic intractable nausea, vomiting, or hiccups are also a feature of NMO with area postrema irritation or involvement [4].

Prior to the discovery of the AQP4-Ab in 2004, Devic’s disease was largely considered to have an absence of brain involvement. However, with advancing studies, certain discoveries may prove otherwise. In fact, hypothalamic lesions may be more prevalent in neuromyelitis optica

Table 7.2 Clinical comparison between neuromyelitis optica and multiple sclerosis


	Neuromyelitis optica (Devic’s disease)	Multiple sclerosis
AVG. age of onset	40s	30s
Brain involvement	Brain MRI is typically nonspecific	Multiple periventricular white matter lesions (Dawson’s fingers)
Optic nerve involvement	Mostly unilateral, however can be bilateral as well; outcome is typically poor	Mostly unilateral with favorable recovery
Spinal cord involvement	Longitudinally extending (>3 vertebral segments)	Smaller lesions
Associated antibody	Associated with serum NMO antibody	Associated with oligoclonal bands in CSF
Overall impairment	Functional impairment after attack is usually severe	Usually less severe outcome

than multiple sclerosis; however, this claim is under study [7]. AQP4-Ab has a sensitivity of ~60% and a specificity of 98% on tissue-based assay [8]. Patients who have AQP4-Ab positive NMOSD with hypothalamic lesions had symptoms of ataxia, intractable hiccups/nausea, SIADH, and encephalopathy, which were more common than those without hypothalamic lesions.

Limbic Encephalitis

The limbic system is a collection of brain structures that aid in many functions including homeostasis, olfaction, memory, and emotion (mnemonic – “HOME”) [1]. Interestingly, these structures are thought to be resultant of evolutionary change as they may have played an important role in survival. Key structures of the limbic system include the hypothalamus, hippocampus, amygdala, thalamus, septal nuclei, olfactory cortex/bulb, limbic cortex, and basal

Fig. 7.2 Limbic structures and cortical pathways. (Source: Ropper AH, Samuels MA, Klein JP: *Adams and Victor's Principles of Neurology, Tenth Edition*: www.accessmedicine.com. Copyright © The McGraw-Hill Companies, Inc. All rights reserved)



Copyright The McGraw-Hill Companies, Inc. All rights reserved.

ganglia. Figure 7.2 depicts a visual representation of many of the limbic system structures [9].

Limbic encephalitis is a neurological disorder in which the limbic system is affected by inflammation and consequently dysfunction. Limbic encephalitis classically presents as short-term memory loss, agitation, cognitive impairment, and seizures. Involvement of the hypothalamus may also include presenting symptoms of sleep disturbances, fatigue, narcolepsy, hyperthermia, weight gain, sexual dysfunction, autonomic dysfunction, and hormonal dysfunction of the hypothalamic-pituitary-adrenal (HPA) axis. Most

causes of inflammation to the limbic system include infectious or autoimmune etiology. There is insufficient data to suggest a prevalence or incidence rate of limbic encephalitis.

Autoimmune forms of limbic encephalitis are further categorized into paraneoplastic limbic encephalitis and non-paraneoplastic limbic encephalitis. Between the two subsets of autoimmune encephalitis, non-paraneoplastic limbic encephalitis is much more common [10]. Non-paraneoplastic limbic encephalitis is caused by circulating antibodies recognizing self-antigens located on or near the limbic system.

Table 7.3 Known antigens associated with autoimmune encephalitis

Intracellular antigens	Neuronal-surface proteins
GAD 65 – glutamic acid decarboxylase	Caspr2 – contactin-associated protein 2
CRMP5 – collapsin response mediator protein 5	LGI-1 – leucine-rich glioma-inactivated 1
Ma1/2 – Ma proteins widely throughout a normal brain	DDPX – dipeptidyl-peptidase-like protein 6
DNER – Delta/Notch-like epidermal growth factor	GFAP – glial fibrillary acidic protein
	Neurexin-3-alpha – synapse membrane proteins
	IgLON5 – a cell surface adhesion molecule

Potential threatening antibodies have been identified to target antigens that play a crucial role in signal transmission in the central nervous system. Table 7.3 details some of the major intracellular antigens and proteins involved in the etiopathogenesis of this form of encephalitis. Intracellular antigens (GAD54, CRMP5, Ma1, Ma2, DNER), extracellular neuronal cell-surface proteins (Caspr2, LGI-1, DDPX, GFAP, neurexin-3-alpha, IgLON5) and receptors (NMDAR, AMPAR, GABAbR, GABAaR, mGluR), and others (MOG, AQP4).

Most non-paraneoplastic limbic encephalitides result from antibodies attacking cell surface proteins or receptors, instead of the intracellular foci. Some of the well-established foci include LGI1, CASPR2, AMPA receptor, GABA(A/B) receptor, AMPA receptor, and NMDA receptor. Receptor antibodies can alter receptor functioning and receptor density in synapses [11].

Table 7.4 details the major autoantibodies and their associated clinical presenting symptomatology and signs in the spectrum of autoimmune encephalitis.

Voltage-gated potassium channels (VGKC) were once believed to be a major antigen associated with limbic encephalitis. The VGKC complex is the combination of LGI1 and CASPR-2 with voltage-gated potassium channels. Patients with LGI1-antibody positivity are more likely to have features of limbic encephalitis and seizures (Table 7.3 shows autoimmune encephalitis anti-

Table 7.4 Autoimmune encephalitis antibodies with associated clinical features

Antibody	Associated clinical features
Anti-NMDA	Memory loss, psychosis, seizures, insomnia, dyskinesias
Anti-Ma2	LE, daytime sleepiness, ophthalmoplegia
Anti-Hu	Sensory neuropathy, LE, cerebellar ataxia
Anti-VGCC	Cerebellar ataxia
Anti-LGI1	LE, faciobrachial dystonic seizures, different epileptic seizures
Anti-GABA-A	Refractory seizures, status epilepticus, cognitive and behavioral changes
Anti-GABA-B	LE, seizures, opsoclonus-mycoclonus
Anti-Caspr2	LE, cerebellar ataxia, neuromyotonia
Anti-AMPA	LE, bipolar features, seizures
Anti-IgLON5	Parasomnias, chorea

LE limbic encephalitis

bodies associated with clinical features), while patients with CASPR2-antibody positivity were more likely to experience symptoms of limbic encephalitis with neuromyotonia, dysautonomia, and neuropathic pains.

The N-methyl-D-aspartate (NMDA) receptor is a glutamate receptor that allows cations (positively charged ions) to flow through the neuronal cell membrane allowing for depolarization. The NMDA receptor aids in synaptic receptor down-/upregulation and memory. Anti-NMDA receptor encephalitis has a higher prevalence in women compared to men but can be seen in children as well [12]. Clinical symptoms may include hallucinations (auditory/visual), agitation, impaired consciousness, seizures, motor dysfunction, and autonomic dysfunction. Anti-NMDA receptor antibodies can also be associated with psychiatric conditions like schizophrenia [12], and caution should be used when considering a diagnosis of NMDA-receptor encephalitis. The α -amino-3-hydroxy-5-methylisoxazolepropionic acid (AMPA) receptor is an ionotropic transmembrane receptor that, when activated, induces NMDA receptor activity. Antibodies against the AMPA receptor typically target the glutamate receptor (GluR1, GluR2) subunits. The AMPA receptors are also expressed widely throughout the CNS, including the hypothalamus.

Glucagon-like peptide-1 (GLP-1)-expressing neurons in the hindbrain have been found to project to the paraventricular nucleus of the hypothalamus and aid in the regulation of food consumption by enhancing AMPA receptor expression. Anti-AMPA encephalitis is a paraneoplastic process in 64% of cases, and malignancy screening should be considered. The typical presenting features are limbic encephalitis or psychosis, and the prognosis and treatment response largely depend on detection of tumors and onconeural antibodies [13].

Another important paraneoplastic limbic encephalitis to mention in special regard to the hypothalamus is anti-Ma2 encephalitis. Anti-Ma2 limbic encephalitis differs from the classical presentation of paraneoplastic limbic encephalitis. This subtype of paraneoplastic encephalitis often presents with diencephalic or brainstem dysfunction. In one study of 34 patients with confirmed anti-Ma encephalitis, excessive daytime sleepiness was seen in 32% of patients, and 34% had additional abnormalities seen in hormonal testing or MRI findings that indicated hypothalamic involvement [14]. Notably, in both anti-Ma1 and anti-Ma2 encephalitis, tumors can be an underlying factor and must be screened for particularly of the testes. Immunosuppressive therapy and oncological treatments have been shown to improve outcomes [14].

The first step in approaching limbic encephalitis is its diagnosis and malignancy screening. Treatment and prognosis of autoimmune limbic encephalitis vary largely on the time of diagnosis, type of antibody, comorbidities, and associated malignancies. Diagnosis can be very challenging as patients often present with nonspecific signs and symptoms. Treatment options can include supportive therapy, glucocorticoids, IVIG, plasma exchange, or immunomodulatory therapies (i.e., azathioprine, mycophenolate, etc.).

Migraine


Migraine is a neurological disorder that most commonly presents as recurrent severe headaches. There are classically four main phases of migraine – migraine aura, premonitory phase,

headache phase, and the postdrome [15]. The International Classification of Headache Disorders, Third Edition (ICHD-3), defines migraine aura as recurrent attacks, lasting minutes, of unilateral fully reversible visual, sensory, or other central nervous system symptoms that usually develop gradually and are usually followed by headache and associated migraine symptoms. Migraines can be with or without aura. About 20–40% of patients with migraines report some form of aura. Roughly 90% of migraine auras are reportedly visual symptoms, and ~10% are sensory, language, and/or motor symptoms (usually in addition to visual symptoms) [15]. The typical duration of migraine aura is 30 minutes; however, it can last anywhere from a few minutes to 4 hours. The etiology of migraine auras is currently unknown, but interestingly, one theory is that visual auras are a consequence of a higher neuron-to-astrocyte ratio in the visual cortex [15]. Figure 7.3 offers another theory of migraine etiology by means of electrophysiological hyperexcitability extending to nerves and glial cells called *cortical spreading depression* [15].

The premonitory, or prodromal, phase is newly defined as a period of nonpainful symptoms which precede the onset of migraine headache usually by hours to a few days. The most common symptoms of the premonitory phase include fatigue, mood changes, and yawning; however, symptoms can largely be categorized as cognitive changes, homeostatic alterations, and sensory sensitivities. Following the premonitory phase, the headache phase of migraines is typically a severe, unilateral headache with associated nausea, vomiting, photophobia, and phonophobia lasting 5–72 hours. Finally, the postdrome phase is the time between resolution of migrainous headache and at which point the patient feels he/she is back to baseline. Similar to the premonitory phase, symptoms constitute non-headache symptoms, most commonly fatigue, lack of concentration, and neck stiffness [16].

One study in 2016 showed evidence that neuropeptide Y may play as a key component in migraines and trigeminovascular activation [16]. Neuropeptide Y is a naturally secreted substance by the hypothalamus that is involved in various

Fig. 7.3 Current understanding of the pathophysiologic mechanisms involved in migraine. Cortical spreading depression is a phenomenon theorized in migraine etiology, which is described as an extending electrophysiological hyperexcitability of nerves and glial cells that propagates alterations in the blood-brain barrier permeability and activation of trigeminal afferent sensory fibers



processes such as appetite regulation, pain, and circadian rhythms. The authors of this study administered neuropeptide Y systemically and noted inhibition of the trigeminocervical complex. Additionally, this study proposed the possible hypothalamic role in the association of migraine with impaired appetite and serum glucose levels [16]. Other studies suggest migrainous hypothalamic roles in sleep/mood disturbances and emotionality [17, 18]. Furthermore, there has been some evidence supporting that stimulation of the A11 nucleus of the hypothalamus lessened nociception by diminishing trigeminocervical complex signals [19]. As previously mentioned, one of the most common reported premonitory phase symptoms is yawning. Yawning is thought to be partly dopamine-mediated as it can be triggered by dopamine-agonists and countered by hypophysectomy. Somatostatin, another hypothalamic hormone, may also play a significant role in primary headache disorders [20].

Hypothalamic involvement in migraines has also been noted radiographically. Maniyan, FH., et al. demonstrated early hypothalamic activation on positron emission tomography (PET) scan during the premonitory phase of nitroglycerin-triggered migraine attacks [21]. Other brain structures included the dorsolateral pons and

multiple cortical areas as well. These findings may explain some of the migraine premonitory phase symptomatology and alterations in homeostasis. Functional MRI (fMRI) studies have shown some evidence of hypothalamic involvement in migraines. In one study [22] an individual with a history of migraines was followed by daily fMRIs for a month. The authors found that hypothalamic activity was altered in the 24 hours preceding the headache onset as well as alterations in the functional signaling between the hypothalamus, spinal trigeminal nuclei, and dorsal rostral pons during the headache phase of migraine. These results are also consistent with hypothalamic involvement of the headache phase in migraines.

Trigeminal Autonomic Cephalgias and Cluster Headache

Trigeminal autonomic cephalgias (TACs) are a group of primary headache disorders that are typically unilateral and are associated with ipsilateral autonomic features such as lacrimation, conjunctival injection, and nasal congestion. This group of primary headache disorders includes (1) cluster headache, (2) short-lasting unilateral neuralgiform headache with conjunctival injection

Table 7.5 Trigeminal autonomic cephalgias

Diagnosis	Features	Duration	Common trigger	Treatment
CH	Periorbital, excruciating pain with autonomic and migrainous features	30 to 90 minutes	Alcohol	Sumatriptan, oxygen
SUNCT	Orbital sharp pain with autonomic features	5 seconds to 6 minutes	Touch	IV lidocaine
SUNA	Orbital sharp pain with autonomic features	15 seconds to 3 minutes	Touch	IV lidocaine
PH	Orbital stabbing pain with autonomic and migrainous features	2 to 30 minutes	Touch	Indomethacin
HC	Mild to severe throbbing pain which is continuous and has exacerbations; can be periorbital, temporal, and/or frontal	>3 months	Food/stress	Indomethacin

TACs trigeminal autonomic cephalgias, CH cluster headache, SUNCT short-lasting unilateral neuralgiform headache with conjunctival injection and tearing, SUNA short-lasting unilateral neuralgiform headache attacks with cranial autonomic symptoms, PH paroxysmal hemicrania, HC hemicrania continua

and tearing (SUNCT), (3) short-lasting unilateral neuralgiform headache with cranial autonomic symptoms (SUNA), (4) paroxysmal hemicrania, and (5) hemicrania continua (Table 7.5 shows major TACs and their characteristic features). While the symptoms of these cephalgias are similar in headache with autonomic features, they differ vastly in duration and treatment [23].

Cluster headaches are more specifically characterized as a unilateral headache disorder with ipsilateral autonomic features and/or restlessness. Cluster headaches are more common than other trigeminal autonomic cephalgias, and interestingly, they are often more severe. There is a higher incidence of attacks in men when compared to women. The usual age of onset is between 20 and 40 years old. Cluster headaches can be episodic in nature (90% of affected individuals), which is defined as having headache-free periods for greater than 3 months, or chronic, that is, having attacks frequently within a 3-month timeframe. Interestingly, one distinguishing feature of cluster headaches compared to other TACs is a clocklike pattern and circannual periodicity of attacks. For example, an affected individual may report having daily headaches for the month of May at a specific time of day and then experience years of resolution until he/she experiences another bout of headaches during May at the same time of day. It is noted that this clocklike pattern is present in 82% of individuals affected by cluster headaches but not classically seen in other TACs [23].

It is important to consider cluster headache therapies in understanding the hypothalamic role in TACs. Preventative therapy for cluster headaches and other trigeminal autonomic cephalgia headaches includes verapamil and lithium. Verapamil, a calcium-channel blocker, should be considered for treatment of cluster headache prevention. Verapamil modulates muscarinic, serotonergic, and dopaminergic receptor activity of central neurons and increases noradrenaline release by inhibiting presynaptic adrenergic receptors [24]. The inhibition of presynaptic adrenergic receptors is most critical in the hypothalamus. While the mechanism of action of lithium is largely unknown, lithium has been found to accumulate in the hypothalamus, which may provide some insight to its use in cluster headaches. There is evidence to suggest that lithium also modulates serotonin pathways and inhibits the spontaneous release of serotonin in the hypothalamus and other areas of the brain. Additionally, lithium inhibits the enkephalin system and circadian sleep cycles [24].

The exact pathophysiology of trigeminal autonomic cephalgias is partially unclear; however, some data suggest the inclusion of the trigeminovascular system, the cranial autonomic system, and the hypothalamus [25]. The trigeminovascular system is largely responsible for the pain experienced in TACs. This system receives signals from the trigeminal nerves (CN V1–3) and projects nociceptive signals to the brainstem and upper cer-

vical spinal cord, then to the thalamus, and lastly to the pain neuromatrix (multiple areas of the brain that alter and register many types of pain). The autonomic symptoms of TACs are a result of sympathetic inactivation or parasympathetic overactivation via the transmissions between the superior salivatory nucleus and the sphenopalatine ganglion [25]. The hypothalamus, conversely, may serve as a large component in TACs. Considering that the hypothalamus is responsible for circadian rhythms and partly responsible for aggressive emotions, the restlessness reported in TACs and the clocklike patterns of cluster headaches may be attributed to the hypothalamus. One study used PET scans to assess cerebral blood flow during nitroglycerin-induced cluster headaches and found repeated dysfunction in the posterior hypothalamus [21]. Reduced melatonin has also been noted in cluster headaches [26]. Additionally, stimulation of the posterior hypothalamus has been shown successful in treating TAC attacks [27]. Based on these outcomes, it is appropriate to assume significant hypothalamic role in the pathophysiology of cluster headaches.

Wernicke-Korsakoff Syndrome

Wernicke's encephalopathy (WE) is a preventable and treatable neurological syndrome in which vitamin B1, or thiamine, has been depleted in the central nervous system. Within the body, thiamine is converted to thiamine pyrophosphate synthetase to the metabolically active form of thiamine, thiamine pyrophosphate. Thiamine pyrophosphate is a crucial cofactor in glucose metabolism throughout the body, including the central nervous system. A reduction of thiamine pyrophosphate activates the pyruvate dehydrogenase kinase (PDK) in the brain, which triggers neuronal phosphorylation of pyruvate dehydrogenase E1 α (p-PDE1 α). Phosphorylated p-PDE1 α consequently causes reduced mitochondrial aerobic metabolism that can lead to neurodegeneration [28].

Thiamine deficiency syndromes like Wernicke's encephalopathy are almost exclusively seen in alcoholics in the United States [29]. Ethanol (EtOH) inhibits the conversion of thiamine to thia-

mine pyrophosphate and impairs absorption of thiamine in the duodenum. Additionally, severe hepatic impairment, as commonly seen in chronic alcoholics, leads to decreased conversion of thiamine to thiamine pyrophosphate and reduced thiamine stores in the liver as the liver becomes infiltrated with fatty acids or cirrhotic [28].

Other causes of thiamine deficiency include malnutrition, malabsorptive syndromes (i.e., inflammatory bowel disease), and eating disorders (i.e., anorexia). The classical presentation of Wernicke's encephalopathy is characterized by the triad of confusion, ataxia, and ophthalmoplegia. However, realistically only ~10–16% of affected patients have all three of the triad symptoms together [30]. Other symptoms include hypotension, tachycardia, hearing loss, daytime somnolence, dysphagia, seizures, memory impairment, psychosis, hypothermia, hyperhidrosis, and peripheral neuropathy [1]. Many structures of the central nervous system that can be affected by Wernicke's encephalopathy are visible on MR sequences including the mammillary bodies, periaqueductal and periventricular gray matter, superior and inferior colliculus, and thalamus. CNS alterations seen on MRI in WE are typically symmetrical [31]. One of the most well-known structural sequelae of chronic Wernicke's encephalopathy is mammillary body hemorrhage and necrosis. Early recognition of Wernicke's encephalopathy is important as it is associated with significant morbidity and mortality and can be reversed with thiamine supplementation. Without diagnosis and/or intervention, Korsakoff syndrome can develop.

Wernicke-Korsakoff syndrome (WKS) is an irreversible neurological condition characterized by cognitive/memory impairment, confabulation, and personality changes [1]. The mortality rate of WKS is near 17% in severe cases; however, given the difficulty in diagnosis with nonspecific and variable signs/symptoms, the true mortality rate is difficult to determine [32]. The treatment of choice is parenteral thiamine (considering gastrointestinal malabsorption of thiamine is a key factor in the disease process) in addition to oral thiamine supplementation as early a diagnosis is suspected.

Neurodegenerative Disorders

Neurodegenerative diseases are a group of progressive disorders characterized by deterioration of brain structures and functions. This group of disorders include age-related Alzheimer’s disease, early-onset Alzheimer’s disease, frontotemporal dementia, Parkinson’s disease, Parkinson-plus syndromes (multiple systems atrophy, progressive supranuclear palsy, corticobasal degeneration), spinocerebellar ataxias, Huntington disease, amyotrophic lateral sclerosis (ALS), and prion diseases (Table 7.6 shows major neurodegenerative conditions and their characteristic features). Patients with neurodegenerative disorders often present with complaints of cognitive/memory impairments, ataxia, and motor dysfunction, lack of energy, endocrine abnormalities,

changes in behavior/mood, and changes in patterns of food intake.

Alzheimer’s disease (AD) is one of the most common types of neurodegenerative disorder/dementia that typically slowly progresses and causes difficulties in memory, cognition, and behavior [33]. The name “Alzheimer’s disease” can be referenced to two subtypes: early-onset or late-onset (Table 7.7 shows differences between early-onset AD and late-onset AD). Early-onset Alzheimer’s disease is much less common than its variant and has been associated with familial causes – mutations of amyloid precursor protein (APP), presenilin 1 (PSEN1), and presenilin 2 (PSEN2) genes [34]. However, some forms of early-onset AD are also sporadic like the classically known late-onset AD. Both variants are characterized by the presence of neurofibrillary

Table 7.6 Neurodegenerative diseases and characteristic features

Neurodegenerative disease	Characteristic features
Alzheimer’s disease	Progressive memory and cognitive impairment with loss of ability to carry out simple tasks
Frontotemporal dementia	Personality changes, impulsivity/disinhibition, loss of language skills
Parkinson’s disease	Resting tremor, bradykinesia, rigidity, postural instability
Parkinson-plus syndromes	<i>Multiple system atrophy</i> – autonomic features (postural hypotension, urinary/bowel dysfunction, sweating abnormalities) <i>Progressive supranuclear palsy</i> – early falls, vertical gaze palsy <i>Corticobasal degeneration</i> – gradual loss of motor skills, apraxia along with cognitive and behavioral deficits
Spinocerebellar ataxia	Progressive loss of coordination of gait, arm movement, eye movement, and speech
Huntington disease	Chorea, dementia
Amyotrophic lateral sclerosis	UMN and LMN signs
Prion diseases	Quickly progressive infectious disease with psychiatric features and abnormal movements

Table 7.7 Feature variations between early-onset Alzheimer’s disease and late-onset Alzheimer’s disease [34]

	Early-onset Alzheimer’s disease	Late-onset Alzheimer’s disease
Age of onset	<65 years old	>65 years old
Genetic predisposition	Significant predisposition	Complex
Progression	Typically accelerated clinical progression; increased mortality after onset	Relatively more gradual in clinical progression
Presenting symptoms	Language, visuospatial, cognitive deficits	Memory loss
MRI	Widespread cortical atrophy with parietal cortical predominance	Bilateral temporal atrophy
PET scan	Increased parietal hypometabolism	Bilateral temporal hypometabolism
CSF analysis	Low amyloid-β, increased total and phosphorylated tau	Low amyloid-β, increased total and phosphorylated tau

tangles, which are hyperphosphorylated tau protein aggregates intracellularly and accumulated amyloid plaques of A β peptides extracellularly. These histopathological findings can lead to parenchymal inflammation, glial dysfunction, microtubule hindrance, and inhibition of neuronal cell signaling [34].

Many studies have noted hypothalamic involvement and atrophy in patients affected by Alzheimer's disease [35–37]. Expectedly, the volume of the hypothalamus was reduced (by an average of about 11%), and many hypothalamic nuclei exhibited amyloid plaques and neurofibrillary tangles [37]. Additionally in AD, orexin levels are slightly reduced in CSF studies, and suprachiasmatic nuclei degeneration has been noted – which may account for sleep disturbances seen in AD [37]. Altered food-intake behaviors are seen in roughly 50–60% of AD cases, declined nutritional status is seen in 14–80%, and weight loss is seen in 20–45% [37]. One theory suggests that the hypothalamus is largely to blame for these weakening sequelae of AD; although, a definitive cause is currently unknown. It is important to mention, however, that obesity and diabetes mellitus are significant risk factors for developing Alzheimer's disease. Paradoxically, patients with Alzheimer's disease are also at risk for developing diabetes, which may augment evidence that suggests metabolic derangements are a consequence of AD. Weight loss and tight glycemic control are imperative for patients already at high risk for AD development.

Frontotemporal dementia (FTD), previously known as Pick's disease, is another subtype of neurodegenerative disease which, as its name implies, affects the frontal and temporal lobes. FTD is the second most prevalent type of neurodegenerative disease, manifesting most commonly between the ages of 45 and 65 [38]. The typical signs and symptoms include personality/behavioral changes, abulia (an absence of will-power or drive), and expressive/receptive aphasia. Interestingly, eating habit changes frequently occur in patients with frontotemporal dementia as well; however, inversely to Alzheimer's dis-

ease, hyperphagia is more prevalent as well as weight gain [39].

It is presumed that FTD affects majorly the posterior hypothalamus, specifically the mammillary bodies, paraventricular nucleus, and dorsomedial nuclei via atrophy [37]. Volume loss of the hypothalamus on average was slightly more than that of Alzheimer's disease (~15–20%). Those seen with more atrophy of the hypothalamus had more significant eating behavioral changes further suggesting the role of the hypothalamus in nutritional intake [37].

Amyotrophic lateral sclerosis (ALS) is also a progressive neurodegenerative disease that affects primarily motor neurons. ALS is a process with no known cure nor definitive causal etiology to date. A large current theory is that there are minimal hereditary factors and variants but a larger unclear environmental component [40]. The hallmark of this disease is the conjunction of both upper motor neuron disease (i.e., hyperreflexia, spasticity) and lower motor neuron disease (i.e., hypotonia, lack of or loss of reflexes, fasciculations) patterns (Table 7.8 demonstrates characteristics of upper and lower motor neuron lesions). While the extent of the disease is not typically confined either to the spinal cord or to neurons within the cranium, the disease process can affect the hypothalamus [41].

One study found pathological amounts of transactive response DNA-binding protein 43 (TDP-43) seen in the hypothalamus in roughly one third of the ALS patient sample population [42]. The amount of these inclusions had no correlation to the severity of the disease state. Additionally, as seen in AD and FTD, atrophy can also be noted; however, in ALS, there was no discrimination in hypothalamic areas – involving the anterior and posterior aspects.

Based on this discussion of few neurodegenerative disease processes, it is important to recognize the importance of eating habit changes in patients affected. While practitioners may commonly focus on the cognitive impairments, motor/sensory abnormalities, and affective medications, it is important to screen for nutritional status changes as both weight loss and weight

Table 7.8 Upper motor neuron vs. lower motor neuron disease

	Upper motor neuron (UMN) lesion	Lower motor neuron (LMN) lesion
Muscle strength	Weakness/paralysis	Weakness/paralysis
Type of weakness	Spastic paresis	Flaccid paralysis
Muscle tone	Spasticity, rigidity	Reduced; hypotonia
Deep tendon reflexes (DTRs)	Increased; hyperreflexia	Reduced; hyporeflexia
Atrophy	Disuse atrophy (slight)	Severe atrophy
Plantar response	Extensor	Normal; flexor
Fasciculation	Absent	Present
Electromyography	Normal; reduced interference pattern and firing rate	Fasciculations and fibrillations

gain can have significant worsening of prognosis. Diabetes and hyperlipidemia should be routinely screened for in all patients and particularly in those with hyperphagic symptoms. Malnutrition and depressive symptoms should also be considered in all patients affected by neurodegenerative diseases and particularly in those with a reduced appetite.

Narcolepsy

Narcolepsy is a neurological syndrome affecting the ability to self-regulate sleep-wake cycles characterized by excessive daytime somnolence, intermittent loss of consciousness, cataplexy (sudden loss of muscle tone), sleep paralysis, and hypnagogic hallucinations (Table 7.9 ICSD-3 Narcolepsy Diagnostic Criteria) [43]. The prevalence of this disorder is roughly 47 individuals per 100,000 in the general population [44]. Age of onset varies but typically peaks in adolescence and the fourth decade of life. Two subtypes of narcolepsy exist: Type 1 and Type 2 narcolepsy. Type 1 narcolepsy is diagnosed when an affected individual has associated cataplexy or low orexin, or hypocretin, hormone in the CSF. Type 2 narcolepsy is diagnosed when a patient experiences excessive daytime somnolence without cataplexy or reduced CSF orexin levels [45]. While the exact cause for narcolepsy remains poorly understood, many theories exist. An important and consistent finding is reduced level of CSF orexin, which is a neuropeptide released by the dorsolateral hypothalamus that plays an important role in

sleep regulation and wakefulness [46]. While it may be understood that individuals living with narcolepsy have these findings, the presumed causal factors include autoimmunity, post-infectious, genetics, or trauma [46, 47].

Diagnosis can be made with a polysomnogram, which may reveal reduced REM latency, or a multiple sleep latency test (MSLT), which may also show a quickness to REM sleep. Treatment typically involves lifestyle modifications, avoidance of triggers, and sometimes stimulant medications, like Modafinil or amphetamine-like medications [46]. It is important to note that in the brain, hypocretin is solely produced by the hypothalamus, which projects mainly to the locus coeruleus, dorsal raphe nuclei, amygdala, suprachiasmatic nucleus, cholinergic nuclei in the brainstem, and spinal cord [48]. The neuropeptide acts mostly as an excitatory neurotransmitter to the aforementioned structures, in turn, increasing serotonin, dopamine, norepinephrine, epinephrine, histamine, and acetylcholine [48]. Secondarily, orexins are believed to have some role in other functioning such as energy regulation, eating habits, metabolism, sexual desire, and gut motility/acid production and have some role in increasing mean arterial blood pressure and pulse rate [46]. One postmortem study comparing hypothalami of individuals with narcolepsy versus controls showed remarkable reductions (85–95%) in the density of hypocretin-producing neurons [49]. An increased amount of gliosis and astrocytes were found suggesting degeneration of these neurons [49]. Other conditions in which hypocretin is believed to be

Table 7.9 ICSD-3 Narcolepsy Diagnostic Criteria

Narcolepsy type 1
<i>Alternate names:</i> Hypocretin deficiency syndrome, narcolepsy-cataplexy, narcolepsy with cataplexy
<i>Diagnostic criteria:</i> Criteria A & B must be met
A. The patient has daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for ≥ 3 months ^a
B. The presence of one or both of the following:
1. Cataplexy (as defined under Essential Features) and a mean sleep latency of ≤ 8 minutes and two or more sleep onset REM periods (SOREMPs) on an MSLT performed according to standard techniques. A SOREMP (within 15 minutes of sleep onset) on the preceding nocturnal polysomnogram may replace one of the SOREMPs on the MSLT ^b
2. CSF hypocretin-1 concentration, measured by immunoreactivity, is either ≤ 110 pg/mL or $< 1/3$ of mean values obtained in normal subjects with the same standardized assay

Notes:

Originally published in the international classification of sleep disorders, third edition, published by the American Academy of sleep medicine. Copyright American Academy of sleep medicine, used with permission

CSF cerebrospinal fluid, MSLT multiple sleep latency test, REM rapid eye movement

^aIn young children, narcolepsy may sometimes present as excessively long night sleep or as resumption of previously discontinued daytime napping

^bIf narcolepsy type 1 is strongly suspected clinically but the MSLT criteria of B1 are not met, a possible strategy is to repeat the MSLT

Narcolepsy type 2
<i>Alternate names:</i> Narcolepsy without cataplexy
<i>Diagnostic criteria:</i> Criteria A–E must be met:
A. The patient has daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for ≥ 3 months
B. A mean sleep latency ≤ 8 minutes and ≥ 2 sleep-onset REM periods (SOREMPs) are found on an MSLT performed according to standard techniques. A SOREMP (within 15 minutes of sleep onset) on the preceding nocturnal polysomnogram may replace 1 of the SOREMPs on the MSLT
C. Cataplexy is absent ^a
D. Either CSF hypocretin-1 concentration has not been measured or CSF hypocretin-1 concentration measured by immunoreactivity is either > 110 pg/mL or $> 1/3$ of mean values obtained in normal subjects with the same standardized assay ^b
E. The hypersomnolence and/or MSLT findings are not better explained by other causes such as insufficient sleep, obstructive sleep apnea, delayed sleep phase disorder, or the effect of medication or substances or their withdrawal

Notes:

Originally published in the international classification of sleep disorders, third edition, published by the American Academy of sleep medicine. Copyright American Academy of sleep medicine, used with permission

CSF cerebrospinal fluid, MSLT multiple sleep latency test, REM rapid eye movement

^aIf cataplexy develops later, then the disorder should be reclassified as narcolepsy type 1

^bIf the CSF hypocretin-1 concentration is tested at a later stage and found to be either ≤ 110 pg/mL or $< 1/3$ of mean values obtained in normal subjects with the same assay, then the disorder should be reclassified as narcolepsy type 1

involved with is Kleine-Levin syndrome, age-related reduction in REM sleep, and other primary sleep disorders (hypersomnolence and insomnia-related).

Kleine-Levin Syndrome

Kleine-Levin syndrome (KLS) is a peculiar and rare neurological disorder affecting typically teenaged boys that is characterized by episodic

hypersomnolence, hypersexualism, impulsive eating, and cognitive impairment. These episodes of hypersomnolence can vary from days to months at a time, and patients affected with KLS are often unable to perform their activities of daily living (i.e., going to school). Given the rarity of KLS, limited information is currently understood about this disease. Based on the symptomatology of KLS, it has long been postulated that the hypothalamus is the culprit. One case report in 1987 [50] showed slightly elevated

prolactin levels and abnormalities in LH and 11-OHCS. Another more recent study of five KLS patients noted normal hormonal patterns in sleep except hGH, suggesting no underlying circadian disorder [51].

Special Thanks With special thanks to the AASM for generously allowing permission of the diagnostic criteria for narcolepsy.

References

1. Ropper, A. H., Adams, R. D., Victor, M., Brown, R. H., & Victor, M. (2005). *Adams and Victor's principles of neurology*. New York: McGraw-Hill Medical Pub. Division.
2. Dawson B. The blood vessels of the human optic chiasma and their relation to those of the hypophysis and hypothalamus. *Brain*. 1958;81:207–17.
3. De Silva K, Silva R, Amaratunga D, Gunasekera W, Jayasekera R. Types of the cerebral arterial circle (circle of Willis) in a Sri Lankan population. *BMC Neurol*. 2011; <https://doi.org/10.1186/1471-2377-11-5>.
4. Flanagan E. Neuromyelitis Optica Spectrum disorder and other non-multiple sclerosis central nervous system inflammatory diseases. *Continuum: Lifelong Learning in Neurology*. 2019;25:815–44.
5. Huitinga I, De Groot C, Van der Valk P, Kamphorst W, Tilders F, Swaab D. Hypothalamic lesions in multiple sclerosis. *J Neuropathol Exp Neurol*. 2001;60:1208–18.
6. Erkut Z, Hofman M, Ravid R, Swaab D. Increased activity of hypothalamic corticotropin-releasing hormone neurons in multiple sclerosis. *J Neuroimmunol*. 1995;62:27–33.
7. Gao C, Wu L, Chen X, Long Y, Zhong R, Yang N, Chen Y. Hypothalamic abnormality in patients with inflammatory demyelinating disorders. *Int J Neurosci*. 2016;126:1036–43.
8. Kandel E, Schwartz J, Jessell T. *Principles of neural science*. 10th ed. New York: McGraw-Hill, Health Professions Division; 2009.
9. Ruiz-Gaviria R, Baracaldo I, Castañeda C, Ruiz-Patiño A, Acosta-Hernandez A, Rosselli D. Specificity and sensitivity of aquaporin 4 antibody detection tests in patients with neuromyelitis optica: a meta-analysis. *Mult Scler Relat Disord*. 2015;4:345–9.
10. Irani S. Limbic encephalitis. In: *The Encephalitis Society*. 2019. <https://www.encephalitis.info/limbic-encephalitis>. Accessed 6 Dec 2019.
11. Lai M, Huijbers M, Lancaster E, Graus F, Bataller L, Balice-Gordon R, Cowell J, Dalmau J. Investigation of LGI1 as the antigen in limbic encephalitis previously attributed to potassium channels: a case series. *Lancet Neurol*. 2010;9:776–85.
12. Pollak T, McCormack R, Peakman M, Nicholson T, David A. Prevalence of anti-N-methyl-D-aspartate (NMDA) receptor antibodies in patients with schizophrenia and related psychoses: a systematic review and meta-analysis. *Psychol Med*. 2013;44:2475–87.
13. Hoftberger R, van Sonderen A, Leypoldt F, et al. Encephalitis and AMPA receptor antibodies: novel findings in a case series of 22 patients. *Neurology*. 2015;84:2403–12.
14. Dalmau J. Clinical analysis of anti-Ma2-associated encephalitis. *Brain*. 2004;127:1831–44.
15. Charles A. The migraine Aura. *Continuum: Lifelong Learning in Neurology*. 2018;24:1009–22.
16. Karsan N, Bose P, Goadsby P. The migraine premonitory phase. *Continuum: Lifelong Learning in Neurology*. 2018;24:996–1008.
17. Brennan K, Romero-Reyes M, López Valdés H, Arnold A, Charles A. Reduced threshold for cortical spreading depression in female mice. *Ann Neurol*. 2007;61:603–6.
18. Eikermann-Haerter K, Baum M, Ferrari M, van den Maagdenberg A, Moskowitz M, Ayata C. Androgenic suppression of spreading depression in familial hemiplegic migraine type 1 mutant mice. *Ann Neurol*. 2009;66:564–8.
19. Charbit A, Akerman S, Goadsby P. Trigemino-cervical complex responses after lesioning dopaminergic A11 nucleus are modified by dopamine and serotonin mechanisms. *Pain*. 2011;152:2365–76.
20. Bartsch T, Levy M, Knight Y, Goadsby P. Inhibition of nociceptive dural input in the trigeminal nucleus caudalis by somatostatin receptor blockade in the posterior hypothalamus. *Pain*. 2005;117:30–9.
21. Maniyan F, Sprenger T, Monteith T, Shankin C, Goadsby P. Brain activations in the premonitory phase of nitroglycerin-triggered migraine attacks. *Brain*. 2013;137:232–41.
22. Schulte L, May A. The migraine generator revisited: continuous scanning of the migraine cycle over 30 days and three spontaneous attacks. *Brain*. 2016;139:1987–93.
23. Burish M. Cluster headache and other trigeminal autonomic cephalalgias. *Continuum: Lifelong Learning in Neurology*. 2018;24:1137–56.
24. Costa A, Antonaci F, Ramusino M, Nappi G. The neuropharmacology of cluster headache and other trigeminal autonomic cephalalgias. *Curr Neuropharmacol*. 2015;13:304–23.
25. Burish M, Rozen T. Trigeminal autonomic cephalalgias. *Neurol Clin*. 2019;37:847–69.
26. Gelfand A, Goadsby P. The role of melatonin in the treatment of primary headache disorders. *Headache J Head Face Pain*. 2016;56:1257–66.
27. Messina G, Broggi G, Levi V, Franzini A. Deep brain stimulation for trigeminal autonomic cephalalgias. *Expert Rev Neurother*. 2018;18:421–6.
28. Butterworth R. Thiamin deficiency and brain disorders. *Nutr Res Rev*. 2003;16:277–84.

29. Sinha S, Kataria A, Kolla B, Thusius N, Loukianova L. Wernicke encephalopathy—clinical pearls. *Mayo Clin Proc.* 2019;94:1065–72.
30. Cook C. Prevention and treatment of Wernicke-Korsakoff syndrome. *Alcohol Alcohol.* 2000;35:19–20.
31. Shigekiyo T, Tani H, Nakajima H, Kimura F. Antecedent diffuse cerebellar lesions on diffusion-weighted MRI in nonalcoholic Wernicke's encephalopathy. *Neurol Sci.* 2017;39:789–90.
32. Day G, del Campo C. Wernicke encephalopathy: a medical emergency. *Can Med Assoc J.* 2013;186:E295.
33. Hampel H, Toschi N, Babiloni C, et al. Revolution of Alzheimer precision neurology. Passageway of systems biology and Neurophysiology. *J Alzheimers Dis.* 2018;64:S47–S105.
34. Rabinovici G. Late-onset Alzheimer disease. *Continuum: Lifelong Learning in Neurology.* 2019;25:14–33.
35. Mann D. Changes in Alzheimer's disease in the magnocellular neurones of the supraoptic and paraventricular nuclei of the hypothalamus and their relationship to the noradrenergic deficit. *Clin Neuropathol.* 1985;4:127–34.
36. Musiek E, Bhimasani M, Zangrilli M, Morris J, Holtzman D, Ju Y. Circadian rest-activity pattern changes in aging and preclinical Alzheimer disease. *JAMA Neurol.* 2018;75:582.
37. Vercruysse P, Vieau D, Blum D, Petersén Å, Dupuis L. Hypothalamic alterations in neurodegenerative diseases and their relation to abnormal energy metabolism. *Front Mol Neurosci.* 2018;11:2. <https://doi.org/10.3389/fnmol.2018.00002>.
38. Finger E. Frontotemporal dementias. *Continuum: Lifelong Learning in Neurology.* 2016;22:464–89.
39. Olney N, Spina S, Miller B. Frontotemporal dementia. *Neurol Clin.* 2017;35:339–74.
40. Zarei S, Carr K, Reiley L, Diaz K, Guerra O, Altamirano P, Pagani W, Lodin D, Orozco G, China A. A comprehensive review of amyotrophic lateral sclerosis. *Surg Neurol Int.* 2015;6:171.
41. Ngo S, Mi J, Henderson R, McCombe P, Steyn F. Exploring targets and therapies for amyotrophic lateral sclerosis: current insights into dietary interventions. *Degener Neurol Neuromuscul Dis.* 2017;7:95–108.
42. Cykowski M, Takei H, Schulz P, Appel S, Powell S. TDP-43 pathology in the basal forebrain and hypothalamus of patients with amyotrophic lateral sclerosis. *Acta Neuropathol Commun.* 2014;2:171. <https://doi.org/10.1186/s40478-014-0171-1>.
43. Ropper, A. H., Adams, R. D., Victor, M., Brown, R. H., & Victor, M. (2005). *Adams and Victor's principles of neurology: "Sleep and its Abnormalities"*. New York: McGraw-Hill Medical Pub. Division.
44. Narcolepsy Fact Sheet | National Institute of Neurological Disorders and Stroke. 2019. In: [Ninds.nih.gov. https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Fact-Sheets/Narcolepsy-Fact-Sheet](https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Fact-Sheets/Narcolepsy-Fact-Sheet). Accessed 6 Dec 2019.
45. Lecendreux M, Lavault S, Lopez R, Inocente C, Konofal E, Cortese S, Franco P, Arnulf I, Dauvilliers Y. Attention-Deficit/Hyperactivity Disorder (ADHD) symptoms in pediatric narcolepsy: a cross-sectional study. *Sleep.* 2015;38:1285–95.
46. Black S, Yamanaka A, Kilduff T. Challenges in the development of therapeutics for narcolepsy. *Prog Neurobiol.* 2017;152:89–113.
47. Mignot E. History of narcolepsy at Stanford University. *Immunol Res.* 2014;58:315–39.
48. Siegel J, Boehmer L. Narcolepsy and the hypocretin system—where motion meets emotion. *Nat Clin Pract Neurol.* 2006;2:548–56.
49. Thannickal T, Moore R, Nienhuis R, Ramanathan L, Gulyani S, Aldrich M, Cornford M, Siegel J. Reduced number of hypocretin neurons in human narcolepsy. *Neuron.* 2000;27:469–74.
50. Gadoth N, Dickerman Z, Bechar M, Laron Z, Lavie P. Episodic hormone secretion during sleep in Kleine-Levin syndrome: evidence for hypothalamic dysfunction. *Brain Dev.* 1987;9:309–15.
51. Mayer G, Leonhard E, Krieg J, Meier-Ewert K. Endocrinological and polysomnographic findings in Kleine-Levin syndrome: no evidence for hypothalamic and circadian dysfunction. *Sleep.* 1998;21:278–84.

Neuropsychiatric and Neurobehavioral Syndromes of the Human Hypothalamus

8

John Wagner III, Noeen Sarfraz, Kunal Maini,
and Amber N. Edinoff

Introduction

The role that the human hypothalamus plays in behavior, as well as psychiatric disorders, has been known for quite some time. As early as the 1930s, Selye studied the physiological response to stress [1]. His work led to our understanding of the hypothalamic-pituitary-adrenal (HPA) axis. Hypothalamic activity influences human behavior [2, 3] and plays a role in a wide range of mental disorders. These include anxiety disorders, affective disorders such as major depression and bipolar, and substance use disorders.

The Hypothalamus and Basic Behavior

The hypothalamus mediates a variety of basic human behaviors, including sexual, food and water intake, fear, and aggressiveness. These are considered as evoked behaviors or “survival needs,” and they are the result of neuronal stimuli upon discrete hypothalamic regions [3]. The

posterior nucleus of the posterior hypothalamus has been linked to the drive for drugs and sex, the mammillary bodies to episodic memory. Pascual et al. describe the result of damage by craniopharyngiomas to various hypothalamic structures and resulting psychiatric disorders. In their systemic review and meta-analysis of work investigating the damage caused by craniopharyngiomas to hypothalamic structures and resulting psychiatric disorders, Pascual et al. describe a loss of episodic memory resulting from damage to the mammillary bodies. In particular, there was a significantly higher proportion of atrophic or absent mammillary bodies in subjects with a psychiatric disorder than in those without, and tumor distortion of the mammillary bodies was noted to the key anatomic distortion associated with episodic memory loss. Memory loss was observed in up to 86% of subjects with atrophic mammillary bodies [4].

Interestingly, some of these behaviors have been observed during neurosurgical procedures. In 1933, Foerster and Gagel published their observations made during the removal of a cyst from the third ventricle. Whenever the ventricle floor was contacted, the patient laughed, whistled, and made obscene remarks. These behaviors ceased when contact with the ventricular floor was removed [2]. The periventricular zone (PVZ) was implicated in mood regulation by Angelergues et al. in a case reported of persistent euphoria after a bilateral infarct [5].

J. Wagner III (✉) · N. Sarfraz · K. Maini
A. N. Edinoff
Department of Psychiatry and Behavioral Medicine,
LSU Health Shreveport, Shreveport, LA, USA
e-mail: JWagne1@lsuhsc.edu; nsarfr@lsuhsc.edu;
kmmaini@lsuhsc.edu; aedino@lsuhsc.edu

Multiple regions of the hypothalamus affect sexual drive and behavior. Small lesions in the medial preoptic area/anterior hypothalamus (MPOA/AH) temporarily affected male rat sexual drive [6]. The dorsomedial and ventromedial nuclei of the hypothalamus (DMH and VMN) affect penile erection through projections to the dorsolateral funiculus in the spinal cord [7]. The ventromedial hypothalamus was a frequent target for stereotactic brain lesion surgery to treat paraphilia and exhibitionism in the 1960s and 1970s in Germany. These procedures drastically reduced sexual drive [8]. The paraventricular nucleus is active during sexual activity, including orgasm, most likely through the production of oxytocin, vasopressin, and dopamine [7]. A recent quantitative meta-analysis of visual cues shows consistent activation of the hypothalamus [8].

The hypothalamus has also been a target for neurosurgical intervention to treat refractory aggressive behaviors. From 1962 to 1977, Sano conducted stereotactic hypothalamotomy surgeries focusing on the posterior hypothalamus on 60 patients with severe aggressive behaviors. He later published the long-term outcomes, which showed a 78% positive result, with no violent or aggressive behaviors exhibited postoperatively [9]. Schwarcz reported on 11 cases of stereotactic posterior hypothalamotomy to treat severe aggressiveness with associated violent behavior. In each case, psychiatric treatment had failed. At follow-up, up to 4 years postop, 10 of the 11 patients had marked improvement with or without psychotropic medication. One was deemed a failure [10].

Although psychosurgery has fallen out of favor due to public outcry, the advent of deep brain stimulation (DBS), which is both less destructive and reversible, is a viable alternative. DBS, along with the development of more advanced stereotactic neurosurgical techniques and improved neuroimaging, provides neurosurgeons with an intervention to address psychopathological symptoms. Thus, the hypothalamus is once again a target for treatment. Several case reports and a cohort of seven patients have been published documenting DBS application to the posterior hypothalamus to treat aggressive behavior, with positive results [11].

The Hypothalamus and Anxiety

The DSM-5 describes a variety of anxiety disorders. These include (with their 12-month prevalence rates) separation anxiety disorder (0.9–1.9%), social anxiety disorder (7%), panic disorder (2–3%), agoraphobia (1.7%), specific phobia (7–9%), and generalized anxiety disorder (2.9%). The DSM-5 classifies post-traumatic stress disorder (PTSD) separately as a trauma-related disorder. Its lifetime prevalence rate is 8.7%. Anxiety disorders occur more frequently in females with a ratio of 2:1. Excessive fear and anxiety as well as behavioral disturbances are common to all anxiety disorders. Fear is the response to a real or perceived threat often with autonomic arousal. Anxiety is associated with internalizing symptoms including muscle tension, hyper-vigilance, and avoidance behaviors [12].

HPA axis dysfunction can lead to the development of anxiety disorders [13]. The HPA axis has been shown to demonstrate dramatic reactivity to psychological and/or physiological stressors, as evidenced by increased production of hormones like adrenocorticotrophic hormone (ACTH), corticotropin-releasing factor (CRF), and cortisol (aptly termed a “stress hormone”). In addition to dysregulation to the HPA axis, insults to the hypothalamic-pituitary-thyroid (HPT) axis have also been shown to play a role in the development of anxiety disorders vis-à-vis an increase in the triiodothyronine (T3)/thyroxine (T4) ratio [14]. In a systematic review, Fischer and Cleare synthesize the literature on the relationship between cortisol levels and response to psychotherapy in individuals with anxiety disorders [15]. They conclude that current evidence is equivocal in terms of the predictive value of basal cortisol levels in response to psychotherapy for the treatment of anxiety disorder. In another systematic review on the role the HPT plays in anxiety disorders, Fischer and Ehler found that nearly all 20 studies included in the review revealed a significant degree of comorbidity between thyroid disorder and symptoms of anxiety [16]. The findings of the systematic review support current clinical guidelines to test for thyroid disorders in patients

presenting with symptoms of anxiety, including, but not limited to, palpitations and diaphoresis.

While the role of the HPA axis in disorders such as generalized anxiety disorder is well understood, such is not the case for PTSD [14, 17]. Similar to other anxiety disorders, plasma levels of CRH are abnormally elevated in PTSD, but in cases of PTSD, the response of ACTH to CRH is blunted. These findings have implications for treatment modalities for PTSD and represent an area of continued research.

The Hypothalamus and Mood Disorders

Hypothalamic-pituitary-adrenal (HPA) axis dysfunction has been implicated in mood disorders, including major depressive disorder, postpartum depression, and bipolar disorder.

Major depressive disorder (MDD) is the most prevalent psychiatric disorder worldwide. The annual incidence rate in developed countries is 10%, while 20% of individuals are depressed during their lifetime [18]. According to the DSM-5, five or more of the following symptoms must be present during the same 2-week period: (1) depressed mood most of the day nearly every day, (2) diminished interest or pleasure in activities, (3) significant weight loss or weight gain, (4) insomnia or hypersomnia, (5) psychomotor agitation or retardation, (6) fatigue, (7) feelings of worthlessness or guilt, (8) decreased concentration or indecisiveness, and (9) recurrent thoughts of death or suicidal ideation (with or without a specific plan) [12]. Once diagnosed, current pharmacologic treatment with antidepressants is effective in up to 70% of patients. Unfortunately, a substantial number of patients (19–34%) fail to respond fully, and up to 34% are treatment-resistant [19].

Psychosocial stressors play a significant role in the development of depression. Childhood exposure to stressors, such as the death of a parent or caregiver, maternal deprivation, paternal abandonment, parental separation, or divorce, increases the risk of adult depressive episodes. The likely mechanism for this relationship

between acute stress and depression lies in the abnormal activation of the HPA axis.

HPA axis dysfunction can result in hypersecretion of cortisol, enlargement of the pituitary and adrenal glands, and elevated corticotropin-releasing factor (CRF), also known as corticotropin-releasing hormone. Depressed patients have shown to have elevated 24-hour urinary free cortisol, elevated serum cortisol, and elevated CSF cortisol. Other abnormalities in depressed patients include non-suppression of cortisol, beta-endorphin, and ACTH after dexamethasone challenge. Postmortem examination has shown evidence of increased CRF mRNA and protein in the PVN of the hypothalamus [20].

There are several theories as to the cause of HPA axis dysfunction in depression. One suggests of increased central drive in the hypothalamus, resulting in hypersecretion of CRF and/or arginine vasopressin (AVP). Interestingly, animal studies have shown that elevated CRF itself is tied to alterations in activity, appetite, and sleep, which are also observed in depressed patients [21]. Chronic stress with associated increased cortisol levels may decrease the hypothalamus' ability to downregulate CRF released by the paraventricular nucleus (PVN) by altering negative feedback of glucocorticoid receptors [4, 22]. Normally, glucocorticoids serve as potent inhibitors of CRF release by the PVN, as well as ACTH by the pituitary, as evidenced by the increased volume of the pituitary in depressed patients [23].

Two types of corticosteroid receptors have been identified, mineralocorticoid receptor (MR) and glucocorticoid receptor (GR), which help regulate the HPA axis. Mineralocorticoid receptors, found mainly in the hippocampus, have a high affinity for endogenous glucocorticoids. Glucocorticoid receptors are found in multiple areas of the brain, including the hippocampus, the amygdala, the brain stem, and the hypothalamus. MR is thought to regulate basal levels of cortisol, and GR regulates at higher or “stressed” levels of cortisol. Thus GR receptors are believed to play a larger role in response to stress. However, the balance of MR and GR activity may affect the set point of the HPA axis. Several studies have shown a decrease in GR response in

depressed patients. It is unknown if this decrease in GR activity is due to the lack of efficacy of the receptor type and a decrease in the number of receptors in the brain. Pariante proposes that the elevated cortisol levels observed in depressed patients are a compensatory mechanism to overcome glucocorticoid resistance [23].

Hypocretins (Hcrts) are neuropeptides produced by the lateral hypothalamus (LH) with projections throughout the brain to include the locus coeruleus, basal forebrain, ventral tegmental area, raphe nuclei, and tuberomammillary nucleus. Animal studies link hypocretins to the HPA axis, showing that stress increases the expression of one type of hypocretin, Hcr1 mRNA in Hcr1 neurons located in the LH [24]. Hyperfunction of the Hcr1 system may be linked to depression, in both unipolar and bipolar depressive episodes. One study showed higher mean levels of CSF Hcr1 levels in these patients. Furthermore, treatment with antidepressants restored Hcr1 to normal levels [25].

Melancholic and Atypical Depressions Types

There are two subtypes of major depressive disorder, the melancholic type and the atypical type. Depression with melancholic features is defined as having either loss of pleasure in all or most activities or lack of emotional reactivity to pleasurable stimuli along with at least three of the following: (1) profound despondency or despair, (2) worse depression in the morning, (3) early-morning awakening, (4) psychomotor agitation or retardation, (5) significant anorexia or weight loss, and (6) excessive guilt. Atypical depression has the following characteristics: mood reactivity and at least two of the following: (1) significant weight gain or increased appetite, (2) hypersomnia, (3) leaden paralysis, and (4) heightened sensitivity to interpersonal rejection [12].

Juruena and colleagues published an extensive review of 48 studies examining the role of HPA functioning in melancholic and atypical depression. They concluded that HPA axis activity differed in these two subtypes of major depression.

Specifically, patients with melancholic depression were shown to have increased cortisol levels, both at baseline and post-challenge. Atypical depressed patients exhibited lower levels of HPA axis activity, suggesting the possibility of enhanced negative feedback [26].

Postpartum Depression

Postpartum depression (PPD) is defined as major depressive disorder with peripartum onset [12]. Criteria for PPD are very similar to that of major depressive disorder, with the additional temporal requirement of onset within 4 weeks of delivery. Risk factors for PPD include ongoing psychosocial stressors such as the lack of social support, marital discord, and socioeconomic status. Other risk factors include a history of childhood trauma and sexual abuse. The severity of symptoms of PPD is related to chronic and acute stress [27].

Bipolar Disorder

Bipolar disorder, historically known as manic depression, is characterized by both episodes of depression and mania. The DSM-5 identifies multiple variations of bipolar disorder due to its heterogeneity in symptom severity and duration. Broadly defined criteria for a bipolar I diagnosis include a manic episode lasting a minimum of 1 week, with a persistently elevated or irritable mood, increased activity or energy, as well as associated symptoms including grandiosity, decreased need for sleep, pressured speech, racing thoughts, and impulsive behavior. Bipolar II can be viewed as a milder form, with no history of a fully manic episode. These individuals have exhibited an episode of hypomania, which may last less than 1 week and is less severe [12].

The hypothalamus may play a role in bipolar disorder through the activity of the neuropeptide oxytocin. Oxytocin, produced predominately by magnocellular cells in the supraoptic and paraventricular nuclei of the hypothalamus, plays a variety of roles including placentation, viviparity, lactation, and smooth muscle contraction of the

uterus during labor. Oxytocin also regulates social interactions, including those between the mother and her offspring, as well as more broad pro-social behaviors such as empathy and trust [28]. Although oxytocin is commonly viewed for its positive impact on psychological functioning, elevated levels of oxytocin may result in increased emotionality and anxiety. Some studies have tied elevated levels of oxytocin to several mental illnesses, including depression, schizophrenia, and bipolar disorder [29]. Yueh-Ju et al. examined oxytocin levels in patients with MDD and bipolar II disorder in a depressed state and found that plasma oxytocin levels were significantly higher in the bipolar II group than either the MDD group or healthy controls. The MDD group's oxytocin levels were elevated compared to controls but were not statistically significant [30].

GABA and Neurosteroids

CRF neuronal activity in the PVN is tightly regulated by GABAergic signaling. Up to one-third of CRF, neuron inputs are GABAergic. GABAergic neurons project to the PVN from multiple hypothalamic sources, including the anterior and lateral areas, the dorsomedial nucleus, the medial preoptic area, as well as the stria terminalis [27]. Culliana et al. showed evidence of the role GABA plays in corticosterone release in rats by microinjection of muscimol into the PVN, which reduces the stress-related elevation of corticosterone with the opposite effect with microinjection of bicuculline [31].

GABA receptors are themselves regulated by allosteric modulators, both positive (PAM) and negative (NAM) types. Neuroactive steroids and neurosteroids can act as both PAMs and NAMs. Several of these have been identified and have been shown to have antidepressant and anxiolytic effects, including allotetrahydrodeoxycorticosterone (THDOC) and allopregnanolone. Both of these are decreased following chronic stress, suggesting that at low levels, the inhibiting effects of GABA on the PVN's release of CRF are decreased [27]. The 5 α -reductase inhibitor finasteride, which blocks neurosteroidogenesis, is

known to cause both anxiety and depression, supporting the model that reduced neurosteroid levels may lead to these disorders. In fact, one possible treatment of anxiety and depression is medications which stimulate neurosteroidogenesis [32].

Schizophrenia

Schizophrenia is a chronic mental health disorder that affects approximately 3.3 million people in the United States. Its symptoms, which must be present for greater than 6 months, comprise disorganized behavior and speech, a diminished capacity to comprehend reality, hearing voices unheard by others, seeing things unseen by others, delusions, and decreased social commitment and motivation. Some individuals may also have magical thinking, such as the belief that others can read their minds, as well as debilitating paranoia with notions that others can read and, thus, control their minds. More severely, certain people can develop catatonic behavior and therefore require help with daily activities [33].

Multiple studies have correlated specific areas of the brain that can be affected in schizophrenia. Evidence not only shows abnormalities in the gray matter but white matter as well. The most evidence-based anatomical brain aberrations found in multiple meta-analysis studies include reduction in total brain volume; hypertrophy of the cavum septum pellucidum and lateral ventricles; atrophy of the midsagittal callosum as well as the thalamus, anterior cingulate cortex, bilateral hippocampus, and amygdala; and, finally, a reduction in size of frontal, parietal, and temporal lobes [34]. Larger basal ganglia and hypothalami were also noted.

Any or all of these abnormalities can be related to glutamate deficiency due to functional deficits in particular neuronal circuits present in these atrophied regions of the brain. Due to these structural abnormalities, patients with schizophrenia can have both hypo- and hyperfunctioning of the hypothalamic-pituitary-adrenal axis, leading to a multitude of complex correlations due to increased or decreased production of

hypothalamic neuropeptides from functionally compromised neuroendocrine neurons.

Studies have shown that the hippocampal changes may be a consequence of the disease process rather than the cause. One theory suggests that the physiological stress induced by the negative symptoms of schizophrenia could lead to altering the structure of the hippocampus, interrupting neurogenesis, decreasing dendritic density, and leading to cell destruction [34].

Schizophrenia typically presents in early adulthood. Females have a later onset of symptoms in comparison to males, due to protective estrogenic effects. Greater exposure to 17 beta-estradiol (E2) at age onset of menarche in females correlates to a later onset of schizophrenia. There was no correlation between the onset of puberty and disease presentation in males. Females typically have less severe symptoms when compared to their male counterparts, including decreased disability in activities of daily living and requiring lower dosing of antipsychotic medication. Furthermore, females admitted for inpatient treatment and stabilization of psychotic symptoms due to schizophrenia also had lower E2 concentrations [35].

Many studies have shown that a continued state of stress can cause HPA axis dysfunction, partly due to decreased feedback of glucocorticoid secretion in the hippocampus or hypothalamus. A study conducted in patients diagnosed with chronic schizophrenia showed much higher basal cortisol levels in comparison to control groups and a larger cortisol surge after dexamethasone pretreatment. Abnormal haloperidol metabolism and a decreased response to antipsychotic treatment were seen due to nonspecific chronic stress combined with cerebral atrophy and ventricular enlargement. Even when these patients have increased basal cortisol levels, they have a reduced response to environmental stressors suggesting a disturbance in HPA axis [36].

These feedback mechanisms that regulate cortisol secretion may also be due to altered or dysfunction of glucocorticoid receptors in the hippocampus. Structural changes, including atrophy, may attribute to NMDA (N-methyl-D-

aspartate) glutamate receptor malfunction and may be the cause of negative symptoms and cognitive impairment. Structural changes in the hippocampus are also worsened by hypercortisolism, thus creating an inverse relationship and a worsening cycle of symptoms [36].

As discussed above, Hcrt neuronal hyperfunction has been linked to affective disorders. Individuals with schizophrenia have significantly higher plasma Hcrt-1 levels versus healthy controls [37]. Antipsychotic medications reduce CSF levels of Hcrt-1 [38].

Schizophrenia may be an NMDA receptor hypofunctioning disorder. NMDA receptors are not able to respond to peptides appropriately. This dysfunction is similar to that seen with ketamine and phencyclidine (PCP) use, which can lead to acute psychosis in otherwise healthy individuals. The atypical antipsychotic clozapine, used for treatment-resistant schizophrenia and other structurally similar atypical antipsychotics, can prevent this NMDA antagonist-induced neurotoxicity making it more effective. Clozapine's effectiveness, particularly concerning addressing negative symptoms, more so than the typical D2 receptor antagonist antipsychotics, is attributed to its effect of decreasing plasma cortisol levels [36]. There is a positive relationship with CNS infections occurring during childhood- and adult-onset schizophrenia and other varying psychosis disorders. Plasma IL-6 and its receptor, TNF-alpha, and IL-2 receptor were shown to be higher in patients with schizophrenia.

Many of the negative symptoms of schizophrenia, which can affect activities of daily living, are correlated to both plasma cortisol levels and multiple cytokines. This data can further be utilized to understand the different causes of negative symptoms better and be used to determine the modalities of the most effective treatment better. Schizophrenia is a multifaceted, complex psychotic disorder, and the correlation of treating with medications that not only affect plasma cortisol levels but glutaminergic receptors as well can lead to a better understanding and higher level of efficacy when considering treatment options.

The majority of these symptoms can be managed with the proper antipsychotic medication with therapy being an adjunctive treatment modality showing an increased improvement in developing insight and awareness of one's condition.

Drugs, Alcohol, and the Hypothalamus

Stress, drugs, and alcohol have been intricately connected in preclinical models using rodents. Stress and its relationship to the HPA axis can lead to both the acquisition and the reinforcement using the reward pathway in both illicit drug and alcohol use.

Substance use, in terms of alcohol or other illicit drugs, can be a life-controlling condition. According to the centers for disease control, the prevalence of illicit drug use in people aged 12 and over is 10.6%. The prevalence of adults who have had at least one heavy drinking day, as defined by five or more drinks for males and four or more drinks for females, is 25.1%. What makes some individuals more prone to drug and alcohol use and/or dependency? Preclinical data suggest that chronic exposure to stress and the resulting release of glucocorticoids via the HPA axis sensitizes dopaminergic response to some drugs. There could be an interplay of stress and drug-induced activation of this axis that allows glucocorticoids to sensitize the reward pathways [39, 40]. When first exposed to drugs, an individual experiences pleasurable feelings. As this progresses to the addicted state, the HPA axis is activated, and high levels of glucocorticoids and stress peptides, which are associated with addiction, create an internal form of stress, which is seen as anxiety-like behaviors.

The pleasurable state of drug intoxication is a positive reinforcer for future use. These reinforcing effects act through signal transduction systems involving the mesocorticolimbic dopaminergic neurons [41]. This leads to increased dopamine expression in the nucleus accumbens. There is evidence that the HPA axis interacts with the mesocorticolimbic dopaminer-

gic system. Preclinical studies have shown that glucocorticoids alter the dopamine signaling in the mesolimbic system and can thereby amplify the positive reinforcing effects of alcohol and illicit drugs [42]. This is known as sensitization of the reward system, which could lead to the user being more responsive to drugs or alcohol and therefore be more vulnerable to the development of addiction [42]. Studies have shown that corticotropin-releasing factor (CRF), released from the paraventricular nucleus of the hypothalamus, as well as the HPA axis end product of glucocorticoids, has led to the increased glutamatergic activity in the ventral tegmental area (VTA). VTA axons extend to the nucleus accumbens resulting in an increase in dopamine release. CRF administration alone can produce the same results [43]. Several studies have shown that stress and corticosterone, as well as substances such as cocaine, methamphetamine, morphine, nicotine, and alcohol, can sensitize VTA dopamine neurons to increase their glutamatergic input [44, 45].

Animal models show that during the acquisition phase of drug-taking behavior, the animal will come into contact with the drug and its rewarding properties for the first time [46]. Stressors applied to rats increase the serum glucocorticoid levels by activation of the HPA axis, ultimately altering the drug self-administration. Tail pinch [39], neonatal isolation [47], and foot shock [48] are examples of stressors that can increase the drug acquisition of stimulants like cocaine and amphetamine and self-administration of opiates. Shaham and Stewart also showed that electric foot shock increases the reinforcing effect of heroin. Researchers further investigated this process by administering the final product of the HPA axis, corticosterone (analogous to cortisol in humans). Increased low-dose cocaine self-administration was observed in foot shock-induced rats with increases in the corticosterone levels [46]. Even more interesting is the fact that self-administration did not occur unless the plasma levels reached a certain threshold of corticosterone [46]. A study also showed that corticosterone administration had been shown to facilitate the psychomotor stimulant effects of both cocaine and morphine [49]. In a related study,

Goeders and Guerin showed that adrenalectomized rats, which eliminated the final step in the HPA axis, actually did not self-administer cocaine at any dose they tested [50]. Pretreatment with corticosterone facilitates the cocaine self-administration in rats [51]. This could be reversed in a dose-dependent manner with the administration of corticosterone [52].

Using an escalation model for cocaine self-administration associated with an increase in reward thresholds, Koob and Kreck reported a positive correlation between pre-session corticosterone levels and the amount of cocaine that was self-administered [53]. In a related experiment, the blockage of corticosterone synthesis with ketoconazole reduced low-dose self-administration of cocaine during acquisition [54]. Plasma corticosterone levels prior to self-administration predicted cocaine self-administration, but this was only observed at low doses of cocaine (0.25 mg/kg) [55]. This indicates that individual differences in cocaine self-administration that occur during the acquisition period are only relevant at low doses of cocaine. Also, during acquisition, cocaine administration induces the activation of the HPA axis, which leads to increased levels of corticosterone in rats. This cocaine-induced increase in corticosterone in the plasma may result from the effects of the drug on CRF secretion in the hypothalamus [56]. With this in mind, it seems reasonable to deduce that low-dose psychostimulant effects may be related to the initial vulnerability to drug use, and the activation of the HPA axis may contribute to this vulnerability. Cocaine use not only stimulates the HPA axis but also may act as a positive feedback loop to its continued use.

Alcohol Consumption and the Hypothalamus

Neuropeptides play a role in both the consumption of alcohol and to encourage increased intake of alcohol. Galatin, enkephalin, and orexin work to enhance alcohol consumption in various ways and to act in a positive feedback loop in terms of consumption [57].

Galatin (GAL) is found in the paraventricular nucleus, arcuate nucleus, lateral hypothalamus, dorsomedial nucleus, and periventricular area in rats. GAL interacts with three different G-protein receptors. In the paraventricular nucleus, GAL promotes alcohol consumption. Injection of GAL into this area in rats has shown to increase moderate alcohol drinking [58]. This is analogous to the way that GAL seems to trigger increased food intake when interacting in the paraventricular nucleus, which this study also noted. There is also evidence that endogenous GAL has a role in promoting alcohol drinking as well. Barson showed that rats with elevated mRNA expression of GAL were correctly predicted to drink higher levels of alcohol [59]. Furthermore, mice overexpressing the GAL gene were shown to drink more alcohol and show a higher preference for alcohol than their wild-type peers [60]. Alternatively, those who were lacking the GAL gene drank less alcohol and showed a lower preference for it [61]. GAL also has a role in the reward pathway. Increased levels of GAL in the paraventricular nucleus have been found to increase extracellular levels of dopamine in the nucleus accumbens [62]. There is also evidence that supports that GAL has a positive feedback relationship with alcohol. Leibowitz et al. showed that alcohol at moderate and excessive levels led to increased gene expression and peptide levels of GAL in multiple areas of the hypothalamus [63].

Enkephalin (ENK), an endogenous opioid, is another neuropeptide widely disturbed throughout the brain. ENK is active in the paraventricular nucleus, arcuate nucleus, lateral hypothalamus, dorsomedial nucleus, and perifornical area. ENK can promote moderate alcohol drinking through its actions on the paraventricular nucleus. It has been shown to have similar actions as GAL in terms of promoting alcohol drinking when active in the paraventricular nucleus; however, it differs from GAL in also increasing the duration of those drinking episodes [58]. This study suggests that not only do rats drink more with elevated endogenous ENK but that they also drink for a longer period of time. Unlike mice lacking GAL, mice lacking the ENK gene showed no difference from their wild-type controls in either their alco-

hol preference or intake [64] and no increase in alcohol drinking in all individuals under any condition. However, ENK does have a role in the reward pathway, which could promote increased drinking behaviors. It does this by its actions on the delta-opioid receptor (DOR), which increases dopamine in the nucleus accumbens and thereby has a role in the positive reinforcement of drinking. A DOR agonist injected into the paraventricular nucleus was shown to increase extracellular levels of dopamine in the nucleus accumbens [65].

Orexin (OX) is found in cell bodies that lie exclusively within the hypothalamus allowing the effects of OX to be traced back to the actions of neurons within the hypothalamus. Projections of these cell bodies lie in various parts of the brain such as the thalamus, locus coeruleus, nucleus accumbens, ventral tegmental area, amygdala, and various nuclei of the hypothalamus which shows that OX has effects throughout the brain. Generally, OX is thought to promote alcohol consumption, but this does not happen in all brain areas. Injection of OX-A into the paraventricular or lateral hypothalamus of rats stimulates moderate alcohol consumption [58]. Outside of the hypothalamus, OX promotes excessive alcohol consumption in the nucleus accumbens core but not the shell [66], the ventral tegmental area but not the substantia nigra, and the anterior but not posterior paraventricular thalamus [67]. OX is different from ENK and GAL, where the latter stimulates alcohol drinking by increasing the amount and duration of alcohol drinking episodes and the former in the hypothalamus stimulates alcohol drinking by increasing the number of drinking episodes [58]. This suggests that rats with elevated OX initiate more drinking responses. Additionally, scenarios that induce reinstatement of alcohol-seeking have been found to stimulate c-Fos activity within OX neurons [68]. Also, rats predicted to drink higher levels of alcohol based on their novelty-induced locomotor activity showed elevated OX mRNA levels in the perifornical LH [59]. However, there is some evidence that shows that OX levels are the same in alcohol-preferring rats compared to non-preferring rats [69]. Evidence that offers a

possible explanation shows that while OX on its own is not necessary for the renewal of alcohol-seeking behaviors, it may work in concert with other neuropeptides to induce this behavior [70]. In terms of the reward system, OX is responsive to conditions associated with rewards. Within the hypothalamus, OX neurons show c-Fos activation in response to condition contextual cues associated with sexual behavior, which is a natural reward [71]. This study also showed that lesions of OX neurons reduced the conditioned place preference for a sexual paired behavior chamber. These effects of OX are attributed to the actions at the OX2R receptor rather than the OX1R since a peripheral injection of an OX2R antagonist reduced the acquisition, expression, and reinstatement of alcohol-conditioned place preference [72], whereas an injection of an OX1R antagonist does not affect the acquisition and only weakly reduces the expression of alcohol-conditioned place preference. There is also a role in positive reinforcement of OX. Studies report that alcohol increases gene expression of OX within the lateral hypothalamus [67, 69]. Chronic alcohol injection in mice also leads to a small increase in c-fos activity in OX neurons [73]. Overall, this indicates that OX can help promote alcohol consumption, while alcohol consumption itself, in turn, increases levels of OX in a positive feedback loop.

There are three other neuropeptides that are associated with the hypothalamus that have a negative effect on alcohol consumption. These are dynorphin (DYN), corticotropin-releasing factor (CRF), and melanocortins (MCs). These neuropeptides have a negative effect on drinking by causing dysphoria, suppress drinking overall, or cause disruptions in the reward pathway.

Dynorphin (DYN) is found throughout the brain but is most densely expressed within the hypothalamus itself with cell bodies located within the paraventricular nucleus, arcuate nucleus, dorso-medial nucleus, lateral hypothalamus, and ventromedial hypothalamus. DYN in the lateral hypothalamus is highly co-expressed with OX, and the two are found packaged within the same synaptic vesicles in the hypothalamus. DYN exhibits kappa opioid receptor (KOR) activity

resulting in inhibiting alcohol consumption. This was proved when a KOR agonist injected to the lateral hypothalamus or paraventricular nucleus suppressed moderate alcohol drinking [74]. One theory is that KOR activity in the lateral hypothalamus induces a conditioned place aversion and subsequent dysphoria [75]. Alcohol consumption itself also stimulates DYN production. It has been shown that alcohol drinking at moderate levels can lead to elevated DYN gene expression in the paraventricular nucleus and that increase can be observed with just a single alcohol injection [76]. This release is thought to curb the subsequent alcohol consumption and counter the drive for further intake that is induced by the simultaneously released OX peptide.

Corticotropin-releasing factor (CRF) is found densely in the extending amygdala and also in the hypothalamus predominantly in the paraventricular nucleus and the lateral hypothalamus. CRF shows a negative feedback relationship with alcohol. Injection into the lateral ventricles has found to reduce moderate and excessive drinking in rats [77] though when injected into the paraventricular nucleus, it has no effect on excessive alcohol drinking [78]. Additionally, levels of CFR are decreased in the hypothalamus of alcohol-dependent rats at the onset of withdrawal [79]. This is when animals are presumed to crave alcohol, which further supports the theory that CFR helps reduce the drive for alcohol consumption. CFR and DYN are similar in that the reduced alcohol consumption could be due to their ability to induce aversion. Ventricular injection of a CRF agonist will elevate the intracranial self-stimulation threshold, which indicates that it reduces brain stimulation in terms of reward [80]. This, therefore, indicates that CRF can mediate some of the aversive aspects of alcohol consumption and can serve to inhibit subsequent alcohol drinking.

Melanocortin (MC) peptides are derived from proopiomelanocortin (POMC), which includes other components including alpha-melanocyte-stimulating hormone, β -MSH, gamma-MSH, and adrenocorticotrophic hormone. Cell bodies containing MC are located primarily in the hypothalamus, especially in the arcuate nucleus [81].

Projections of the cell bodies terminate in all areas of the hypothalamus as well as in a variety of other brain areas such as the amygdala, periaqueductal gray, and septum. MC shows a double-negative feedback loop in relation to alcohol, which generally is noted in decreasing alcohol consumption and also being endogenously reduced by alcohol. Injection of an MC receptor agonist in the nucleus accumbens, ventral tegmental area, or amygdala showed a reduction in moderate and excessive alcohol consumption [82]. Other studies have shown that this also occurs in the nuclei of the medial hypothalamus, such as the paraventricular nucleus and the dorsomedial nucleus. The ability of MC to inhibit alcohol consumption may be due to the suppression of reward or reinforcement from this behavior. Ventricular injection of alpha-MSH reduces the anxiolytic effect of excessive alcohol consumption, as demonstrated with an experiment using an elevated plus maze [83]. This study also showed that an MC4R antagonist enhanced the anxiolytic effect and reduced the anxiety associated with alcohol withdrawal. Furthermore, an MC4R agonist in the nucleus accumbens reduced the hedonic reactions to alcohol while increasing aversive ones [84]. Therefore, MCs, like CFR, may enhance the aversive aspects of alcohol consumption while also reducing the rewarding effects. Unlike other neuropeptides previously listed, endogenous MC is generally inhibited by alcohol. Peptide levels of POMC in the arcuate nucleus are decreased after long-term exposure to high levels of alcohol via a vapor chamber or alcohol-containing diet [85, 86]. Peptide levels of alpha-MSH after long-term consumption of alcohol-containing diet are also decreased in the arcuate nucleus and possibly in the lateral hypothalamus and in certain limbic regions, including the amygdala and substantia nigra [85]. With this double-negative feedback loop, the reduction of MC levels with alcohol consumption may, in fact, be permissive to further alcohol consumption by reducing the aversive aspects of its intake.

The hypothalamus plays a central role in terms of substance abuse. Its involvement in the HPA axis in stress reactions can prime the use of a sub-

stance. The hypothalamus can also potentiate substance use since it is involved in the reward pathway. This priming of the use and positive feedback loop can have negative consequences for individuals who have been exposed to life stressors such as abuse, neglect, or early trauma. Practitioners must be aware of early stressors and address them with appropriate treatment if substance use treatment is to be successful. Simply removing a person from the substance may not be enough to stop their use in the long term. Treatment must address their behaviors, thoughts, and emotions.

Conclusion

The hypothalamus plays an integral role in the body's response to physiological as well as emotional stress. Chronic exposure to such stressors can have lasting effects on these individuals, including predisposing them for risk of various psychiatric disorders. Pharmacological treatment interventions may be effective by altering either hypothalamic function or the consequence of its dysfunction [32, 87].

References

1. Selye H. A syndrome produced by diverse nocuous agents. 1936. *J Neuropsychiatry Clin Neurosci*. 1998;10(2):230–1.
2. Barbosa DAN, de Oliveira-Souza R, Monte Santo F, de Oliveira Faria AC, Gorgulho AA, De Salles AAF. The hypothalamus at the crossroads of psychopathology and neurosurgery. *Neurosurg Focus*. 2017;43(3):E15.
3. Stemson SM. Hypothalamic survival circuits: blueprints for purposive behaviors. *Neuron*. 2013;77(5):810–24.
4. Pascual JM, Prieto R, Castro-Dufourny I, Mongardi L, Rosdolsky M, Strauss S, et al. Craniopharyngiomas primarily involving the hypothalamus: a model of neurosurgical lesions to elucidate the neurobiological basis of psychiatric disorders. *World Neurosurg*. 2018;120:e1245–e78.
5. Angelergues R, De Ajuriaguerra J, Hecaen H. [Paralysis of the upward movement of the eyes of vascular origin; anatomoclinical study]. *Rev Neurol (Paris)*. 1957;96(4):301–19.
6. Paredes RG, Tzschentke T, Nakach N. Lesions of the medial preoptic area/anterior hypothalamus (MPOA/AH) modify partner preference in male rats. *Brain Res*. 1998;813(1):1–8.
7. Calabro RS, Cacciola A, Bruschetta D, Milardi D, Quattrini F, Sciarrone F, et al. Neuroanatomy and function of human sexual behavior: A neglected or unknown issue? *Brain Behav*. 2019;9:e01389.
8. Fuss J, Auer MK, Biedermann SV, Briken P, Hacke W. Deep brain stimulation to reduce sexual drive. *J Psychiatry Neurosci*. 2015;40(6):429–31.
9. Sano K, Mayanagi Y. Posteromedial hypothalamotomy in the treatment of violent, aggressive behaviour. *Acta Neurochir Suppl (Wien)*. 1988;44:145–51.
10. Schwarcz JR, Driollet R, Rios E, Betti O. Stereotactic hypothalamotomy for behaviour disorders. *J Neurol Neurosurg Psychiatry*. 1972;35(3):356–9.
11. Cleary DR, Ozpinar A, Raslan AM, Ko AL. Deep brain stimulation for psychiatric disorders: where we are now. *Neurosurg Focus*. 2015;38(6):E2.
12. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. Fifth ed. Arlington: American Psychiatric Association; 2013.
13. Eiland L, Romeo RD. Stress and the developing adolescent brain. *Neuroscience*. 2013;249:162–71.
14. Malikowska-Racia N, Salat K. Recent advances in the neurobiology of posttraumatic stress disorder: a review of possible mechanisms underlying an effective pharmacotherapy. *Pharmacol Res*. 2019;142:30–49.
15. Fischer S, Cleare AJ. Cortisol as a predictor of psychological therapy response in anxiety disorders-systematic review and meta-analysis. *J Anxiety Disord*. 2017;47:60–8.
16. Fischer S, Ehler U. Hypothalamic-pituitary-thyroid (HPT) axis functioning in anxiety disorders. A systematic review. *Depress Anxiety*. 2018;35(1):98–110.
17. Naughton M, Dinan TG, Scott LV. Corticotropin-releasing hormone and the hypothalamic-pituitary-adrenal axis in psychiatric disease. *Handb Clin Neurol*. 2014;124:69–91.
18. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003;289(23):3095–105.
19. Souery D, Papakostas GI, Trivedi MH. Treatment-resistant depression. *J Clin Psychiatry*. 2006;67(Suppl 6):16–22.
20. Juruena MF. Early-life stress and HPA axis trigger recurrent adulthood depression. *Epilepsy Behav*. 2014;38:148–59.
21. Owens MJ, Nemeroff CB. The role of corticotropin-releasing factor in the pathophysiology of affective and anxiety disorders: laboratory and clinical studies. *Ciba Found Symp*. 1993;172:296–308; discussion -16.
22. Price JL, Drevets WC. Neurocircuitry of mood disorders. *Neuropsychopharmacology*. 2010;35(1):192–216.

23. Pariente CM. The glucocorticoid receptor: part of the solution or part of the problem? *J Psychopharmacol.* 2006;20(4 Suppl):79–84.
24. Ida T, Nakahara K, Murakami T, Hanada R, Nakazato M, Murakami N. Possible involvement of orexin in the stress reaction in rats. *Biochem Biophys Res Commun.* 2000;270(1):318–23.
25. Salomon RM, Ripley B, Kennedy JS, Johnson B, Schmidt D, Zeitzer JM, et al. Diurnal variation of cerebrospinal fluid hypocretin-1 (Orexin-A) levels in control and depressed subjects. *Biol Psychiatry.* 2003;54(2):96–104.
26. Juruena MF, Bocharova M, Agustini B, Young AH. Atypical depression and non-atypical depression: is HPA axis function a biomarker? A systematic review. *J Affect Disord.* 2018;233:45–67.
27. Maguire J. Neuroactive steroids and GABAergic involvement in the neuroendocrine dysfunction associated with major depressive disorder and postpartum depression. *Front Cell Neurosci.* 2019;13:83.
28. Dinsdale NL, Crespi BJ. Revisiting the wandering womb: oxytocin in endometriosis and bipolar disorder. *Horm Behav.* 2017;96:69–83.
29. Crespi BJ. Oxytocin, testosterone, and human social cognition. *Biol Rev Camb Philos Soc.* 2016;91(2):390–408.
30. Lien YJ, Chang HH, Tsai HC, Kuang Yang Y, Lu RB, See CP. Plasma oxytocin levels in major depressive and bipolar II disorders. *Psychiatry Res.* 2017;258:402–6.
31. Cullinan WE, Ziegler DR, Herman JP. Functional role of local GABAergic influences on the HPA axis. *Brain Struct Funct.* 2008;213(1–2):63–72.
32. Schule C, Eser D, Baghai TC, Nothdurfter C, Kessler JS, Rupprecht R. Neuroactive steroids in affective disorders: target for novel antidepressant or anxiolytic drugs? *Neuroscience.* 2011;191:55–77.
33. National Institute of Mental Health (U.S.). Schizophrenia. [Bethesda, MD]: National Institute of Mental Health, U.S. Dept. of Health and Human Services, National Institutes of Health; 2009. Available from: <http://utils.louislibraries.org/cgi-bin/lz0050.x?sitecode=LASH?http://purl.access.gpo.gov/GPO/LPS123332>.
34. Bernstein H-G, Dobrowolny H, Bogerts B, Keilhoff G, Steiner J. The hypothalamus and neuropsychiatric disorders: psychiatry meets microscopy. *Cell Tissue Res.* 2019;375(1):243–58.
35. Du X, Hill RA. Hypothalamic-pituitary-gonadal axis dysfunction: an innate pathophysiology of schizophrenia? *Gen Comp Endocrinol.* 2019;275:38–43.
36. Altamura AC. Hypothalamic-pituitary-adrenal axis in schizophrenia. *Biol Psychiatry.* 1996;40. United States:560–1.
37. Chien YL, Liu CM, Shan JC, Lee HJ, Hsieh MH, Hwu HG, et al. Elevated plasma orexin A levels in a subgroup of patients with schizophrenia associated with fewer negative and disorganized symptoms. *Psychoneuroendocrinology.* 2015;53:1–9.
38. Borgland SL, Labouebe G. Orexin/hypocretin in psychiatric disorders: present state of knowledge and future potential. *Neuropsychopharmacology.* 2010;35(1):353–4.
39. Piazza PV, Le Moal M. The role of stress in drug self-administration. *Trends Pharmacol Sci.* 1998;19(2):67–74.
40. Tidey JW, Miczek KA. Acquisition of cocaine self-administration after social stress: role of accumbens dopamine. *Psychopharmacology (Berl).* 1997;130(3):203–12.
41. Uhart M, Wand GS. Stress, alcohol and drug interaction: an update of human research. *Addict Biol.* 2009;14(1):43–64.
42. Marinelli M, Piazza PV. Interaction between glucocorticoid hormones, stress and psychostimulant drugs. *Eur J Neurosci.* 2002;16(3):387–94.
43. Ungless MA, Singh V, Crowder TL, Yaka R, Ron D, Bonci A. Corticotropin-releasing factor requires CRF binding protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. *Neuron.* 2003;39(3):401–7.
44. Overton PG, Tong ZY, Brain PF, Clark D. Preferential occupation of mineralocorticoid receptors by corticosterone enhances glutamate-induced burst firing in rat midbrain dopaminergic neurons. *Brain Res.* 1996;737(1–2):146–54.
45. Saal D, Dong Y, Bonci A, Malenka RC. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron.* 2003;37(4):577–82.
46. Goeders NE. Stress and cocaine addiction. *J Pharmacol Exp Ther.* 2002;301(3):785–9.
47. Kosten T, Miserendino M, Kehoe P. Enhanced acquisition of cocaine self-administration in adult rats with neonatal isolation stress experience. *Brain Res.* 2000;875(1–2):44–50.
48. Shaham Y, Stewart J. Exposure to mild stress enhances the reinforcing efficacy of intravenous heroin self-administration in rats. *Psychopharmacology (Berl).* 1994;114(3):523–7.
49. Marinelli M, Piazza PV, Deroche V, Maccari S, Le Moal M, Simon H. Corticosterone circadian secretion differentially facilitates dopamine-mediated psychomotor effect of cocaine and morphine. *J Neurosci.* 1994;14(5 Pt 1):2724–31.
50. Goeders N, Guerin G. Effects of surgical and pharmacological adrenalectomy on the initiation and maintenance of intravenous cocaine self-administration in rats. *Brain Res.* 1996;722(1–2):145–52.
51. Mantsch J, Saphier D, Goeders N. Corticosterone facilitates the acquisition of cocaine self-administration in rats: opposite effects of the type II glucocorticoid receptor agonist dexamethasone. *J Pharmacol Exp Ther.* 1998;287(1):72–80.
52. Deroche V, Marinelli M, Le Moal M, Piazza P. Glucocorticoids and behavioral effects of psychostimulants. II: cocaine intravenous self-administration and reinstatement depend on glucocorticoid levels. *J Pharmacol Exp Ther.* 1997;281(3):1401–7.

53. Koob G, Kreek MJ. Stress, dysregulation of drug reward pathways, and the transition to drug dependence. *Am J Psychiatry*. 2007;164(8):1149–59.
54. Goeders N, Peltier R, Guerin G. Ketoconazole reduces low dose cocaine self-administration in rats. *Drug Alcohol Depend*. 1998;53(1):67–77.
55. Mantsch J, Ho A, Schlussman S, Kreek M. Predictable individual differences in the initiation of cocaine self-administration by rats under extended-access conditions are dose-dependent. *Psychopharmacology (Berl)*. 2001;157(1):31–9.
56. Sarnyai Z, Shaham Y, Heinrichs S. The role of corticotropin-releasing factor in drug addiction. *Pharmacol Rev*. 2001;53(2):209–43.
57. Barson J, Leibowitz S. Hypothalamic neuropeptide signaling in alcohol addiction. *Prog Neuropsychopharmacol Biol Psychiatry*. 2016;65:321–9.
58. Chen Y, Barson J, Chen A, Hoebel B, Leibowitz S. Hypothalamic peptides controlling alcohol intake: differential effects on microstructure of drinking bouts. *Alcohol*. 2014;48(7):657–64.
59. Barson J, Fagan S, Chang G, Leibowitz S. Neurochemical heterogeneity of rats predicted by different measures to be high ethanol consumers. *Alcohol Clin Exp Res*. 2013;37 Suppl 1(0 1):E141–51.
60. Karatayev O, Leibowitz J, Leibowitz SF. Increased intake of ethanol and dietary fat in galanin overexpressing mice. *Alcohol*. 2009;43(8):571–80.
61. Karatayev O, Baylan J, Weed V, Chang S, Wynick D, Leibowitz S. Galanin knockout mice show disturbances in ethanol consumption and expression of hypothalamic peptides that stimulate ethanol intake. *Alcohol Clin Exp Res*. 2010;34(1):72–80.
62. Rada P, Mark G, Hoebel B. Galanin in the hypothalamus raises dopamine and lowers acetylcholine release in the nucleus accumbens: a possible mechanism for hypothalamic initiation of feeding behavior. *Brain Res*. 1998;798(1-2):1–6.
63. Leibowitz S, Avena N, Chang G, Karatayev O, Chau D, Hoebel B. Ethanol intake increases galanin mRNA in the hypothalamus and withdrawal decreases it. *Physiology & Behavior*. 2003;79:103–11.
64. Koenig H, Olive M. Ethanol consumption patterns and conditioned place preference in mice lacking preproenkephalin. *Neurosci Lett*. 2002;325(2):75–8.
65. Rada P, Barson J, Leibowitz S, Hoebel B. Opioids in the hypothalamus control dopamine and acetylcholine levels in the nucleus accumbens. *Brain Res*. 2010;1312:1–9.
66. Brown R, Khoo S, Lawrence A. Central orexin (hypocretin) 2 receptor antagonism reduces ethanol self-administration, but not cue-conditioned ethanol-seeking, in ethanol-preferring rats. *Int J Neuropsychopharmacol*. 2013;16(9):2067–79.
67. Barson J, Fagan S, Chang G, Leibowitz S. Anterior thalamic paraventricular nucleus is involved in intermittent access ethanol drinking: role of orexin receptor 2. *Addict Biol*. 2015;20(3):469–81.
68. Dayas C, McGranahan T, Martin-Fardon R, Weiss F. Stimuli linked to ethanol availability activate hypothalamic CART and orexin neurons in a reinstatement model of relapse. *Biol Psychiatry*. 2008;63(2):152–7.
69. Lawrence A, Cowen M, Yang H, Chen F, Oldfield B. The orexin system regulates alcohol-seeking in rats. *Br J Pharmacol*. 2006;148(6):752–9.
70. Prasad A, McNally G. Effects of vivo morpholino knockdown of lateral hypothalamus orexin/hypocretin on renewal of alcohol seeking. *PLoS One*. 2014;9(10):e110385.
71. Di Sebastiano A, Wilson-Pérez H, Lehman M, Coolen L. Lesions of orexin neurons block conditioned place preference for sexual behavior in male rats. *Horm Behav*. 2011;59(1):1–8.
72. Shoblock J, Welty N, Aluisio L, Fraser I, Motley S, Morton K, et al. Selective blockade of the orexin-2 receptor attenuates ethanol self-administration, place preference, and reinstatement. *Psychopharmacology (Berl)*. 2011;215(1):191–203.
73. Macedo G, Kawakami S, Vignoli T, Sinigaglia-Coimbra R, Suchecki D. The influence of orexins on ethanol-induced behavioral sensitization in male mice. *Neurosci Lett*. 2013;551:84–8.
74. Barson J, Carr A, Soun J, Sobhani N, Rada P, Leibowitz S, et al. Opioids in the hypothalamic paraventricular nucleus stimulate ethanol intake. *Alcohol Clin Exp Res*. 2010;34(2):214–22.
75. Bals-Kubik R, Ableitner A, Herz A, Shippenberg T. Neuroanatomical sites mediating the motivational effects of opioids as mapped by the conditioned place preference paradigm in rats. *J Pharmacol Exp Ther*. 1993;264(1):489–95.
76. Chang G, Karatayev O, Ahsan R, Avena N, Lee C, Lewis M, et al. Effect of ethanol on hypothalamic opioid peptides, enkephalin, and dynorphin: relationship with circulating triglycerides. *Alcohol Clin Exp Res*. 2007;31(2):249–59.
77. Thorsell A, Slawecki C, Ehlers C. Effects of neuropeptide Y and corticotropin-releasing factor on ethanol intake in Wistar rats: interaction with chronic ethanol exposure. *Behav Brain Res*. 2005;161(1):133–40.
78. Knapp D, Overstreet D, Huang M, Wills T, Whitman B, Angel R, et al. Effects of a stressor and corticotropin releasing factor on ethanol deprivation-induced ethanol intake and anxiety-like behavior in alcohol-preferring P rats. *Psychopharmacology (Berl)*. 2011;218(1):179–89.
79. Zorrilla E, Valdez G, Weiss F. Changes in levels of regional CRF-like-immunoreactivity and plasma corticosterone during protracted drug withdrawal in dependent rats. *Psychopharmacology (Berl)*. 2001;158(4):374–81.
80. Macey D, Koob G, Markou A. CRF and urocortin decreased brain stimulation reward in the rat: reversal by a CRF receptor antagonist. *Brain Res*. 2000;866(1-2):82–91.
81. Dubé D, Lissitzky J, Leclerc R, Pelletier G. Localization of alpha-melanocyte-stimulating

- hormone in rat brain and pituitary. *Endocrinology*. 1978;102(4):1283–91.
82. Lerma-Cabrera J, Carvajal F, de la Torre L, de la Fuente L, Navarro M, Thiele T, et al. Control of food intake by MC4-R signaling in the lateral hypothalamus, nucleus accumbens shell and ventral tegmental area: interactions with ethanol. *Behav Brain Res*. 2012;234(1):51–60.
83. Kokare D, Chopde C, Subhedar N. Participation of alpha-melanocyte stimulating hormone in ethanol-induced anxiolysis and withdrawal anxiety in rats. *Neuropharmacology*. 2006;51(3):536–45.
84. Lerma-Cabrera J, Carvajal F, Chotro G, Gaztañaga M, Navarro M, Thiele T, et al. MC4-R signaling within the nucleus accumbens shell, but not the lateral hypothalamus, modulates ethanol palatability in rats. *Behav Brain Res*. 2013;239:51–4.
85. Navarro M, Cubero I, Thiele T. Decreased immunoreactivity of the polypeptide precursor pro-opiomelanocortin (POMC) and the prohormone convertase pc1/3 after chronic ethanol exposure in Sprague-Dawley rats. *Alcohol Clin Exp Res*. 2013;37(3):399–406.
86. Scanlon M, Lazar-Wesley E, Grant K, Kunos G. Proopiomelanocortin messenger RNA is decreased in the mediobasal hypothalamus of rats made dependent on ethanol. *Alcohol Clin Exp Res*. 1992;16(6):1147–51.
87. Bradley AJ, Dinan TG. A systematic review of hypothalamic-pituitary-adrenal axis function in schizophrenia: implications for mortality. *J Psychopharmacol*. 2010;24(4 Suppl):91–118.

Neurosurgical Aspects of Hypothalamic Disease

9

Ketan R. Bulsara, Joshua Knopf, Rebecca Calafiore,
and Anzhela D. Moskalik

Anatomy of the Hypothalamus

The hypothalamus, anatomically, poses many difficulties for a neurosurgeon. The hypothalamus is situated beneath the thalamus, separated by the hypothalamic sulcus of Monro – formed by the lateral wall of the third ventricle, extending anteroposteriorly from the foramina of Monro to the cerebral aqueduct. Anteriorly, the hypothalamus is bordered by the lamina terminalis. Superiorly, the inferolateral wall of the third ventricle and the fornix form the upper limits to the hypothalamus. Posteriorly, the hypothalamus extends to the tegmentum of the midbrain. The lateral wall of the hypothalamus extends to the optic tract(s) [19].

K. R. Bulsara (✉)

Department of Surgery, Division of Neurosurgery,
University of Connecticut Health Center, UConn
Health, Farmington, CT, USA
e-mail: bulsara@uchc.edu

J. Knopf

School of Medicine, UConn Health,
Farmington, CT, USA
e-mail: knopf@uchc.edu

R. Calafiore

School of Medicine, University of Connecticut
Health Center, Farmington, CT, USA
e-mail: rcalafiore@uchc.edu

A. D. Moskalik

University of Connecticut School of Medicine,
Farmington, CT, USA
e-mail: moskalik@uchc.edu

The hypothalamus itself is divided into eight nuclei, each with their own specific function. The supraoptic, dorsomedial, and paraventricular nuclei sit in the anterior aspect of the hypothalamus, dubbed the “supraoptic area,” as this portion of the hypothalamus sits above the optic chiasm and the pituitary gland. The central hypothalamus consists of the paraventricular, dorsomedial, arcuate, and ventromedial nuclei. The posterior aspect of the pituitary contains the mammillary bodies/nuclei, along with the posterior hypothalamic nuclei [19].

Clearly, the hypothalamus sits in an incredibly dense area of the brain, surrounded on all sides by eloquent structures. Surgical approach to these lesions can be incredibly difficult and requires advanced surgical techniques and approaches. We will discuss common diseases of the hypothalamus requiring surgical intervention, as well as the well-reported surgical approaches to treat these lesions.

Hypothalamic Diseases Requiring Surgical Management

Hypothalamic Hamartoma

A hamartoma is a benign noncancerous mass that consists of disorganized growth of native tissue. Although considered a tumor, this lesion rarely exhibits neoplastic characteristics and/or tendencies (i.e., no accelerated growth, no local invasion

of surrounding area(s), nor do hamartomas spread). These lesions consist of a mix of hyperactive neuronal cells and glial cells within the hypothalamus, usually arising from the floor of the third ventricle [25, 31]. Morphologically, there are two main subtypes: pedunculated and sessile hypothalamic hamartomas – the chief difference being the normal location of the lesion. Pedunculated lesions stem from a common stalk and extend into the third ventricle, whereas the sessile form grows entirely within the hypothalamus [25].

These are associated with a number of endocrine abnormalities as well as gelastic epilepsy. Additionally, these people may suffer from behavioral, psychiatric, and cognitive impairments causing developmental delay in childhood and can complicate therapeutic approach to these patients [16].

The diagnosis is made through clinical symptoms of endocrine abnormalities, precocious puberty, or epileptic symptoms. Appropriate workup includes relevant endocrine labs (electrolytes, pituitary hormones, etc.), EEG, and MR imaging studies. It is important to note that EEG studies can often fail to show ictal/interictal abnormalities (gelastic seizures can present without these findings).

Surgical intervention has limited evidence-based management data due to the novelty of the disease. Common practice and relevant case-report literature indicate that surgical intervention is considered in patients with uncontrollable, medication-refractory seizures in the setting of a known hypothalamic hamartoma. Case reports identify the standard pterional approach, transcallosal approach, and subfrontal translaminar terminalis approach as common surgical approaches to these lesions [21, 31]. Intraoperative guidelines and reports suggest aggressive resection is not advised due to the delicate nature of resection within the region, as injury to the subfrontal region, anterior cerebral artery (and associated perforating arteries), and chiasmatic structures [26].

Pilocytic Astrocytoma

Pilocytic astrocytomas are widely known for being very common ventricular tumors, typically arising

within the posterior fossa (cerebellum), extending into the fourth ventricle. These tumors also can originate from the midbrain, thalamus, and hypothalamus – these tumors often extend into the third ventricle. These tumors have good prognosis (5-year survival 80–90%), likely attributable to the slow growth of the mass, which allows for complete surgical resection. Surgery is a staple for effective management of these tumors [23].

Although the majority of pilocytic astrocytomas are found along the fourth ventricle, these tumors can also be found along the third ventricle. The hypothalamus can both complicate an appropriate surgical approach and be involved via direct extension of the tumor. Surgical approach to a third ventricular pilocytic astrocytoma remains relatively consistent regardless of the specific origin of the tumor. Commonly used surgical approaches include suprasellar transventricular approaches as well as a subfrontal translaminar approach or transcallosal approach [22]. The location of the astrocytoma within the third ventricle can affect which approach is chosen. It is essential to identify the hypothalamus when resecting these third ventricular-based lesions, as any that extend anteriorly or laterally to the ventricle risks involving the hypothalamus. Careless resection in these areas risks damaging the nearby, or potentially involved, hypothalamus. An important surgical note in these resections is that traditional practices include draining the cystic portion of the tumor prior to microsurgical resection/dissection, thereby opening additional room to resect and relieving pressure on nearby structures [22, 28].

Expansive Craniopharyngioma

Craniopharyngiomas exist as metaplastic adenohypophyseal cells, remnants of the hypophyseopharyngeal duct (Rathke's pouch). Embryologically, Rathke's pouch is a derivative of the ectoderm, exists as an invagination of the developing mouth, and eventually becomes the anterior pituitary gland and the pars distalis/tuberalis [11, 33]. During this process, cysts can develop during embryologic transformation within Rathke's pouch. In some people, these

cells persist and can proliferate/cause mass effect symptoms in the parasellar region.

The vast majority of craniopharyngiomas sit within the parasellar chiasmatic cisterns, with a minority (less than 5%) existing solely intraventricularly. These tumors are often caught once they grow to be very large, as the bulk of their growth extends into the chiasmatic cisterns and surrounding cisterns. This growth pattern causes large and expansive growth prior to affecting the surrounding parenchyma [24].

Histologically, craniopharyngiomas have two main subtypes: adamantinomatous and papillary subtypes. Adamantinomatous tumors are seen in young children and are cystic, classically known to contain cholesterol and dark cystic fluid. This variant is known to encase/surround numerous deep hemispheric structures, including the pituitary gland, optic chiasm, third ventricle, and hypothalamus. Papillary tumors, on the other hand, can be seen in both child and adult populations and are more solid and well-circumscribed [11, 17].

In patients with craniopharyngioma, evidence of neurocognitive changes best indicates involvement of the hypothalamus, as symptoms like endocrine dysfunction do not help physicians and surgeons differentiate between hypothalamic and pituitary function [17]. Common surgical approaches are transnasal or transcallosal, the decision between the two typically made by the location/size of the lesion. Should the tumor sit too far laterally, an endonasal approach could potentially not provide a wide-enough window to the full extent of the tumor, making maximum resection unlikely. Hypothalamic involvement of these tumors also complicates a surgical approach and can even call into question the need for an operation [33].

Typically, craniopharyngiomas respond extremely well to full resection; however involvement of the hypothalamus limits the surgeon's ability to be aggressive with surgical excision. In these patients, safe, yet incomplete, resection is more valued than gross total resection due to the efficacy of radiotherapy. Radiotherapy and radiosurgery have been found to protect vital structures while still adequately treating the tumor and decreasing morbidity and mortality [29].

Hypothalamic Optic Glioma/ Astrocytoma

Optic pathway and hypothalamic gliomas are rare astrocytic tumors, more commonly seen in children than adults. They comprise 3–5% of pediatric tumors (2% of brain tumors in all ages) and are commonly deemed unresectable. Common presenting symptoms are attributable to local interruption of visual/hypothalamic pathways – visual disturbances, endocrinopathies, hydrocephalus, and hypothalamic dysfunction [10].

To date, surgical resection of these lesions results in middle-term survival, tumor control, and functional outcome equivalent to that of chemotherapy – although the current fold standard treatment includes chemotherapy and as-needed debulking surgery.

Surgical treatment of these gliomas is controversial, likely because the rates of cure and successful treatment are wholly dependent on chemotherapy. Surgery is currently used when the tumor has exhibited growth following chemotherapy, or the size/location of the tumor permits surgical removal – unilaterality of the mass and involvement of only one optic pathway. Tumors localized on the optic chiasm or tumors extending into both optic pathways are deemed inoperable [34].

Common surgical approaches include an interhemispheric transcallosal, translaminar terminalis approach, transcortical, pterional, and, occasionally, an endonasal transsphenoidal approach. Maximal resection is critical, as these lesions commonly regrow and recur [3].

Deep Subcortical Arteriovenous Malformations

Arteriovenous malformations are aberrant connections between arterial and venous blood vessels and can cause numerous issues, such as headaches, seizures, stroke, and, potentially, hemorrhage. Deep subcortical AVMs can be a large headache for surgeons, due to the highly eloquent and functional parenchymal location and surgically difficult-to-reach areas [4].

Blood vessels to the hypothalamus have been implicated as spots for AVMs, both arising from the anterior and posterior circulation. Anterior circulation AVMs affecting the hypothalamus include the median lenticulostriate vessels (branching from the A1 anterior cerebral artery segment *or* the proximal M1 middle cerebral artery segment), which perfuse the suprachiasmatic hypothalamus, and the anterior thalamoperforators, arising from the M1 MCA segment. Posterior circulation AVMs affecting the hypothalamus are from the posterior thalamoperforators. Posterior thalamoperforating arteries are a set of small arteries arising from the proximal P1 segment of the posterior cerebral artery, affecting the posterior hypothalamus, thalamus, and medial midbrain [15].

These lesions are very difficult to approach, and guidelines suggest that resection should only be attempted if the lesion is small enough and the nidus of the AVM is both compact and lies along the ependymal surface [9]. Patients with these lesions who have no experienced hemorrhage nor preexisting neurologic deficits should not be operated on – the risk of the operation outweighs the risk of bleeding/deficits of such benign AVMs. Overaggressive surgical approach leaves the surgeon at risk for entering/rupturing the nidus, leaving the surgeon in a dangerous situation to stop torrential, life-threatening bleeding in a narrow operative window [27].

Should it be the best treatment option, surgical intervention and management of these lesions can best be approached via the transcortical approach or via a transcallosal transforaminal transvenous transchoroidal route. Following the medial wall of the third ventricle, the thalamoperforators, both anterior and posterior, can be identified at the inferior margins [15].

Supratentorial Cavernous Malformation

Cavernous malformations are clusters of small, abnormal blood vessels that can displace neurologic tissue, filled with slow-moving/stagnant blood that are very fragile. These malformations

predominantly arise (80%) above the tentorium. The majority of these reside in the cortical and/or subcortical space, with a small percentage of these (5–10%) originating in the thalamic/hypothalamic region. The cavernous malformations in this area are commonly associated with endocrine and thalamic abnormalities and rarely cause changes in cognition [18].

Supratentorial CMs are typically incidental findings but can also present with seizures or progressive neurologic deficits (secondary to hemorrhage). This constellation of symptoms due to SCMs is especially seen in younger patients – those in their 20s and 30s [14].

The approach to treatment about these malformations is heavily debated [14, 20]. Studies have advocated for early/immediate operative intervention, citing better surgical outcomes for those who underwent surgery closer to the initial onset of any seizure activity. On the flip side, studies have also showed that the risk of hemorrhage per year is as low as 0.25% per person per year in patients with no prior history of hemorrhage, calling into question the risk/benefit ratio of an operation to remove these lesions, especially those deep in the hypothalamus.

Several pre-op evaluations are critical: these malformations need to be assessed for an A-V fistula via arteriogram, and a functional MRI may be necessary to assess for functionality within the malformation [18]. Intraoperative image guidance improvements have allowed for more effective excision of deep cavernous malformations, such as those within the hypothalamus. Gross total resection of these CMs is essential, as this effectively removes chance for future hemorrhage and recurrent/continuing seizures.

Histoplasmosis

Histoplasmosis capsulatum is a dimorphic fungus commonly found in the midwestern United States. In immunocompromised states this fungus can disseminate to the brain and form histoplasmoses – small, expansile lesions within the hypothalamus, thalamus, or chiasmatic region [8]. These present on imaging as ring-enhancing

lesions and can cause local destruction and affect the function of these deep structures, resulting in hypothalamic dysfunction. These are very rare, as there are only ten reported cases of thalamic or hypothalamic involvement.

Surgical intervention in these patients has been documented, yet there are too few cases to come to consensus on whether surgical intervention is an effective treatment strategy [30]. For those who have underwent surgery for the fungal masses related to histoplasmosis, the literature suggests that most the cases that went to surgery were done via transcallosal or transocortical approach with great success [8, 30].

Surgical Approaches

Standard Pterional/Trans-sylvian Approach

A pterional approach, also known as a frontotemporal craniotomy, is a very commonly used approach for supratentorial lesions. It allows surgeons a wide range of variability and flexibility in the surgical approach to reach the anterior and middle skull base. From this opening, the surgeon has many options for surgical approach, most commonly trans-sylvian. The key steps of the pterional approach are positioning, skin incision, interfascial dissection, craniotomy, drilling of the sphenoid wing, and the dural opening. After the bone removal, the pterional approach proceeds with the opening of the sylvian fissure and the basal cisterns [32]. The trans-sylvian approach involves a small corticotomy with minimal extension to reach the hypothalamic region. Large landmarks with this approach include the internal carotid artery, the optic nerve/chiasm, and the infundibulum inferiorly. This lateral approach can be a large advantage for surgeons approaching lateral lesions [7].

This approach, however, has its limitations. First and foremost, this approach provides a somewhat narrow surgical window between the ICA, the optic nerve, and the infundibulum, limiting visualization and access of the third ventricle and medially placed surgical targets. Additionally, this exposure increases the risk for

additional morbidity – literature has reported increased rates transient third nerve palsy, post-op central diabetes insipidus, and hyperphagia with this approach [34]. Furthermore, Palmini et al. reported that 3 of their 13 patients achieved seizure freedom, and a > 90% reduction in seizure frequency was noted in the remaining 10 children [34].

When approaching hypothalamic lesions requiring surgical intervention that lie too far lateral for an endonasal or transventricular approach, a pterional approach with trans-sylvian approach to the lateral hypothalamus should be considered [1].

Supraorbital Craniotomy

In the mid-2000s, surgeons at the neurosurgery center in Helsinki Finland, most notably professor Juha Hernesniemi, were developing an alternative way to approach craniopharyngiomas, seeking to improve upon the standard pterional approach [1, 2]. Their efforts helped create and establish the lateral supraorbital approach to the parasellar, para-chiasmatic, and intra-sylvian regions. The lateral supraorbital approach involves an eyebrow incision with a minimally invasive keyhole approach to the frontal and frontal temporal region. The trajectory from this angle gives the surgeon a longer operative distance; however it simultaneously requires less brain manipulation and retraction. The benefit to this is that you have a window with many possible working angles to reach interior circulation aneurysms, school-based tumors, as well as parasellar, suprachiasmatic, and third ventricular lesions [1, 2]. Within this window lies the hypothalamus.

This approach is, naturally, not without its limitations. Approaching these deep structures from this angle requires corticectomy of the frontal lobe, which can lead to disinhibition and worsening cognitive processing and decision-making. Additionally, mediobasal injury can lead to amnesia, both anterograde and retrograde. Cosmetically, the incision around the eyebrow can lead to scarring. Similarly, should the incision run too far beyond the zygomatic process,

there is increased risk of injuring the frontal branch of the facial nerve and causing forehead weakness unilaterally (either temporarily or permanently) [7].

When compared directly with a pterional approach, the lateral supraorbital approach offers similar exposure while reducing surgical trauma and manipulation of the brain cortex and simplifying the surgical procedure [33].

Transcallosal Approach

A transcallosal approach involves approaching the hypothalamus/structures between the cerebral hemispheres, retracting each hemisphere laterally, and approaching the corpus callosum from above. The callosum is then split and the third ventricle, hypothalamus, thalamus, etc. all come into view. The patient is positioned with the dominant/affected hemisphere up, with the patient positioned such that the falx runs parallel to the floor. This positioning aids visualization and allows for avoidance of large frontal bridging veins and excellent visualization of the third ventricle, hypothalamus, mammillary bodies, and pituitary stalk beneath. This technique offers reduced rates of injury to the mammillary bodies, pituitary stalk, and optic chiasm. Furthermore, the minimal manipulation of the cerebral hemispheres and neurovascular structures within the suprasellar cisterns decrease the likelihood of cerebral infarction and oculomotor nerve palsy [7].

The inherent risks of the surgery involve the path to the hypothalamus – the septal and fornical injury can result in postsurgical memory deficits [12]. A 2003 study done by the Melbourne group analyzing complications following hypothalamic hamartoma resection using the transcallosal approach showed common complications, including thalamic infarction, increased appetite, and short-term memory deficits. Subsequent studies show an increased correlation between extent of resection and postoperative seizures [21]. Although these complications have been well-reported, their likelihood is low [21, 31].

Additional limitations to this approach include a restricted access and visualization to

laterally reaching tumors due to the presence of the pericallosal arteries. For medially sitting tumors, however, the transcallosal approach offers fantastic surgical access to this region with limited and infrequent long-term complications.

Subfrontal Translaminar Terminalis Approach

The lamina terminalis is a thin layer of gray matter extending backward from the corpus callosum (above the optic chiasm) and forms the median border of the third ventricle. Perhaps best known for its utility in third ventricle surgery, a subfrontal approach to the hypothalamus extending through this layer of the cortex involves superior retraction of the frontal lobe to expose the hypothalamus and third ventricle. The subfrontal translaminar terminalis approach offers excellent views and visualization of the lamina terminalis, optic nerves, optic chiasm, bilateral carotids, ACA segments, the hypothalamus, and the pituitary stalk [6, 7]. For anteriorly lying hypothalamic lesions, the lamina terminalis approach offers the surgeon great exposure within the surgical corridor. The largest advantage for this approach is the minimal manipulation of the cortex necessary to expose structures within the region of the hypothalamus. Additionally, virtually no manipulation of visual nerves is necessary, thereby reducing the risk of harm to these structures during surgery [6, 31].

The same reasons that make this approach great are also the reasons that make it problematic. The clean surgical window that provides great visualization of midline structures is also long, restricted, and non-flexible. The length of the approach limits the ability of the surgeon to extend laterally. The critical anatomy that sits just on the periphery of this approach makes extension beyond the surgical window very dangerous.

When compared to other approaches, the subfrontal terminalis approach is an excellent choice for a small subset of hyper-fire lesions, namely, those that sit close to the midline, whereas lesions that lay nearly anywhere else or

those that extend laterally are best operated on using a different approach.

Endoscopic Endonasal Approach

An endoscopic endonasal approach is a commonly used surgical approach to reach the sellar and suprasellar regions of the brain. This approach involves approaching the sellar and parasellar regions via drilling through the ethmoid sinus via nasal approach. This view has inline access and superior visualization of the pituitary stalk and hypothalamus. This visualization, traditionally used in removal of sellar/suprasellar craniopharyngiomas, offers excellent surgical access to midline lesions [13, 34].

The caveat with the endonasal approach, however, is that masses that extend laterally or posteriorly complicate resection. The laterality of the mass forces the surgeon to have to traverse the cavernous sinus and the cavernous portion of the carotid artery. This anatomy provides a dangerous road and can cause significant increases in morbidity. When these tumors remain midline and the whole capsule can be accessed readily, the endoscopic endonasal approach is preferred. For lesions that do not sit midline, other approaches will give better visualization.

Literature surrounding the capabilities of the endoscopic endonasal approach to lesions is growing. There are case reports and studies focused on the use of the endoscopic endonasal approach to not only pituitary lesions but also craniopharyngiomas (sellar, parasellar, hypothalamic, etc.), hypothalamic hamartomas, optic or hypothalamic gliomas, and meningiomas [31, 34]. Many of these lesions are traditionally operated on via a pterional subfrontal, supraorbital, or transcortical approach. This literature emphasizes that the endoscopic endonasal approach minimizes the manipulation of brain parenchyma or retraction; however it increases the likelihood of a postoperative CSF leak (into the sinus). Although many papers pointed to the increased risk of endocrinologic morbidity of this approach, no study had data or analysis to completely support this claim [31].

Additional Considerations to Surgery

When managing hypothalamic lesions, specifically operative lesions, there must be good discussion and preparation of the possible long-term effects of the operation. Injury to surrounding structures, loss of function of the hypothalamus, and morbidities surrounding intracranial surgery are all worth considering when weighing the risks and benefits of surgery on the hypothalamus. Limitations to surgery or surgical management, however, are not necessarily due to the risks of the surgery, but the circumstances surrounding an operation.

A solid foundational knowledge of the anatomy of the hypothalamus and surrounding structures is imperative for understanding the risks of surgery to this region. The visual pathways, including the optic chiasm, are structures that must be identified and avoided during surgery. Injury to the structures, naturally, will cause visual impairment and possible blindness. The classic presentation of this would be a bitemporal hemianopsia secondary to injury at the chiasm itself. Furthermore, injury to the seller region can cause pituitary dysfunction, as severe as panhypopituitarism due to destruction of the pituitary infundibulum. The close proximity of the hypothalamus to basal ganglia structures can lead to seizures, choreiform movements, involuntary tics, and spasticity, which have all been documented in the literature [5]. Vascular injury can cause several different, yet equally dangerous, outcomes. First and foremost, intraoperative damage to blood vessels can cause bleeding and related ischemic deficits. Additionally, hemorrhage can occur in the postoperative period, both epidural and subdural, dependent on the approach. Postoperative imaging should be completed both to record the extent of resection and also to assess for this undesired outcome [28].

The hypothalamus itself plays roles in numerous bodily functions, namely, multiple endocrine axes, appetite, sleep-wake cycle, behavior, memory, and emotion. The hypothalamus' close relationship with the pituitary gland via endocrine signaling is the source of many side effects and

morbidity surrounding hypothalamic surgery. Many of these complications arise with or are already complications of the lesion itself, as the hypothalamic process disrupts normal hypothalamic function. Manifestations of this include thyroid abnormalities, Cushing syndrome, diabetes insipidus, sexual dysfunction, growth abnormalities, and amenorrhea [5, 7, 19]. Additionally, the hypothalamus is the hub for energy homeostasis and food regulation. Disruption of this network can lead to hyperphagia and morbid obesity, best documented in craniopharyngioma literature, but also noted elsewhere. Regulation of blood pressure, thirst, and body temperature have also been seen and well-reported. Daytime sleepiness and disruption of circadian rhythm is a known warning sign for hypothalamic dysfunction, and the rates of this following surgical treatment spike – one study notes an increase by up to 65–80% [7, 31]. Many of these hypothalamic injury-induced sequelae persist even beyond hormone replacement treatment.

Finally, it is worth noting that the individual risks associated with any particular approach to this region must not be ignored and should be compared on a case-by-case basis to determine the best surgical approach to the lesion. Many of these approach-specific sequelae are detailed above.

Regarding the nonoperative considerations to surgery in many of these patients, there are numerous factors that can influence the decision to operate or not to operate. A proper assessment of comorbid diseases and operative risk is essential to a neurosurgical worker. Although surgery may be the best option to manage the specific lesion, the patient must be considered as a whole, and the decision whether to undergo surgery should be done under such pretense. Additionally, workup of these lesions should include a discussion with the patient on or ability and mortality of their disease both before and after a possible surgery. Neurosurgery can be incredibly taxing to the body, and proper understanding of this will help the patient make the most appropriate decision regarding their care. If at any point it becomes clear that the patient will not be able to tolerate the operation or postoperative therapy, open dis-

cussion with the patient is warranted, including a discussion on long-term goals of care.

Conclusion

As neurosurgical technology grows and expands, the ability to surgically manage patients with hypothalamic diseases and lesions becomes more and more possible and prevalent. Common diseases that affect the hypothalamus include tumors (gliomas, hamartomas, astrocytomas, expansive craniopharyngiomas) and vascular abnormalities (SCMs, AVMs) and can be approached by either endonasal techniques or cranial approaches (pterional, supraorbital, subfrontal, etc.).

The decision regarding the appropriate approach to a lesion depends heavily on the location of the tumor. Lateral hypothalamic lesions are best approached via pterional approach, whereas medial lesions can be approached via endonasal or translaminar approaches. Superior lesions and lesions affecting the third ventricle can be approached via the transcallosal approach or the pterional/modified pterional approaches. The ultimate decision made by the surgeon should consider major side effects or possible complications, as well.

The most common postoperative sequelae are due to the loss of hypothalamic function secondary to removal of tissue within the region. Other complications – namely, damage to surrounding structures – can be minimized and mitigated by good surgical technique and good surgical approach. The hypothalamus' location makes clear operations and wide surgical windows difficult to obtain.

New prospects to possibly improve these operations include increased image guidance capabilities. Improvement to intraoperative image guidance will help maximize excision of unwanted tissue while minimizing the loss of extraneous tissue or damage to other structures. Similarly, advanced and innovative operative approaches can provide better surgical windows than we know now.

References

1. Acioly M, Cohen-Gadol A. Pterional craniotomy. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v2.ch02>.
2. Acioly M, Cohen-Gadol A. Supraorbital craniotomy. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v2.ch04>.
3. Aihara Y, Chiba K, Eguchi S, Amano K, Kawamata T. Pediatric optic pathway/hypothalamic Glioma. *Neurol Med Chir (Tokyo)*. 2018;58:1–9. <https://doi.org/10.2176/nmc.ra.2017-0081>.
4. Brain Arteriovenous Malformations (AVM) [WWW Document]. n.d. http://brainavm.uhnres.utoronto.ca/malformations/brain_avm_index.htm. Accessed 24 Dec 2019.
5. Carmel PW. Surgical syndromes of the hypothalamus. *Clin Neurosurg*. 1980;27:133–59. https://doi.org/10.1093/neurosurgery/27.cn_suppl_1.133.
6. Cohen-Gadol A. Subfrontal Translamina Terminalis approach. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2016. <https://doi.org/10.18791/nsatlas.v4.ch05.5.3.2>.
7. Day JD. Surgical approaches to suprasellar and parasellar tumors. *Neurosurg Clin N Am*. 2003;14:109–22. [https://doi.org/10.1016/s1042-3680\(02\)00071-2](https://doi.org/10.1016/s1042-3680(02)00071-2).
8. Dodson S, Kamer A, Cohen-Gadol A. Histoplasmosis. In: Cohen-Gadol A, editor. Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v1.03.02.17>.
9. Frost E. Seven AVMs tenets and techniques for resection. *J Neurosurg Anesthe* [WWW Document]. 2015. https://journals.lww.com/jnsa/Citation/2015/07000/Seven_AVMs_Tenets_and_Techniques_for_Resection.12.aspx. Accessed 24 Dec 2019.
10. Goodden J, Pizer B, Pettorini B, Williams D, Blair J, Didi M, Thorp N, Mallucci C. The role of surgery in optic pathway/hypothalamic gliomas in children. *J Neurosurg Pediatr*. 2014;13:1–12. <https://doi.org/10.3171/2013.8.PEDS12546>.
11. Hendricks B, Cohen-Gadol A. Craniopharyngioma (Transcranial approach). In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2016. <https://doi.org/10.18791/nsatlas.v5.ch09.2>.
12. Hendricks B, Cohen-Gadol A. Ventricular tumors. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2016. <https://doi.org/10.18791/nsatlas.v4.ch05.2>.
13. Hendricks B, Cohen-Gadol A. Introduction to third ventricular tumors. In: Neurosurgical atlas. Neurosurgical Atlas Inc.; 2016. <https://doi.org/10.18791/nsatlas.v4.ch05.5.1>.
14. Maraire JN, Awad IA. Intracranial cavernous malformations: lesion behavior and management strategies. *Neurosurgery*. 1995;37:591–605. <https://doi.org/10.1227/00006123-199510000-00001>.
15. Mirza F, Cohen-Gadol A. Deep subcortical AVMs. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v3.ch02.8>.
16. Mullatti N, Selway R, Nashef L, Elwes R, Honavar M, Chandler C, Morris R, Jarosz J, Buchanan C, Polkey C. The clinical spectrum of epilepsy in children and adults with hypothalamic Hamartoma. *Epilepsia*. 2003;44:1310–9. <https://doi.org/10.1046/j.1528-1157.2003.04103.x>.
17. Müller HL. Craniopharyngioma and hypothalamic injury: latest insights into consequent eating disorders and obesity. *Curr Opin Endocrinol Diabetes Obes*. 2016;23:81–9. <https://doi.org/10.1097/MED.0000000000000214>.
18. Parish J, Cohen-Gadol A. Supratentorial cavernous malformation resection. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2016. <https://doi.org/10.18791/nsatlas.v3.ch04.2>.
19. Pop MG, Crivii C, Opincariu I. Anatomy and function of the hypothalamus. In: Hypothalamus in health and diseases; 2018. <https://doi.org/10.5772/intechopen.80728>.
20. Rigamonti D, Hadley MN, Drayer BP, Johnson PC, Hoenig-Rigamonti K, Knight JT, Spetzler RF. Cerebral cavernous malformations. Incidence and familial occurrence. *N Engl J Med*. 1988;319:343–7. <https://doi.org/10.1056/NEJM198808131906005>.
21. Rosenfeld JV, Harvey AS, Wrennall J, Zacharin M, Berkovic SF. Transcallosal resection of hypothalamic Hamartomas, with control of seizures, in children with Gelastic epilepsy. *Neurosurgery*. 2001;48:108–18. <https://doi.org/10.1097/00006123-200101000-00019>.
22. Sawamura Y, Kamada K, Kamoshima Y, Yamaguchi S, Tajima T, Tsubaki J, Fujimaki T. Role of surgery for optic pathway/hypothalamic astrocytomas in children. *Neuro-Oncology*. 2008;10:725–33. <https://doi.org/10.1215/15228517-2008-033>.
23. Seltman R, Kamer A, Cohen-Gadol A. Pilocytic Astrocytoma (PA). In: Cohen-Gadol A, editor. Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v1.03.01.31>.
24. Seltman R, Kamer A, Cohen-Gadol A. Craniopharyngioma. In: Cohen-Gadol A, editor. Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v1.03.01.09>.
25. Shaikh K, White I, Cohen-Gadol A. Hypothalamic Hamartoma. In: Neurosurgical atlas. Neurosurgical Atlas, Inc.; 2017. <https://doi.org/10.18791/nsatlas.v7.ch06>.
26. Shim K-W, Chang J-H, Park Y-G, Kim H-D, Choi J-U, Kim D-S. Treatment modality for intractable epilepsy in hypothalamic hamartomatous lesions. *Neurosurgery*. 2008;62:847–56; discussion 856. <https://doi.org/10.1227/01.neu.0000318170.82719.7c>.
27. Spetzler RF, Kondziolka DS, Higashida RT, Kalani MYS. Comprehensive management of arteriovenous malformations of the brain and spine. Cambridge University Press. Cambridge, England, United Kingdom; 2015.
28. Sutton LN, Molloy PT, Sernyak H, Goldwein J, Phillips PL, Rorke LB, Moshang T, Lange B, Packer

- RJ. Long-term outcome of hypothalamic/chiasmatic astrocytomas in children treated with conservative surgery. *J Neurosurg.* 1995;83:583–9. <https://doi.org/10.3171/jns.1995.83.4.0583>.
29. Tang B, Xie SH, Xiao LM, Huang GL, Wang ZG, Yang L, Yang XY, Xu S, Chen YY, Ji YQ, Zeng EM, Hong T. A novel endoscopic classification for craniopharyngioma based on its origin. *Sci Rep.* 2018;8:1–21. <https://doi.org/10.1038/s41598-018-28282-4>.
30. Venger BH, Landon G, Rose JE. Solitary Histoplasmoma of the thalamus: case report and literature review. *Neurosurgery.* 1987;20:784–7. <https://doi.org/10.1227/00006123-198705000-00019>.
31. Wait SD, Abla AA, Killory BD, Nakaji P, Rekat HL. Surgical approaches to hypothalamic hamartomas. *Neurosurg Focus.* 2011;30:E2. <https://doi.org/10.3171/2010.11.FOCUS10250>.
32. Wen HT, de Oliveira E, Tedeschi H, Andrade FJC, Rhoton AL. The pterional approach: surgical anatomy, operative technique, and rationale. 2001; <https://doi.org/10.1053/otns.2001.25567>.
33. Zhao C, Chen Z, Xu N, Xue T, Wu X, You W, Zhu Y, Wang Z. Comparative analysis on microsurgical removal of craniopharyngioma via lateral supraorbital approach and standard pterional approach. *Chin Neurosurg J.* 2018;4:16. <https://doi.org/10.1186/s41016-018-0126-7>.
34. Zoli M, Mazzatenta D, Valluzzi A, Marucci G, Acciarri N, Pasquini E, Frank G. Expanding indications for the extended endoscopic endonasal approach to hypothalamic gliomas: preliminary report. *Neurosurg Focus.* 2014;37:E11. <https://doi.org/10.3171/2014.7.FOCUS14317>.

Hormone Excess Syndromes of the Hypothalamic-Pituitary Axis

10

Nidhi Agrawal, Hyon Kim, Kyla Wright,
and Sonal Mehta

Prolactinomas

Hyperprolactinemia is a condition of excess prolactin, which could be physiologic, pathologic, or idiopathic. Prolactinomas are the most frequent cause of chronic hyperprolactinemia [1]. They are also the most common type of pituitary adenomas and account for approximately half of pituitary tumors requiring medical attention [2]. Based on the size of the tumors, prolactinomas can be classified as microadenomas if less than 10 mm in size, macroadenomas if greater than 10 mm in size, or giant adenomas if larger than 4 cm in size [3].

Clinical Manifestations

The clinical presentation of prolactinomas differs based on the size of the adenoma and the sex of

the patient. Most clinically diagnosed prolactinomas are found in premenopausal women and are microadenomas. In men, prolactinomas typically present as macroadenomas or giant adenomas [4]. Patients with macroprolactinomas or giant prolactinomas typically present with neurologic symptoms caused by mass effect. These include headaches, vision changes, cranial neuropathies, hypopituitarism, seizures, and cerebrospinal fluid rhinorrhea [1, 2, 5]. Prolactinomas predominantly affect women between the ages of 20 and 50 years [6]. In premenopausal women, classic symptoms include oligomenorrhea or amenorrhea, galactorrhea, infertility, decreased libido, and decreased bone mass [1, 2]. Postmenopausal women typically do not present with typical hyperprolactinemia-related symptoms as these are dependent on intact ovarian function. These patients therefore usually present with mass effect symptoms when their adenomas become large enough [7]. In men, hyperprolactinemia causes hypogonadotropic hypogonadism. This can manifest as decreased libido, impotence and erectile dysfunction, infertility, gynecomastia, and rarely galactorrhea [3, 8, 9]. Males sometimes present with small testes and prostate [9]. They can also present with decreased bone mass and over time can have diminished energy, reduced muscle mass, and increased risk of osteopenia [1]. In children, prolactinomas typically occur during the start of puberty. The most common presenting symptoms are headaches and

N. Agrawal (✉) · H. Kim
Division of Endocrinology, Diabetes and Metabolism,
NYU Langone Medical Center, New York, NY, USA
e-mail: Nidhi.Agrawal@nyulangone.org; Hyon.Kim@nyulangone.org

K. Wright
NYU Grossman School of Medicine,
New York, NY, USA
e-mail: Kyla.Wright@nyulangone.org

S. Mehta
Department of Medicine, NYU Langone Medical
Center, New York, NY, USA
e-mail: Sonal.Mehta@nyulangone.org

Table 10.1 Clinical characteristics of prolactinomas

Type of patient	Clinical manifestations
Premenopausal women	Menstrual irregularities such as amenorrhea and oligomenorrhea, galactorrhea, infertility, decreased libido, decreased bone mass, anxiety, depression, fatigue, and emotional instability
Postmenopausal women	Mass effect symptoms including headaches, vision changes, cranial neuropathies, seizures
Males	Hypogonadism, decreased libido, erectile dysfunction, infertility, gynecomastia, galactorrhea (rare), anxiety, depression, fatigue, and emotional instability
Children	Girls: menstrual irregularities, galactorrhea Boys: delayed pubertal development, hypogonadism
Patients with macroadenomas and giant prolactinomas	Mass effect symptoms including headaches, vision changes, cranial neuropathies, seizures

Modified with information from Majumdar et al. [1], Chanson et al. [2], Yatavelli et al. [3], Luciano, AA [5], Shimon et al. [7], Carter et al. [8], Segal et al. [9], Hoffman et al. [10], Melmed et al. [11]

visual impairment from mass effect [10]. In girls, menstrual irregularities and galactorrhea can be seen, while boys present with delayed pubertal development and hypogonadism [11]. Girls are more often affected than boys; however, prolactinomas in boys are typically larger and more aggressive [10]. Other clinical manifestations in males and females include anxiety, depression, fatigue, and emotional instability. Approximately 10% of prolactinomas also co-secrete growth hormone so gigantism and symptoms of acromegaly can also be seen in patients [3]. The clinical manifestations of prolactinomas are summarized in Table 10.1 [1–3, 5, 7–11].

Diagnosis

When clinical manifestations of prolactin excess are suspected, laboratory evaluation begins with measurement of prolactin levels. The normal prolactin range is approximately

5–20 mcg/L. Prolactin levels are typically elevated and generally correlate with the prolactinoma size [2]. Prolactin levels greater than 250 mcg/L typically indicate the presence of a prolactinoma over other causes, and levels greater than 500 mcg/L are usually diagnostic of a macroprolactinoma. However, select medications such as risperidone and metoclopramide can cause prolactin levels greater than 200 mcg/L [11].

In some cases, there may be an artificially low prolactin level when a prolactinoma is present, caused by the “hook effect.” Prolactin levels are measured using an assay based on a sandwich principle in which a prolactin molecule reacts with a capture antibody and then binds to a detection antibody. Each antibody is specific for a particular epitope on the prolactin molecule. The “hook effect” is caused by an assay artifact when there are extremely high levels of prolactin that saturate both the capture and detection antibodies used in the assay, preventing the antibody-prolactin-antibody sandwich formation, resulting in a falsely low reported value (Fig. 10.1) [12–14]. In order to determine whether the low prolactin is accurate or as a result of the hook effect, serial dilution of the prolactin serum sample should be obtained, which will result in an elevated prolactin if the patient has a prolactinoma [3, 14]. There are also cases in which there is an elevated measured prolactin level; however, the true prolactin level is actually low. This occurs when patients have a higher molecular weight prolactin called macroprolactin. Macroprolactin forms by condensation of prolactin molecules or complex formation between prolactin and plasma proteins. Therefore, in patients with asymptomatic hyperprolactinemia, macroprolactin levels should be obtained to assess for this. The lab can precipitate the macroprolactin by pretreating the serum with polyethylene glycol before obtaining the immunoassay for prolactin [3, 15].

In patients suspected of having a hypothalamic-pituitary cause of their hyperprolactinemia, pituitary MRI with gadolinium enhancement should be obtained as it provides the best visualization of the sellar area [1].




Fig. 10.1 Hook effect sandwich assay. (a) shows the normal sandwich assay with the capture antibody in blue, the detection antibody in red, and the prolactin molecule in green. (b) shows the hook effect where there is excessive prolactin antigen (in green) which then binds up sites on both the capturing antibody (in blue) and detection antibody (in red), thereby reducing detection levels. (Modified from Binart et al. [12], Romijn [13], and Vilar et al. [14])




Fig. 10.2 Causes of hyperprolactinemia. (Modified with information from Majumdar et al. [1], Syro et al. [4], Melmed et al. [11], Molitch ME [16])

Differential Diagnosis

There is a wide differential for hyperprolactinemia and many etiologies need to be consid-

ered (Fig. 10.2) [1, 4, 11, 16]. Physiologic causes of elevated prolactin levels include pregnancy, lactation, chest wall stimulation, sleep, exercise, coitus, and stress. Hyperprolactinemia in these

cases is usually mild to moderate [1, 11]. While prolactin is synthesized in and secreted by the anterior pituitary, it is also regulated by the hypothalamus via inhibition by dopamine. Therefore, any factors that cause loss of dopamine inhibition via the effects of medications or destruction of the hypothalamus or hypothalamic-hypophyseal tract will result in prolactin excess [3, 17]. Thus, there are a broad range of pathologic etiologies including pituitary causes, hypothalamic-pituitary stalk diseases, medication-induced, and systemic or other causes.

Pituitary causes include prolactinomas, acromegaly due to co-secretion with growth hormone, surgery, and trauma. Hypothalamic-pituitary stalk disease etiologies include pituitary stalk compression, lymphocytic hypophysitis, irradiation, trauma, sarcoidosis infiltration, and tumors such as craniopharyngiomas, meningiomas, dysgerminomas, dermoid cysts, and pineal gland tumors. Medication-induced causes include dopamine receptor blocking agents (such as phenothiazines, haloperidol, metoclopramide, and atypical antipsychotics), dopamine-depleting agents (such as reserpine, methyldopa, and opiates), histamine receptor antagonists (such as ranitidine), stimulators of the serotonergic pathway (such as amphetamine and hallucinogens), estrogens and antiandrogens, serotonin reuptake inhibitors (such as fluoxetine), and calcium channel blockers (such as verapamil). Systemic or other causes include chronic renal failure, hypothyroidism, cirrhosis, pseudocyesis, epileptic seizures, polycystic ovarian disease (PCOS), and adrenal insufficiency [1, 4, 11, 16]. Idiopathic hyperprolactinemia accounts for approximately 40% of cases [1].

Management

The preferred initial treatment for prolactinomas is dopamine agonists. Dopamine agonists bind to the dopamine D2 receptors and inhibit the synthesis and secretion of prolactin from the anterior pituitary gland [18]. These have been shown to lower prolactin levels, decrease tumor size, and restore gonadal function for symptomatic patients

[11]. Cabergoline and bromocriptine are the most commonly used dopamine agonists. Others include pergolide, quinagolide, terguride, and metergoline. All are ergot alkaloids except quinagolide, which is a nonergot dopamine agonist [1]. Cabergoline has a long half-life ranging from 63 to 69 hours and the initial dose is usually 0.25 mg twice weekly. The dose can be increased slowly by 0.25 mg twice weekly at a minimum of every 4 weeks [18]. Bromocriptine treatment is usually started at a dose of 1.25–2.5 mg per day and can be titrated up by 2.5 mg per day every 2–7 days. The maintenance dose ranges from 2.5 to 15 mg per day [18]. Side effects of dopamine agonists include nausea, vomiting, headaches, constipation, dizziness, orthostatic hypotension, depression, digital vasospasm, and nasal stuffiness [1]. Regarding efficacy of treatment, bromocriptine treatment has been shown to cause normoprolactinemia and restoration of normal menses in 80–90% of patients [19]. Cabergoline is used more often as it has been shown to achieve higher rates of normalizing prolactin levels and higher frequency of shrinking tumors [11]. Normal prolactin levels are achieved in 80–100% of patients with microprolactinomas and 75–95% of patients with macroprolactinomas. Cabergoline can also normalize prolactin levels in patients with resistance to other dopamine agonists [20]. Treatment with dopamine agonists should be tapered and stopped if prolactin levels normalize, if the tumor is not visible on MRI after at least 2 years of treatment, or if the tumor shrinks by 50% with a mass at least 5 mm from the optic chiasm and no evidence of cavernous sinus or other critical area invasion [3, 11, 21].

In pregnant patients, dopamine agonists should be avoided if possible as they can cross the placenta. However, treatment is indicated during pregnancy if the patient has a large adenoma or symptomatic growth during pregnancy [3, 20]. There do not appear to be any adverse pregnancy outcomes or risks for fetal malformations with either bromocriptine or cabergoline [22].

Resistant prolactinomas are those in which there is failure to achieve normal prolactin levels, reduce prolactin levels by greater than 50%, induce ovulation in women, or reduce symptoms

or normalize prolactin despite high cabergoline doses [20]. Symptomatic patients with prolactinomas resistant to medical therapy may benefit from transsphenoidal surgery [11, 23]. Other indications for surgical therapy include patients with psychiatric conditions with contraindications to dopamine agonist treatment, women with adenomas impinging on the optic chiasm seeking fertility, and patients presenting with CSF leak or pituitary apoplexy [20]. Radiation therapy is typically only used in patients with prolactinomas resistant to cabergoline therapy and who have undergone surgery and have residual tumor present. There is a risk of causing hypopituitarism with radiation therapy [1, 3].

Cushing's Syndrome

Cushing's syndrome results from prolonged glucocorticoid (GC) excess, from either exogenous GCs or endogenous sources of cortisol. Exogenous Cushing's syndrome is the most frequent cause of hypercortisolism. Most cases are iatrogenic, as high-dose or chronic use of GCs is common in the management of a number of inflammatory, autoimmune, and neoplastic diseases. All modes of exogenous GC delivery (oral, inhaled, topical, subcutaneous) have been implicated in the development of Cushing's syndrome and thus must be ruled out initially [24, 25]. Conversely, endogenous Cushing's syndrome is rare, with an estimated incidence of 0.7–2.4 million per year [26, 27]. More frequently affecting women (female/male ratio 3:1) [28], endogenous Cushing's syndrome results from excessive adrenal cortisol secretion that can be either adrenocorticotrophic hormone (ACTH)-dependent or ACTH-independent. A majority are ACTH-dependent, making up about 80–85% of cases [29]. The causes of ACTH-dependent Cushing's syndrome include ACTH-secreting pituitary adenomas, ectopic ACTH secretion, and ectopic corticotropin-releasing hormone (CRH) secretion (Fig. 10.3) [30]. ACTH-secreting pituitary adenomas account for most cases of endogenous hypercortisolism and are referred to as “pituitary-dependent Cushing's syndrome” or




Fig. 10.3 Causes of endogenous Cushing's syndrome. (With permission from Tritos et al. [30])

“Cushing's disease,” in which hypersecretion of ACTH by the adenoma results in pathologic adrenal cortisol secretion [31]. ACTH-independent Cushing's syndrome is even rarer and is characterized by hypersecretion of cortisol by adrenal pathologies such as adrenal adenomas and adrenal carcinomas [30].

Clinical Manifestations

The clinical presentation of Cushing's syndrome is variable and in part is dependent on the extent and duration of cortisol excess. The characteristic features of Cushing's syndrome are depicted in Fig. 10.4 [26, 32–35]. In adults, the signs that best discriminate Cushing's syndrome include easy bruising, facial plethora, proximal myopathy, and purple striae [34]. Other features are less discriminatory but common, such as weight gain, fatigue, depression, facial fullness, acne, decreased libido, and menstrual changes. Central obesity or weight gain is the most common finding in Cushing's syndrome and is seen in approximately 95% of adults [26].

Furthermore, the initial presentation may be characterized by the presence of comorbidities [36–39]. Cushing's syndrome is often complicated by metabolic syndrome (consisting of systemic arterial hypertension, visceral obesity, impaired glucose metabolism, and dyslipidemia),