

NCBI Bookshelf. A service of the National Library of Medicine, National Institutes of Health.

StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-.

# Neuroanatomy, Amygdala

## Authors

Qais AbuHasan<sup>1</sup>; Vamsi Reddy<sup>2</sup>; Waquar Siddiqui<sup>3</sup>.

## Affiliations

<sup>1</sup> An-Najah National University

<sup>2</sup> McKinsey & Company

<sup>3</sup> UMKC

Last Update: July 17, 2023.

## Introduction

The limbic system is vital for one's normal functioning. This system acts as the center of emotions, behavior, and memory. It is also a contributor to the control of reactions to stress, attention, and sexual instincts. It comprises a set of complex structures anatomically divided into the limbic cortex, cingulate gyrus, parahippocampal gyrus, hippocampal formation, dentate gyrus, hippocampus, subauricular complex, septal area, hypothalamus, and amygdala.[1]

The amygdala gets its name from its resemblance to almonds; it is an almond-shaped structure formed by many nuclei sorted into five major groups; basolateral nuclei, cortical-like nuclei, central nuclei, other amygdaloid nuclei, and extended Amygdala.[1]

The amygdala has been implicated in many diseases, such as depression,[2] sleep debt and anger,[3] as well as other neuropsychiatric diseases.

## Structure and Function

The amygdala is an almond-shaped structure that lies in the temporal lobe, lying just beneath the uncus. The amygdala is diverse and complex in structure and comprises approximately 13 nuclei. They further subdivide into extensive internuclear and intranuclear connections. These nuclei functionally sort into five major groups: basolateral nuclei, cortical-like nuclei, central nuclei, other amygdaloid nuclei, and extended amygdala. Amygdala is one of the components of the limbic system, which is responsible for the control of emotions and behavior besides memory formation. Anatomically, the amygdala lies at the anterior border of the hippocampal formation and the anterior aspect of the lateral ventricle's inferior horn where it merges with the peri-amygdaloid cortex, which forms part of the surface of the uncus.[1]

Amygdala manages the processing of information between prefrontal-temporal association cortices and the hypothalamus. Amygdala has neural circuits to carry out its different functions with two major output pathways; the Dorsal route via stria terminalis that projects to the septal area and hypothalamus, and the ventral route via the ventral amygdalofugal pathway which terminates in the septal area, hypothalamus, and the medial dorsal thalamic nucleus.[1] The amygdala also has connections with the basal ganglia circuit via its projections to the ventral pallidum and ventral striatum; these projections are relayed back to the cortex via the dorsomedial nucleus of the thalamus.[1] The basolateral circuit includes the amygdala (especially the basolateral amygdala), the orbitofrontal and anterior temporal cortex, and in the thalamus, the magnocellular division of the dorsomedial nucleus (frontothalamic pathway), which serves as a relay back to the orbitofrontal cortex.[4] The circuit has been proposed as a substrate for the human ability to infer the intentions of others from their language, gaze, and gestures (Theory of mind and social cognition),[5] and helps with social interactions.

The amygdala also functions in regulating anxiety, aggression, fear conditioning, emotional memory, and social cognition.[1] Electrical stimulation of the amygdala evokes fear and anxiety responses in humans while lesions block certain types of unconditioned fear. For example, rats with lesions in the amygdala show reduced freezing in response to cats, or cat hair, attenuated analgesia, heart rate responses to loud noise, and have reduced taste neophobia. However, amygdala lesions do not affect other measures of fear such as an open arm avoidance in an elevated plus-maze in rats or analgesia to shock. The amygdala is also necessary for learning by fear, amygdala lesions disrupt the acquisition of both active avoidance (escape from fear), and passive avoidance of conditioned responses, but do not affect retention. The amygdala processes not only emotions of fear and aversive stimuli, but it is also involved in conditioning using stimuli of appetite such as food, sex, and drugs. As for its role in memory, the activation of the amygdala has a modulatory effect on the acquisition and consolidation of memories that evoke an emotional response.[6]

Some parts of the amygdala have even more specific functions. The basolateral nucleus (BLA) is a cortical-like structure in the dorsal amygdala, and it regulates behavioral and physiological responses to stress.[7] The central amygdala (CeA) plays a crucial role in physiological responses to stressors, such as fearful stimuli, stressful stimuli, and some drug-related stimuli.[8] Meanwhile, the extended amygdala, named the bed nucleus of the stria terminalis (BNST), is involved in anxiety and stress.[9]

## Embryology

At approximately the third week of gestation, the notochord induces neurulation, a process by which ectoderm above the notochord becomes the neural ectoderm which will later form the neural tube and crest. *Noggin*, *chordin*, *BMP4*, and *FGF8* are some of the genes involved. The neural tube closes by week six. The rostral end will be the lamina terminalis. In addition to the spinal cord, the neural tube differentiates into three primary vesicles for the forebrain, midbrain, and hindbrain. The forebrain further differentiates into the telencephalon and diencephalon; the midbrain continues to be the mesencephalon, and the hindbrain becomes the metencephalon and myelencephalon. These structures continue to differentiate into adult brain structures. The origin of the amygdaloid body or complex called the amygdala is traced to populations of diencephalic and telencephalic cells that form the floor of the lateral ventricle about three weeks after conception.[10] The telencephalon gives rise to the amygdala while neurons from the diencephalon migrate to further develop it.[11]

## Blood Supply and Lymphatics

The anterior choroidal artery is the preterminal branch of the internal carotid and provides blood supply to the amygdala, and it is drained by the posterior choroidal vein which ends eventually to form the great cerebral vein that drains into the straight sinus.[12]

Astroglia podocytes form the blood-brain barrier by wrapping podocytes around capillaries. These cells protect the brain from toxins in the blood and facilitate nutrient transport to the neurons. Astroglia also forms a system of microscopic perivascular channels permeating the brain that transmit a cerebrospinal fluid (CSF)-like lymphatic vessels. The system enables CSF to clear the metabolic waste and distribute glucose, amino acids, lipids, and neurotransmitters. This system is most active during sleep, contributing to its restoration function. Arterial pulsation drives lymphatic flow, suggesting that exercise may also enhance it: aging, brain trauma, and ischemia decrease that CSF flow. Also, larger lymphatic vessels in the meninges help absorb interstitial fluid into the dural venous sinuses.[13][14]

## Surgical Considerations

For the management of drug-resistant mesial temporal lobe epilepsy, amygdalohippocampectomy, either selective or combined with anterior temporal lobectomy is being considered with much importance.

## Clinical Significance

The amygdala has been associated with many diseases, mainly neuropsychiatric. Many studies showed its effects on depression.[2][15] Others stressed the involvement of the amygdala in post-traumatic stress disorder as there is a bilateral reduction of the hippocampus and amygdala in PTSD.[16] Neural functioning among patients with PTSD is characterized by attenuated prefrontal inhibition on the limbic system, resulting in emotional dysregulation, and suggests that amygdala neurofeedback may not only be therapeutic for this patient group but may also be used as a future adjunctive treatment.[17] The amygdala and the limbic system might also be involved in chronic pain and has an association with the emotional effects of such pain.[18]

## Review Questions

- [Access free multiple choice questions on this topic.](#)
- [Comment on this article.](#)

## References

1. Rajmohan V, Mohandas E. The limbic system. Indian J Psychiatry. 2007 Apr;49(2):132-9. [PMC free article: PMC2917081] [PubMed: 20711399]
- 2.

- Ruiz NAL, Del Ángel DS, Olguín HJ, Silva ML. Neuroprogression: the hidden mechanism of depression. *Neuropsychiatr Dis Treat*. 2018;14:2837-2845. [PMC free article: PMC6214587] [PubMed: 30464468]
3. Saghir Z, Syeda JN, Muhammad AS, Balla Abdalla TH. The Amygdala, Sleep Debt, Sleep Deprivation, and the Emotion of Anger: A Possible Connection? *Cureus*. 2018 Jul 02;10(7):e2912. [PMC free article: PMC6122651] [PubMed: 30186717]
  4. Deakin JF, Slater P, Simpson MD, Gilchrist AC, Skan WJ, Royston MC, Reynolds GP, Cross AJ. Frontal cortical and left temporal glutamatergic dysfunction in schizophrenia. *J Neurochem*. 1989 Jun;52(6):1781-6. [PubMed: 2566649]
  5. Frith C. Brain mechanisms for 'having a theory of mind'. *J Psychopharmacol*. 1996 Jan;10(1):9-15. [PubMed: 22302722]
  6. Sah P, Faber ES, Lopez De Armentia M, Power J. The amygdaloid complex: anatomy and physiology. *Physiol Rev*. 2003 Jul;83(3):803-34. [PubMed: 12843409]
  7. Bhatnagar S, Vining C, Denski K. Regulation of chronic stress-induced changes in hypothalamic-pituitary-adrenal activity by the basolateral amygdala. *Ann N Y Acad Sci*. 2004 Dec;1032:315-9. [PubMed: 15677440]
  8. Gilpin NW, Herman MA, Roberto M. The central amygdala as an integrative hub for anxiety and alcohol use disorders. *Biol Psychiatry*. 2015 May 15;77(10):859-69. [PMC free article: PMC4398579] [PubMed: 25433901]
  9. Li C, Pleil KE, Stamatakis AM, Busan S, Vong L, Lowell BB, Stuber GD, Kash TL. Presynaptic inhibition of gamma-aminobutyric acid release in the bed nucleus of the stria terminalis by kappa opioid receptor signaling. *Biol Psychiatry*. 2012 Apr 15;71(8):725-32. [PMC free article: PMC3314138] [PubMed: 22225848]
  10. Müller F, O'Rahilly R. The amygdaloid complex and the medial and lateral ventricular eminences in staged human embryos. *J Anat*. 2006 May;208(5):547-64. [PMC free article: PMC2100220] [PubMed: 16637878]
  11. García-Moreno F, Pedraza M, Di Giovannantonio LG, Di Salvio M, López-Mascaraque L, Simeone A, De Carlos JA. A neuronal migratory pathway crossing from diencephalon to telencephalon populates amygdala nuclei. *Nat Neurosci*. 2010 Jun;13(6):680-9. [PubMed: 20495559]
  12. Kiernan JA. Anatomy of the temporal lobe. *Epilepsy Res Treat*. 2012;2012:176157. [PMC free article: PMC3420617] [PubMed: 22934160]
  13. Harrison IF, Siow B, Akilo AB, Evans PG, Ismail O, Ohene Y, Nahavandi P, Thomas DL, Lythgoe MF, Wells JA. Non-invasive imaging of CSF-mediated brain clearance pathways via assessment of perivascular fluid movement with diffusion tensor MRI. *Elife*. 2018 Jul 31;7 [PMC free article: PMC6117153] [PubMed: 30063207]
  14. Dave RS, Jain P, Byraredy SN. Functional Meningeal Lymphatics and Cerebrospinal Fluid Outflow. *J Neuroimmune Pharmacol*. 2018 Jun;13(2):123-125. [PMC free article: PMC5930060] [PubMed: 29464588]
  15. Helm K, Viol K, Weiger TM, Tass PA, Grefkes C, Del Monte D, Schiepek G. Neuronal connectivity in major depressive disorder: a systematic review. *Neuropsychiatr Dis Treat*. 2018;14:2715-2737. [PMC free article: PMC6200438] [PubMed: 30425491]
  16. Ahmed-Leitao F, Spies G, van den Heuvel L, Seedat S. Hippocampal and amygdala volumes in adults with posttraumatic stress disorder secondary to childhood abuse or maltreatment: A systematic review. *Psychiatry Res Neuroimaging*. 2016 Oct 30;256:33-43. [PubMed: 27669407]
  17. Nicholson AA, Rabellino D, Densmore M, Frewen PA, Paret C, Kluetsch R, Schmahl C, Théberge J, Neufeld RW, McKinnon MC, Reiss J, Jetly R, Lanius RA. The neurobiology of emotion regulation in posttraumatic stress disorder: Amygdala downregulation via real-time fMRI

neurofeedback. Hum Brain Mapp. 2017 Jan;38(1):541-560. [PMC free article: [PMC6866912](#)] [PubMed: 27647695]

18. Thompson JM, Neugebauer V. Cortico-limbic pain mechanisms. Neurosci Lett. 2019 May 29;702:15-23. [PMC free article: [PMC6520155](#)] [PubMed: 30503916]

**Disclosure:** Qais AbuHasan declares no relevant financial relationships with ineligible companies.



**Disclosure:** Vamsi Reddy declares no relevant financial relationships with ineligible companies.

**Disclosure:** Waquar Siddiqui declares no relevant financial relationships with ineligible companies.

## Figures

### Amygdala

The Amygdala is in the Limbic system of the brain. It is the epicenter for emotions, most notably, fear.



Amygdala diagram

Contributed by Katherine Humphreys

Copyright © 2023, StatPearls Publishing LLC.

This book is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits others to distribute the work, provided that the article is not altered or used commercially. You are not required to obtain permission to distribute this article, provided that you credit the author and journal.

