

- improves the amyloid pathology in a mouse model of Alzheimer's disease. *J Neuropathol Exp Neurol*. 2010;20:313–22.
66. Bowen RL, Perry G, Xiong C, Smith MA, Atwood CS. A clinical study of lupon depot in the treatment of women with Alzheimer's disease: preservation of cognitive function in patients taking an acetylcholinesterase inhibitor and treated with high dose lupon over 48 weeks. *J Alzheim Dis*. 2015;44:549–60.
67. Williams KW, Elmquist JK. From neuroanatomy to behavior: central integration of peripheral signals regulating feeding behavior. *Nat Neurosci*. 2012;15:1350–5.
68. White H, Piper C, Schmader K. The association of weight change in Alzheimer's disease with severity of disease and mortality: a longitudinal analysis. *J Am Geriatr Soc*. 1998;46:1223–7.
69. Friedman JM, Mantzoros CS. 20 years of leptin: from the discovery of the leptin gene to leptin in our therapeutic armamentarium. *Metab Clin Exp*. 2015;64:1–4.
70. Huttenbrink K-B, Hummel T, Berg D, Gasser T, Hahner A. Olfactory dysfunction: common in later life and early warning of neurodegenerative disease. *Dtsch Arztebl Int*. 2013;110(1–2):1–7.
71. Biessels GJ, Staekenborg S, Brunner E, Brayne C, Scheltens P. Risk of dementia in diabetes mellitus: a systematic review. *Lancet Neurol*. 2006;5:64–74.
72. Liu J, Rutten-Jacobs L, Liu M, Markus HS, Traylor M. Causal impact of type 2 diabetes mellitus on cerebral small vessel disease: a Mendelian randomization analysis. *Stroke*. 2018;49(6):1325–31.
73. Attems J, Jellinger KA. The overlap between vascular disease and Alzheimer's disease—lessons from pathology. *BMC Med*. 2014;206:1–12.
74. La Monte de SM. Type 3 diabetes is sporadic Alzheimer disease: mini-review. *Eur Neuropsychopharmacol*. 2014;24:1954–60.
75. Claxton A, Baker LD, Hanson A, Trittschuh EH, Cholerton B, Morgan A, et al. Long-acting intranasal insulin detemir improves cognition for adults with mild cognitive impairment or early-stage Alzheimer's disease dementia. *J Alzheim Dis*. 2015;44:897–906.
76. Satoh A, Brace CD, Ben-Josef G, West T, Wozniak DF, Holtzman DM, Herzog ED, Imai S-I. SIRT1 promotes the central adaptive response to diet restriction through activation of the dorsomedial and lateral nuclei of the hypothalamus. *J Neurosci*. 2010;30:10220–32.
77. Shannon OM, Blossom CMS, Granic A, Lentjes M, Hayat S, Mulligan A, et al. Mediterranean diet adherence and cognitive function in older UK adults: the European Prospective Investigation into Cancer and Nutrition-Norfolk (EPIC-Norfolk) Study. *Am J Clin Nutr*. 2019;110:938–48.
78. Morris MC, Tangney CC, Wang Y, Sacks FM, Barnes LL, Bennett DA, Aggarwal NT. MIND diet slows cognitive decline with aging. *Alzheimers Dement*. 2015;11(9):1015–22.
79. Musiek ES, Xiong DD, Holtzman DM. Sleep, circadian rhythms, and the pathogenesis of Alzheimer's disease. *Exp Mol Med*. 2015;47(148):1–6.
80. Saper CB. The central circadian timing system. *Curr Opin Neurobiol*. 2013;23:747–51.
81. Tranah GJ, Blackwell T, Stone KL, Ancoli-Israel S, Paudel ML, Ensrud KE, et al. Circadian activity rhythms and risk of incident dementia and mild cognitive impairment in older women. *Ann Neurol*. 2011;70:722–32.
82. Sakurai T. Orexin deficiency and narcolepsy. *Curr Opin Neurobiol*. 2013;23:760–6.
83. Fronczek R, van Geest S, Frolich M, Overeem S, Roelandse FWC, Lammers GJ, Swaab DR. Hypocretin (orexin) loss in Alzheimer's disease. *Neurobiol Aging*. 2012;33:1642–50.
84. Gallone S, Boschi S, Rubino E, De Martino P, Scarpini E, Galimberti D, et al. Is HCRTR2 a genetic risk factor for Alzheimer's disease? *Dement Geriatr Cogn Disord*. 2014;38:245–53.
85. Ballard C, Corbett A. Agitation and aggression in people with Alzheimer's disease. *Curr Opin Psychiatry*. 2013;26:252–9.
86. Haller J. The neurobiology of abnormal manifestations of aggression—a review of hypothalamic mechanisms in cats, rodents and humans. *Brain Res Bull*. 2013;93:97–109.
87. Song J, Sieurin J, Wirdefeldt K, Pedersen NL, Almqvist C, Larsson H, Valdimarsdottir UA, Fang F. Association of stress-related disorders with subsequent neurodegenerative diseases. *JAMA Neurol*. 2020;77:411–8.
88. Canet G, Hernandez C, Zussy C, Chevalier N, Desrumaux C, Givalois L. Is AD a stress-related disorder? Focus on the HPA axis and its promising therapeutic targets. *Front Aging Neurosci*. 2019;11:1–9.
89. Carrol JC, Iba M, Bangasser DA, Valentino RJ, James MJ, Brunden KR, Lee VM-Y, Trojanowski JQ. Chronic stress exacerbates tau pathology, neurodegeneration, and cognitive performance through a corticotropin-releasing factor receptor-dependent mechanism in a transgenic mouse model of tauopathy. *J Neurosci*. 2011;31:14436–49.
90. Swaab DF, Bao A-M, Lucassen PJ. The stress system in the human brain in depression and neurodegeneration. *Age Res Rev*. 2005;4:141–94.
91. Thal DR, Del Tredici K, Ludolph AC, Hoozemans JJM, Rozemuller AJ, Braak H, Knippschild U. Stages of granulovacuolar degeneration: their relation to Alzheimer's disease and chronic stress response. *Acta Neuropathol*. 2011;122:577–89.
92. Mulak A, Bonaz B. Brain-gut-microbiota axis in Parkinson's disease. *World J Gastroenterol*. 2015;21(37):10609–20.
93. Mulak A. An overview of the neuroendocrine system in Parkinson's disease: what is the impact on diagnosis and treatment? *Expert Rev Neurotherapeut*. 2020;20:127–35.
94. Javoy-Agid F, Ruberg M, Pique L, Bertagna X, Taquet H, Studler JM, Cesselin F, Epelbaum J, Agid

- Y. Biochemistry of the hypothalamus in Parkinson's disease. *Neurology*. 1984;34(5):672–5.
95. Shannak K, Rajput A, Rozdilsky B, Kish S, Gilbert J, Hornykiewicz O. Noradrenaline, dopamine and serotonin levels and metabolism in the human hypothalamus: observations in Parkinson's disease and normal subjects. *Brain Res*. 1994;639(1):33–41.
96. Moore RY, Whone AL, Brooks DJ. Extrastriatal monoamine neuron function in Parkinson's disease: an 18F-dopa PET study. *Neurobiol Dis*. 2008;29(3):381–90.
97. Gurevich EV, Joyce JN. Distribution of dopamine D3 receptor expressing neurons in the human forebrain: comparison with D2 receptor expressing neurons. *Neuropsychopharmacology*. 1999;20(1):60–80.
98. Politis M, Piccini P, Pavese N, Koh S-B, Brooks DJ. Evidence of dopamine dysfunction in the hypothalamus of patients with Parkinson's disease: an *in vivo* ¹¹C-raclopride PET study. *Exp Neurol*. 2008;214:112–6.
99. Farde L, Hall H, Ehrin E, Sedvall G. Quantitative analysis of D2 dopamine receptor binding in the living human brain by PET. *Science*. 1986;231(4735):258–61.
100. Tandberg E, Larsen JP, Karlsen K. A community-based study of sleep disorders in patients with Parkinson's disease. *Mov Disord*. 1998;13:895–9.
101. Fronczek R, Overeem S, Lee SY, Hegeman IM, van Pelt J, van Duinen SG, Lammers GJ, Swaab DF. Hypocretin (orexin) loss in Parkinson's disease. *Brain*. 2007;130:1577–85.
102. Thannickal TC, Lai YY, Siegel JM. Hypocretin (orexin) cell loss in Parkinson's disease. *Brain*. 2007;130:1586–95.
103. Bensaid M, Tande D, Fabre V, Michel PP, Hirsch EC, Francois C. Sparing of orexin-A and orexin-B neurons in the hypothalamus and of orexin fibers in the substantia nigra of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated macaques. *Eur J Neurosci*. 2015;41:129–36.
104. Liu M-F, Xue Y, Liu Y-H, Diao H-L, Wang Y, Pa Y-P, Chen L. Orexin-A exerts neuroprotective effects via OX1R in Parkinson's disease. *Front Neurosci*. 2018;12:1–18.
105. Oliveira LM, Henrique E, Bustell I, Netto NFC, Moreira TS, Takakura AC, Caetano AL. Depletion of hypothalamic hypocretin/orexin neurons correlates with impaired memory in a Parkinson's disease animal model. *Exp Neurol*. 2020;323:1–12.
106. Van Raamsdonk IM, Murphy Z, Selva DM, Hamidzadeh R, Pearson J, Petersen A, et al. Testicular degeneration in Huntington disease. *Neurobiol Dis*. 2007;26:512–20.
107. Kandasamy M, Couillard-Despres S, Raber KA, Stephan M, Lehner B, Winner B, et al. Stem cell quiescence in the hippocampal neurogenic niche is associated with elevated transforming growth factor-beta signaling in an animal model of Huntington disease. *J Neuropathol Exp Neurol*. 2010;69:717–28.
108. Selvaraj K, Manickam N, Kumaran E, Thangadurai K, Elumalai G, Sekar A, Radhakrishnan RK, Kandasamy M. Deterioration of neuroregenerative plasticity in association with testicular atrophy and dysregulation of the hypothalamic-pituitary-gonadal (HPG) axis in Huntington's disease: a putative role of the huntingtin gene in steroidogenesis. *J Steroid Biochem Mol Biol*. 2020;197:1–7.
109. Mochel F, Haller RG. Energy deficit in Huntington disease: why it matters. *J Clin Invest*. 2011;121:493–9.
110. Gorges M, Vercruysse P, Muller H-P, Huppertz H-J, Rosenbohm A, Nagel G, et al. Hypothalamic atrophy is related to body mass index and age at onset of amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry*. 2017;88:033–41.
111. Gallo V, Wark PA, Jenab M, Pearce N, Brayne C, Vermeulen R, et al. Prediagnostic body fat and risk of death from amyotrophic lateral sclerosis: the EPIC cohort. *Neurology*. 2013;80:829–38.
112. O'Reilly EJ, Wang H, Weisskopf MG, Fitzgerald KC, Falcone G, McCullough ML, et al. Premorbid body mass index and risk of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Frontotemporal Degener*. 2013;14:205–11.
113. Dupuis L, Pradat PF, Ludolph AC, Loeffler JP. Energy metabolism in amyotrophic lateral sclerosis. *Lancet Neuro*. 2011;10:75–82.
114. Debray S, Race V, Crabbe V, Herdewyn S, Matthijs G, Goris A, et al. Frequency of C9orf72 repeat expansions in amyotrophic lateral sclerosis: a Belgian cohort study. *Neurobiol Aging*. 2013;34:2897–902.
115. Leko MB, Zupunski V, Kirinich J, Smilovic D, Hortobagyi T, Hof PR, Simic G. Molecular mechanisms of neurodegeneration related to C9orf72 hexanucleotide repeat expansion. *Behav Neurol*. 2019;2019:1–18.
116. Renton AE, Chio A, Traynor BJ. State of play in amyotrophic lateral sclerosis genetics. *Nat Neurosci*. 2014;17:17–23.
117. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, et al. Ubiquitinated TDP-43 in frontotemporal degeneration and amyotrophic lateral sclerosis. *Science*. 2006;314:130–3.
118. Lillo P, Matamala JM, Valenzuela D, Verdugo R, Castill JL, Ibanez A, Slachevsky A. Overlapping features of frontotemporal dementia and amyotrophic lateral sclerosis. *Rev Med Chil*. 2014;142(7):867–79.
119. Cykowski MD, Takei H, Schulz PE, Appel SH, Powell SZ. TDP-43 pathology in the basal forebrain and hypothalamus of patients with amyotrophic lateral sclerosis. *Acta Neuropathol Comm*. 2014;2:171–82.
120. Vercruysse P, Sinniger J, El Oussini H, Scekic-Zahirovic J, Dieterle S, Dengler R, et al. Alterations in the hypothalamic melanocortin pathway in amyotrophic lateral sclerosis. *Brain*. 2016;139:1106–22.
121. Siddique T. Molecular genetics of familial amyotrophic lateral sclerosis. *Adv Neuro*. 1991;56:227–31.

122. Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, et al. Mutations in Cu/Zn superoxide dismutase are associated with accumulation of neurofilaments in amyotrophic lateral sclerosis. *Nature*. 1993;362:59–62.
123. Gonzalez de Aguilar J-L, Gordon JW, Rene F, Lutz-bucher B, Kienlen-Cmpard P, Loeffler J-P. A mouse model of familial amyotrophic lateral sclerosis expressing a mutant superoxide dismutase 1 shows evidence of disordered transport in the vasopressin hypothalamo-neurohypophysial axis. *Eur J Neurosci*. 1999;11:4179–87.
124. Dedeene L, Van Schoor E, Vandenberghe R, Van Damme P, Poesen K, Thal DR. Circadian sleep/wake-associated cells show dipeptide repeat protein aggregates in C9orf72-related ALS and FTLN cases. *Acta Neuropathol Comm*. 2019;7:189–200.
125. Pellecchia MT, Pivonello R, Monsurro MR, Trojsi F, Longo K, Piccirillo G, et al. The GH-IGF system in amyotrophic lateral sclerosis: correlations between pituitary GH secretion capacity, insulin-like growth factors and clinical features. *Eur J Neurol*. 2010;17:666–71.
126. Sacca F, Quarantell M, Tucci RC, Piro R, Perrotta G, et al. A randomized controlled clinical trial of growth hormone in amyotrophic lateral sclerosis: clinical, neuroimaging, and hormonal results. *J Neuro*. 2012;259(1):132–8.
127. Sperling R. Functional MRI studies of associative encoding in normal aging, mild cognitive impairment and Alzheimer's disease. *Ann N Y Acad Sci*. 2007;1097:146–55.
128. Werner CT, Williams CJ, Fermelia MR, Lin D-T, Li Y. Circuit mechanisms of neurodegenerative diseases: a new frontier with miniature fluorescence microscopy. *Front Neurosci*. 2019;13:1–9.
129. Inda MC, Joshi S, Wang T, Bolaender A, Gandu S, Koren J, et al. The epichaperome is a mediator of toxic hippocampal stress and leads to protein connectivity-based dysfunction. *Nat Comm*. 2020;11:319–30.
130. Do K, Laing BT, Landry T, Bunner W, Mersaud N, Matsubara T, et al. The effects of exercise on hypothalamic neurodegeneration of Alzheimer's disease mouse model. *PLoS One*. 2018;13(1):e0190205.

Iatrogenic Hypothalamic Disorders

22

Zachary P. Wetsel, Ronak Patel, Roberto Rey Dios,
Vishnu Garla, and Christa O'Hana S. Nobleza

Non-surgical Iatrogenic Hypothalamic Disorders

The hypothalamus is a small structure that is packed with neurons that interface with various structures in the brain and control diverse functions in the human body. Various medications can affect hypothalamic function by modulating the activity of neurotransmitters or by causing structural damage. Multiple medications can cause similar effects on the hypothalamic-pituitary-adrenal axis. The iatrogenic effects of these medications can be classified according to the specific hypothalamic dysfunction they may cause. Knowledge of the various pharmaceutical hypothalamic affectations should guide decision-making on its administration and discussion with patients and family regarding adverse effects.

Z. P. Wetsel · R. R. Dios
Department of Neurosurgery, University of
Mississippi Medical Center, Jackson, MS, USA
e-mail: zwetsel@umc.edu; rrey-dios@umc.edu

R. Patel · C. O' H. S. Nobleza (✉)
Department of Neurology, Division of Neurosciences
Critical Care, University of Mississippi Medical
Center, Jackson, MS, USA
e-mail: rpatel4@umc.edu; csanluis@umc.edu

V. Garla
Department of Medicine, Division of Endocrinology,
University of Mississippi Medical Center,
Jackson, MS, USA
e-mail: vgarla@umc.edu

Hyperprolactinemia

Prolactin is a hormone secreted by lactotrophs located in the anterior pituitary gland, which facilitates lactation. Dopamine (D2) receptors in the tuberoinfundibular area are one of the most important aspects of prolactin regulation through the tuberoinfundibular dopaminergic system [1]. Hyperprolactinemia is classically defined as detecting prolactin level more than 20 ng/ml in men and greater than 25 ng/ml in women at least 2 h after waking up in a fasting state [2]. The basic mechanism for hyperprolactinemia involves either eliminating or reducing inhibitory pathways or through direct stimulation of lactotrophic prolactin production [3]. Hyperprolactinemia is characterized by amenorrhea and galactorrhea in women, and in men, low libido and fatigue predominate [4, 5].

Antipsychotics

Antipsychotics have been the foundation of schizophrenia therapy since the advent of chlorpromazine use 50 years ago [1]. These groups of medications are the most common ones to induce hyperprolactinemia, and their most common adverse effect is hyperprolactinemia [5]. The antipsychotic efficacies of the older antipsychotics (chlorpromazine, thioridazine, fluphenazine) have been shown to be directly proportional to their prolactin-raising potency [6–9]. These classical antipsychotics affect all

dopamine pathways involved in prolactin regulation because they are not selective [3]. Of the classical antipsychotics that include butyrophenones, phenothiazines, thioxanthenes, and pimozide that cause early and sustained prolactin increase during therapy, it is worth mentioning that flupenthixol deviates from this profile in that after the first increase above baseline in the first month of therapy, prolactin level normalization has been found more than 6 months after continued use [10, 11].

Atypical antipsychotics are known for improved antipsychotic effect and adverse event profile [3]. Possible explanations for this pharmacologic profile include receptor binding specificity, regional dopamine receptor preference, different D2 receptor affinity, combined dopamine agonistic and antagonistic activity, and combined antagonism of dopamine and serotonin receptors [3, 11–14]. Of the newer antipsychotics, risperidone has a significant prolactin raising potency [3] and can potentially be mistaken for a prolactinoma. Therefore, it is crucial to establish an accurate timeline of symptoms and risperidone use, as this may help avoid unnecessary imaging or treatment [15–17]. Clozapine, olanzapine, ziprasidone, and aripiprazole not only bind weakly but also have antagonist-agonist effect at the D2 receptor [18–22]. Consequently, with these medications, prolactin levels rise transiently and return to baseline within 8 h [23, 24].

Antidepressants

Significantly less common than antipsychotics, antidepressant medications by which their mechanism of action is through serotonin activity can cause asymptomatic hyperprolactinemia [5, 25]. Tricyclic antidepressants (TCA) have been known to cause modest hyperprolactinemia. Clomipramine raises prolactin levels by 60% in men and 87.5% in women [26]. The monoamine oxidase (MAO) inhibitors, pargyline and clorgyline, have been shown to increase prolactin levels by 5.8 and 8.6 ng/ml, respectively [27]. Selective serotonin reuptake inhibitors (SSRIs) have not been known to be associated with hyperprolactinemia [28].

Opiates

Opiates inhibit dopamine release from the hypothalamus and cause hyperprolactinemia indirectly by inhibiting prolactin-inhibiting factor release or stimulating prolactin-releasing factor production [29]. This effect of opioids is mediated via the μ receptor [30]. It has also been reported to control prolactin secretion through its action on the synchrony of pulsatile pattern of prolactin [31]. Morphine and morphine analogs cause a persistent rise in prolactin levels, whereas methadone use results only in transient elevations [32].

Antiemetics or Prokinetics

Metoclopramide and domperidone are dopamine antagonists used for the treatment of nausea, vomiting, or decreased gastric motility. Through their dopamine antagonistic mechanism, they prevent dopamine from binding to the D2 receptor and result in hyperprolactinemia [3]. Around half of the patients taking these medications develop hyperprolactinemia and present with amenorrhea, decreased libido, and galactorrhea [33, 34].

Antihypertensives

Alpha-methyldopa acts as a false neurotransmitter and prevents L-dopa conversion to dopamine leading to dopamine synthesis decrease [35]. Reserpine on the other hand affects dopamine hypothalamic storage [36]. Verapamil acts on the neuronal N-type calcium channel and inhibits the production of dopamine from the hypothalamus via an unknown mechanism [37]. This inhibition of dopamine results in hyperprolactinemia [37]. Romeo et al. studied 449 subjects on verapamil and observed that 8.5% of them developed hyperprolactinemia compared to 3% in the control group [38]. A common beta-blocker used for emergent blood pressure control is labetalol that may cause hyperprolactinemia via central pathways as it was shown that dopamine agonist pretreatment decreases prolactin response [39].

H2 Receptor Antagonists

Cimetidine and ranitidine stimulate prolactin secretion [3, 40, 41]. Notably, cimetidine has

been found to decrease hypothalamic dopamine release [42] and also acts on neurotransmitters such as pituitary GABAergic system [43].

Cholinomimetic Medications

It has been shown that peptide modulation of the hypothalamic-pituitary regulation of prolactin plays a role in the increase in prolactin after intravenous infusion of physostigmine, a cholinomimetic medication [44]. It was reported that cholinergic stimulation of beta-endorphins in the hypothalamus was associated with prolactin level increase [44].

Hypopituitarism

Hypopituitarism is the general decrease of pituitary hormone production which can be due to tumors, inflammation, infection, infarction, or autoimmune conditions or as a complication of medication or surgical procedures. Because the hypothalamus primarily exerts control over the pituitary gland, any medication affecting the hypothalamic function may in turn result in hypopituitarism.

Immune Checkpoint Inhibitors

Immuno-tolerance is achieved by co-stimulatory pathways involving the programmed death-1 receptor (PD-1) and its ligand (PD-L) and the cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) [45]. The PD-1/PD-L pathway inhibits the activation and proliferation of T cells and stops the production of cytokines [45] while cell cycle arrest and cellular death in regulatory and activated T cells [45–47]. The CTLA-4 pathway binds on CD28 molecules with B7-1 (CD80) and/or B7-2 (CD86) molecules on the antigen-presenting cell (APC) surface in lymph nodes, which is the early stage of T cell activation [45]. The PD-1 pathway on the other hand affects the last part of the immune response, by binding to PD-L1 and programmed death ligand 2 (PD-L2), in peripheral tissues and blocks T cells [48]. Its adverse effects, including endocrinopathies, are related to its immune-mediated effects [45]. These immune checkpoint inhibitors (ICIs) may cause dysfunction of self-reactive quiescent T cells [49].

Secondary adrenal insufficiency due to failure of the hypothalamic-pituitary axis has been reported [50, 51]. Pituitary hypophysitis can present due to symptoms associated with the hormonal deficiencies or actual mass effect from the gland [50]. Direct hypothalamic involvement can be due to mass effect or direct autoimmune dysfunction that may present as panhypopituitarism as well. A high index of suspicion is needed as the presentation may be subtle until severe crisis occurs. ICI-induced hypophysitis affects the corticotrophs, gonadotrophs, thyrotrophs, and rarely lactotrophs [52]. The incidence of hypophysitis with ICIs is 0.4–1.7%. The highest rate is noted with ipilimumab (5.6%), followed by tremelimumab (1.8%). Nivolumab and pembrolizumab have a hypophysitis incidence of only 0.5% [53]. The mechanism of ICI-induced hypophysitis is unclear; however, complement activation by the antibodies could play a role. This also may explain the higher incidence of hypophysitis seen with ipilimumab as it is an IgG1 antibody that can activate the complement pathway [54]. Hypophysitis presents 4–10 weeks after treatment. Clinical features include headache, weakness, and fatigue. Pituitary hormonal evaluation could show a low adrenocorticotrophic hormone (ACTH), cortisol, free t4, luteinizing hormone (LH), and follicle stimulating hormone (FSH). MRI of the pituitary gland may show an enlarged pituitary with homogenous or heterogeneous enlargement; however, a normal appearance of the pituitary gland does not rule out hypophysitis [55, 56]. Treatment with high-dose glucocorticoids is initiated when there is a mass effect or visual defects.

Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH)

Vasopressin or antidiuretic hormone (ADH) is synthesized in the hypothalamus and is stored in the posterior pituitary. ADH is secreted in response to an increase in osmolality detected by the osmoreceptors in the anterior hypothalamus [57, 58]. Its criteria were first established about 40 years ago [59] and include hyponatremia, low

plasma osmolality, high urine osmolality, and high urine sodium in the absence of volume depletion and normal renal and adrenal function. Its clinical manifestation depends on the acuity and severity of hyponatremia and may present as lethargy, cramping, anorexia, nausea, vomiting, seizures, coma, and death [57]. It is treated by fluid restriction, salt tablets, or V2 receptor antagonists [60]. SIADH can be caused by a several groups of medications.

Diuretics

Thiazide diuretics are one of the most common causes of hyponatremia [61]. The exact mechanism of thiazide-induced hyponatremia is unclear; however, the proposed mechanisms include increased secretion of ADH via direct stimulation [57, 62, 63] or indirectly by inducing hypovolemia. Hypokalemia may also increase osmoreceptor-mediated release of ADH [57, 64]. Old age, female gender, and low body mass are risk factors for the development of thiazide-induced hyponatremia [57, 65].

Antidepressants and Antipsychotics

Antidepressants (SSRI and TCA) and antipsychotics (phenothiazines and butyrophenones) have been associated with hyponatremia associated with SIADH by increasing the release of ADH [57, 58]. The incidence of hyponatremia is highest with SSRIs ranging from 0.5% to 32%. However, care must be taken to rule out primary polydipsia, which is co-prevalent with psychiatric disorders before making the diagnosis of SIADH [66, 67].

Carbamazepine and Oxcarbazepine

Used for partial and generalized seizures, mood disorders, psychotic disorders, and trigeminal neuralgia, carbamazepine and oxcarbazepine can both increase the secretion of ADH. The incidence of hyponatremia with carbamazepine is 4.8–41.5% and correlates with the dose and serum level of the same [58, 68, 69].

Antineoplastic Agents

Cyclophosphamide causes hyponatremia secondary to the release of ADH. Patients on cyclophos-

phamide therapy are advised to drink lots of fluids to prevent the development of hemorrhagic cystitis, and this can lead to severe hyponatremia [70, 71]. Other antineoplastic agents that are associated with SIADH include vincristine, vinblastine, and cisplatin [72–74].

Opiates

Opiates cause SIADH due to direct stimulation of ADH as well as indirect stimulation due to opiate-induced nausea and hypotension [26].

Sulphonylureas

Chlorpropamide, through its direct action to increase ADH production centrally, and stimulation of adenylate cyclase generation or inhibition of cyclic adenosine monophosphate at the level of the kidneys, result in increased ADH effect, and antidiuresis is produced [75].

Impulse Control Disorders

The hypothalamus, along with several other regions of the brain, is involved in controlling behavior. Stimulation of the ventral region of the hypothalamus results in aggression, while the lateral hypothalamic neurons have a role to play in controlling impulses [76, 77]. Impulse control disorders such as pathological gambling, hypersexuality, and compulsive shopping have been reported with the use of dopamine agonist therapy for the treatment of Parkinson's disease, restless leg syndrome, or hyperprolactinemia [78]. Moore et al. studied 1580 events of impulse control disorders and observed a strong association with the use of dopamine agonists. The association was strongest for pramipexole, ropinirole, and aripiprazole, which had a preferential affinity for the D3 receptor [79].

Drug-Induced Hyperthermia

Control of the body temperature is within the hypothalamus. The anterior hypothalamus receives and processes the afferent information, while the posterior hypothalamus initiates

effluent responses based on the input from the anterior hypothalamus. A number of different neurotransmitters, including norepinephrine, dopamine, prostaglandin, and neuropeptides, play an important role in the mediation of temperature [80, 81].

Neuroleptic Malignant Syndrome

Neuroleptic malignant syndrome (NMS) is a rare and potentially fatal complication of drugs that can block dopamine receptors most especially those that block D2 receptors [82]. Hyperthermia, a part of the syndrome, can specifically be explained by the hypothalamic pathway [82]. Antipsychotic medications are the most common cause of NMS. However, it can also occur in association with prochlorperazine, metoclopramide, and droperidol [83–85]. NMS is secondary to the depletion or blockade of dopamine in the hypothalamus, which results in decreased heat loss and increased heat production. The hallmark of NMS is muscular rigidity and hyperthermia with a history of neuroleptic medication. Other symptoms include diaphoresis, dysphagia, tremors, incontinence, mutism, and labile blood pressure. The onset of NMS is typically 24 h after the drug has been taken [86, 87]. Management includes supportive measures to achieve normothermia and administration of bromocriptine and dantrolene [88].

Sympathomimetic Syndrome

The thermoregulatory function of the hypothalamus can be affected by noradrenaline (norepinephrine), serotonin, or dopamine [89–91]. Increased heat production by peripheral acting medication causes a sympathomimetic syndrome that is life-threatening and implicates 3,4-methylenedioxymethamphetamine (MDMA/ecstasy), amphetamine, or cocaine [91]. These sympathomimetic substances are believed to cause alteration in the levels of norepinephrine, dopamine, or 5-hydroxytryptamine, thereby affecting the hypothalamic-pituitary regulation of temperature. Clinical features include agitation, confusion, and hallucinations and quickly progress to status epilepticus and death. Other features include myocardial infarction, disseminated intravascular coagulation (DIC), and renal fail-

ure. Management consists of aggressive cooling, anticonvulsants, and benzodiazepines for agitation [88].

Other Medication-Induced Hypothalamic Disorders

Estradiol treatment has been shown to affect beta-endorphin neurons in the arcuate nucleus selectively [92].

Amphetamines have been found in animals to induce hyperphagia and subsequent obesity via lateral hypothalamic injections as a local neurotoxic effect [93].

Surgical Iatrogenic Hypothalamic Disorders

Iatrogenic hypothalamic injuries during neurosurgical procedures can cause devastating consequences and should be on the forefront of every neurosurgeons' mind when operating within the vicinity of the hypothalamus and its surrounding structures. Although hypothalamic damage is rare, the resulting functional outcome may be devastating and may affect a patient's quality of life [94, 95]. The general mechanisms of complications lie with the surgical approach to the sellar and suprasellar region and the manipulation of the hypothalamus in association with the pituitary gland and the optic structures [95, 96], most especially in cases when mass lesions are strongly adherent to the hypothalamus [94].

Neurosurgical Anatomy Considerations

As a minute structure with a volume of 4 cc containing multiple functional nuclei and fibers, located at the base of the brain [97–99], its preservation during neurosurgical procedures cannot be overstated. More specifically, it is under the thalamus separated by the hypothalamic sulcus of Monroe, lying on the wall and floor of the third ventricle, bordered by the lamina terminalis, a

gray matter layer above the optic chiasm anteriorly, and by the periaqueductal gray matter and the brainstem tegmentum posteriorly [100–102]. It is composed of several distinct nuclei with widespread neural interconnections within the brain and governs many of the human body's homeostatic, regulatory, and motivational behaviors (Fig. 22.1). Along the rostro-caudal axis, the distinct hypothalamic areas based on nuclei-anatomical landmarks include the anterior (supraoptic, pre-optic, or chiasmal) region located above the optic chiasm, medial (tuberal or infundibular) region located above the tuber cinereum, and posterior (mammillary) region [97, 100–102]. Mediolateral divisions include the periventricular, medial, and lateral zones [97, 101, 102]. Generally, the anterior region with the supraoptic nucleus produces vasopressin, while the suprachiasmatic nucleus is involved in the circadian rhythm [100–102]. The medial region is associated with controlling habits of eating and feeling of satiety, secretion of orexigenic peptides, and other motivational behaviors [100–102]. Whereas, the posterior region is involved in feeding behavior, gastrointestinal motility, car-

diovascular regulation, learning, thermoregulation, memory, and emotions [100–102]. This brief anatomical overview demonstrates the value of taking perioperative caution and ultimately for understanding the following outcomes of various hypothalamic brain injuries that can occur intraoperatively.

History of Neurosurgical Approaches

The first transsphenoidal surgery for pituitary surgery was done in 1907 by an Austrian surgeon named Hermann Schloffer [95, 103–106]. This approach leads to the development of the endonasal and sublabial approaches [95, 106, 107]. Subsequent approaches including endonasal transseptal transsphenoidal approach and transnasal/submucosal approach were then performed [95, 104, 106, 108, 109]. A French neurosurgeon named Guiot combined the transsphenoidal approach with intraoperative fluoroscopy and introduced this approach to Jules Hardy, a Canadian surgeon, in 1965 [95, 106]. Hardy then incorporated the operating microscope to the use




Fig. 22.1 Hypothalamus and its nuclei. (With permission Pop et al. [100])

of fluoroscopy [95, 106, 110, 111]. Neurosurgical approach via endoscopy was first utilized in 1806 for paranasal approaches [95, 112] and did not gain ground until the 1990s when there was an improvement in techniques and instrument technology [113]. Jankowski utilized the endoscopic endonasal approach for pituitary adenoma resection for the first time in the 1990s [95, 106, 114–117]. This approach allowed neurosurgeons to have a closer magnified view of the ventricles from within its walls [118]. Its initial use was mostly for hydrocephalus [119], but currently, this approach has expanded to skull base surgery [119, 120]. Carrau et al. reported the first application of neuroendoscopy for skull base tumors with their publication on endonasal transsphenoidal hypophysectomy [118, 121]. The use of the operating microscope improved microsurgical techniques by increased magnification and illumination [120], and as the years passed, in addition to this, new techniques, neuronavigation and intraoperative imaging, broaden the range of lesions that can be resected [120, 122].

Risk of hypothalamic injury during these surgeries is increased with suprasellar extension. This is common with macroadenomas and craniopharyngiomas [95, 123–125]. Although pituitary adenomas are the most common types of tumors approached surgically, the invasive behavior of craniopharyngiomas should be highlighted because several prognostic factors are considered based on hypothalamic involvement preoperatively. Hyperphagia, obesity, and behavioral dysfunction [126–129] are more commonly seen in craniopharyngiomas than as part of pituitary tumor extension. The same is true for malignant tumors of the suprasellar region.

Neurosurgical Approaches

Transcranial Surgery

Historically, transcranial approaches have been the most common approach to surgical resection in the sellar/suprasellar region prior to the introduction of transsphenoidal techniques, initially microscopic and later endoscopic. For pituitary adenomas, the vast majority of cases are now

done transsphenoidally; however, there is still a role for transcranial surgery for other pathologies, such as craniopharyngioma, hypothalamic tumors, and intraventricular tumors. Several transcranial approaches have been described in literature. The most common approaches to the suprasellar region are the subfrontal and pterional (frontosphenotemporal). Transventricular approaches (micro- or endoscopic) are used for resection of intraventricular and primary hypothalamic tumors [130–133]. The pterional approach is the most commonly used transcranial approach [95]. This approach starts with a curvilinear incision from the root of the zygoma following the hairline toward the midline and extended contralaterally as necessary. Most commonly, a myocutaneous flap is elevated, and if further temporal exposure is needed, a sub-fascial dissection can be performed to reflect the temporalis muscle posteriorly. The craniotomy is centered around the “pterion,” which is the confluence of the frontal, sphenoidal, and temporal bones. After the craniotomy is elevated, depending on the sites of the lesion, the Sylvian fissure might or might not need to be opened. Another common approach is the subfrontal, which is similar to the pterional, but the craniotomy advances more into the frontal bone and includes more elevation of the frontal lobe than the temporal side. This exposure is used more for tumors that are purely suprasellar. A progression of these approaches is the orbitozygomatic craniotomy, in which the orbital rim is removed to facilitate superior visualization of tumors with a very large suprasellar component, which are located in the third ventricle. In general, transcranial approaches to the suprasellar region are preferred for tumors such as craniopharyngiomas, meningiomas, and pituitary tumors of fibrous or invasive nature. The degree of suprasellar extension is not so much a limitation for transsphenoidal surgery any more since the advent of endoscopic techniques. More and more tumors that were considered non-operable through a transsphenoidal route can now be resected via an endoscopic transsphenoidal approach. For tumors of the third ventricle and primary tumors for the hypothalamic region, such as gliomas or hamartomas, transventricular

surgery can be considered. This can be accomplished with a microscope or endoscope. The most common microscopic approaches are interhemispheric transcallosal and fronto-transparenchymal approaches. Intraventricular endoscopy can be used to biopsy and resect some types of tumors. Some hypothalamic tumors, such as hamartomas (which are non-neoplastic in nature), can cause intractable seizures, and in those cases, surgical resection can be considered. This area can be approached through transventricular or subfrontal/pterional routes. However, with the present-day introduction of laser interstitial thermal therapy (LITT), these tumors can be treated in a minimally invasive fashion with excellent results [134].

Transsphenoidal Surgery

The transsphenoidal route is the most used for the resection of pituitary adenomas. With the introduction of endoscopy, however, the indications for transsphenoidal surgery have been expanded to multiple pathologies that once could not be treated through this route. Currently, two different techniques are used for transsphenoidal approaches. One includes using the operating microscope and a retractor system to access the sella through the nose. The second one uses endoscopes, which improve visualization and illumination of the sellar region.

Transsphenoidal Microscopic Surgery

Since the introduction of the operative microscope and fluoroscopic guidance in transsphenoidal procedures in the 1960s [111, 135, 136], this technique became the most used among neurosurgeons for resection of sellar tumors. Many surgeons have adopted the use of endoscopes; however, microscopic techniques are still broadly used. Advantages of the microscope include faster access to the sella, three-dimensional view of the anatomy, and ability to use the microsurgical techniques that most neurosurgeons are familiar with. The drawbacks include limited illumination due to poor light penetration, in a narrow field, deep exposure, and limited field of view, which doesn't allow panoramic visualization of the sella [137]. Several studies have

compared outcomes for pituitary adenomas between microscopic and endoscopic techniques, and no significant difference in outcomes has been found for tumors with minimal or no suprasellar extension. Endoscopy does appear to increase the extent of resection for larger tumor with significant suprasellar extension [138].

For the microscopic techniques, the main principle involves creating submucosal access to the sphenoid sinus, which can be accomplished via sublabial or intranasal, septal mucosal incision. The sublabial technique has been for the most part abandoned due to increased patient discomfort and the risk of complications, such as numbness and paresthesias in the upper lip and teeth [139]. In the endonasal approach, the mucosa over the cartilaginous septum is injected with lidocaine with epinephrine, and a small vertical incision is made. The mucosa is then dissected back to the rostrum of the sphenoid bone. A transsphenoidal speculum is introduced, and the vertical plate of the ethmoid is fractured to the contralateral side. This allows for the blades of the speculum to open, thus exposing submucosally the front of the sphenoid sinus. Adequate position of the retractor system can be verified using intraoperative fluoroscopy or navigation. The sphenoid ostia are identified, and the entire front of the sphenoid sinus is removed with Kerrison punches. The top of the vomer bone is removed as well. The sphenoid sinus usually contains one or more septations, which need to be removed. The anatomy is identified based on the preoperative imaging and navigation. In most cases, an infrasellar recess can be identified. It is important to be aware of where the midline is to avoid injury to the carotid or cavernous sinuses when opening the sella. In case of large tumors, the bone of the sella is usually eroded, and upon removal of the sphenoid mucosa, the thin layer of the bone can usually be easily elevated with a dissector. If the bone is thick, a drill might need to be used to open the anterior wall of the sella. Once the bone over the anterior portion of the sella is removed, the dura is cauterized and opened sharply. Pituitary adenomas tend to be soft and can be delivered using ring curettes and suction. For fibrous tumors, an ultrasonic aspira-

tor can be used as well. Once the tumor is removed, the distended diaphragma sellae will drop into the sella under the pressure of the intracranial cerebrospinal fluid. Once this occurs, the surgeon can be confident that the suprasellar tumor has been removed. The tumor must be removed in a specific sequence, inferior and posterior portions first, to prevent early descent of the diaphragm which would obliterate the field of view. If the diaphragm doesn't drop, the surgeon cannot know for sure if the suprasellar portion of the tumor has been completely removed. This is where the use of endoscopes has an advantage, since angled lenses can be used to visualize the suprasellar component of the tumor.

Once the resection is complete, the sella can be packed with fat and reconstructed with a piece of vomer or a synthetic plate. Once the speculum is removed, the mucosa is re-approximated onto the septum, and nasal packing is applied to keep the mucosa in place. If residual suprasellar tumor is identified in postoperative imaging, options include to wait and see if the suprasellar component will eventually descend, perform a second transsphenoidal approach, consider a transcranial approach, or use of endoscopy. If a large portion of tumor is left in the suprasellar space, it can turn into a hemorrhagic mass, causing increased mass effect and edema in the optic apparatus and hypothalamus during the postoperative period, leading to deterioration of the patient [123].

Transsphenoidal Endoscopic Surgery

In the past two decades, there has been a shift toward transsphenoidal endoscopic surgery from microscopic surgery [140]. This approach offers improved illumination and visualization of the anatomy through its panoramic, wide-angle view. The introduction of endoscopy has expanded the indications of transsphenoidal surgery and has even changed the surgical management of some skull base pathologies. The main advantage of the endoscope is improved illumination and visualization with the ability to use angled lenses to “look around the corners,” which, in combination with angled instruments, allows to reach tumors beyond what is possible with a microscope. Disadvantages include loss of three-dimensional

vision, which requires a steeper learning curve from the surgeon, and a more involved access to the sella which often necessitates the collaboration of an ears, nose, and throat (ENT) surgeon. Extended approaches to the sellar and suprasellar region have been described to treat large tumors with suprasellar, hypothalamic, and third ventricular involvement. The main complication seen after resection of these large tumors through a large skull base defect has been cerebrospinal fluid fistula. The development of vascularized nasal septum mucosal flaps has allowed for this complication to become less frequent [141].

The patient is positioned supine with the head elevated above the level of the heart to decrease bleeding. Intraoperative navigation is commonly used (Fig. 22.2). An irrigating sheath helps keep the lens clean.

A zero-degree endoscope is advanced through the nostril. The middle and superior turbinates are outfractured to expand the corridor. The ostium of the sphenoid is identified and expanded. A bi-nostril approach is favored by most surgeons. This involves resection of the posterior portion of the nasal septum, which allows the surgeon to use bimanual technique introducing instruments through both nostrils. A wide sphenoid sinus opening is performed, and the septations in the sinus are thoroughly removed. The endoscope allows for close inspection of the anatomy of the sphenoid sinus. The main ana-



Fig. 22.2 Operating room setup for endoscopic transsphenoidal approach. A bi-nostril approach with posterior septectomy is used. The surgeon is holding a navigated suction on the right hand, showing the anterior wall of the sella. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)




Fig. 22.3 Endoscopic view of the posterior wall of the sphenoid sinus after a wide sphenoidotomy and removal of septations. C clivus, CP carotid prominence, ON optic nerve, OCR optico-carotid recess, PS planum sphenoidale. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)

tomical landmarks are the infrasellar recess, carotid prominences, optic canal prominences, and optico-carotid recesses (Fig. 22.3).

Similar to the microscopic technique, the bone over the sella is removed all the way to the cavernous sinus on either side. For tumors with limited suprasellar extension, the bone removal stops at the level of the tuberculum sellae. If the tumor has a large suprasellar component, the tuberculum is removed and as much planum sphenoidale as needed (Fig. 22.4).

Once the dura is exposed, the position of the carotid arteries can be confirmed using Doppler and/or navigation. The dura is cauterized and opened sharply (Fig. 22.5). In microadenoma surgery, heavy venous bleeding is usually encountered from the intercavernous sinus. This can be controlled packing with hemostatic agents. Adenomas which enlarge the sella usually displace and obliterate the intercavernous sinus, so dural bleeding is usually not an issue.

Pituitary macroadenomas are removed using the same techniques described for microscopic surgery. Resection of microadenomas often requires incision of the normal pituitary (Fig. 22.5). Extracapsular dissection is preferred




Fig. 22.4 Endoscopic view of the sellar region after bone removal. Pit Dura over the anterior pituitary, PS planum sphenoidale, SN suprasellar notch, TS tuberculum sellae. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)




Fig. 22.5 The dura over the pituitary gland has been opened. Note the normal, yellow-colored pituitary gland. Pit pituitary gland, SN suprasellar notch, PS planum sphenoidale. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)

for these tumors, especially when they are functioning, to ensure complete resection [142].

Firm adenomas, craniopharyngiomas, meningiomas, and other tumors of fibrous consistency require a different dissection strategy. In these cases, additional bone is removed, and the dura is




Fig. 22.6 Dissection of the tumor from the optic chiasm. The pituitary gland is being retracted inferiorly. OC optic chiasm, ACA anterior cerebral artery, T tumor. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)

opened to the level of the planum. Sharp dissection along with debulking with an ultrasonic aspirator is used to separate the tumor from the optic apparatus, surrounding vessels, stalk, and third ventricle (Figs. 22.6 and 22.7).

In very large tumors that reach the third ventricle, angled endoscopes are used to inspect the interior of the third ventricle to look for residual tumor and possible obstructions to the CSF circulation (Figs. 22.8 and 22.9).

Closure techniques are tailored to the size of the defect created in the skull base and the integrity of the diaphragm. If the diaphragm is preserved, then the sella can be packed with fat, and a synthetic polymer plate is used as a strut to contain the graft in the sella. If the defect is very large or the CSF fistula observed in surgery is high flow, a vascularized nasoseptal flap can be developed and placed over the defect to facilitate healing.

In cases where the diaphragm is violated and there is communication with the third ventricle, fat packing should not be used as there is risk of migration into the ventricular system resulting in




Fig. 22.7 The tumor has been removed and the interior of the third ventricle is seen. ACAs anterior cerebral arteries, FL frontal lobe, OC optic chiasm, FM foramen of Monro, MI massa intermedia (interthalamic adhesion), H hypothalamus (note raw appearance of the neural tissue after dissection from the tumor). (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)




Fig. 22.8 View of the roof of the third ventricle with a 30-degree endoscope. FM foramen of Monro, CP choroid plexus, MI massa intermedia (interthalamic adhesion), H hypothalamus (note raw appearance of the neural tissue after dissection from the tumor). (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)




Fig. 22.9 View of the posterior third ventricle. Note the raw appearance of the third ventricular walls, corresponding to the hypothalamus (H). The pituitary stalk (PS) has been reduced to a very thin layer, through which the right posterior communicating artery (PCo) can be seen. MI massa intermedia, PR pineal recess, AQ cerebral aqueduct, MB mammillary bodies. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)

ventriculitis or obstruction. In these cases, our preferred closure method is the gasket seal [143].

A fascia lata graft is harvested (synthetic materials can be used as well). A polymer plate is fashioned to match the size of the defect. Then the fascial graft is trimmed to be about 30% larger than the plate. The plate is countersunk over the fascia creating a “gasket” closure. The nasoseptal flap is then placed over the sealed defect. In these cases, a lumbar drain is usually placed and kept open to drain 5–10 ml/h for 48–72 h.

At the end of the procedure, the nasal passages are cleaned from debris, and the turbinates are returned to their position. Minimal packing is used, usually with hemostatic resorbable materials.

Improvement in endoscopic techniques with the use of extended approaches and angled endoscopes and instruments has allowed for transnasal resection of tumors that previously could only be reached transcranially. In experienced hands, the vast majority of pituitary adenomas can be

now removed endoscopically regardless of the degree of suprasellar component. Limitations to this approach include lateral extension which cannot be reached from midline approach, extensive cavernous sinus involvement not accessible endoscopically, and tumors that encase blood vessels, especially if they are fibrous in nature.

Craniopharyngiomas represent a good example of how endoscopic techniques have proven an advantage over the transcranial approach. Extended endoscopic skull base approaches have allowed for improved extent of resection of these tumors with lower morbidity. The transsphenoidal approach allows for the tumor to be removed from its inferior portion, following its pattern of growth and allowing dissection from the optic apparatus and hypothalamus under direct visualization, rather than blind dissection from above typically done with the transcranial approach.

Special Neurosurgical Considerations Regarding Hypothalamic Injury

Preoperative Considerations

Based on magnetic resonance imaging (MRI), hypothalamic involvement can be evaluated based on the degree of involvement [144] (Figs. 22.10 and 22.11). In the case of craniopharyngioma, which is the tumor with the highest incidence of surgery-related hypothalamic injury, several classifications have been created in order to predict outcomes and to tailor the extent of resection in order to minimize injury. In the Müller classification, Grade 0 demonstrates no hypothalamic involvement. Grade 1 involves the anterior hypothalamus not involving mammillary bodies and hypothalamic areas beyond mammillary bodies, and Grade 2 involves both anterior and posterior hypothalamic areas including mammillary bodies and hypothalamic areas beyond mammillary bodies [144]. The grade of hypothalamic involvement has been shown to be associated with postoperative increase in BMI and a decrease in quality of life for these patients with craniopharyngioma [144]. For these reasons, for those with Grade 2 hypothalamic involvement in the setting of child-




Fig. 22.10 Preoperative MRI T1 with contrast sagittal (a) and coronal (b) showing the craniopharyngioma growing in the suprasellar region into the third ventricle. Note

the normal pituitary and normal size of the sella. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)



Fig. 22.11 Preoperative axial non-contrasted CT showing the heavy calcification of this craniopharyngioma. (Courtesy of Roberto Rey Dios, MD, University of Mississippi Medical Center, Jackson, MS)

hood craniopharyngioma, Müller et al. discouraged radical excision surgery [144, 145]. Mortini et al. [146] found that in addition to the Müller classification of mammillary body involvement,

other MRI findings can predict outcomes. Hypothalamic hyperintensity, unidentifiable pituitary stalk, dislocated chiasm, visibility of infundibular recess, ability to recognize the supraoptic recess, and retrochiasmatic tumor extension are associated with the development of postoperative obesity. Peritumoral edema, unidentifiable hypothalamus, hypothalamic compression, and fornix displacement in addition to inability to recognize the supraoptic recess were associated with post-surgical hypothalamic syndrome [146].

Mechanisms of Hypothalamic Injury

Injury to the hypothalamus, infundibulum, and pituitary stalk during surgery can happen through direct or indirect mechanisms. Direct injury is most often a result of accidental resection of hypothalamic tissue adherent to the tumor. This is more likely to occur when tumor dissection is done with suboptimal visualization, leading to traction and tearing of the neural tissue. This tends to happen more often in transcranial approaches for adherent tumors such as craniopharyngioma or dermoid cysts. Transsphenoidal resection of pituitary adenomas is usually done in an intracapsular fashion, without violating the diaphragm and arachnoid, thus resulting in lower risk of neural tissue injury.

Another mechanism of injury is ischemia secondary to vascular injury, vasospasm, or edema. The hypothalamus receives its blood supply from anteromedial branches of the anterior cerebral artery, the posteromedial branches of the posterior communicating artery, and the thalamoperforating branches of the posterior cerebral artery. Venous drainage occurs via the intercavernous sinuses [147].

In tumors requiring extracapsular dissection, avulsion of perforators can lead to immediate ischemia and death of the neural tissue. Ischemia can also occur in a delayed fashion as a result of vasospasm of these tiny perforators in the postoperative period. Severe edema after tumor resection can be seen, for example, when residual tumor is left behind. Arterial supply to the remaining tumor is compromised due to devascularization and venous drainage impaired after surgical manipulation. The result is tumor infarction, often with hemorrhagic transformation (i.e., pituitary tumor apoplexy) resulting in tumor swelling and compression of adjacent structures. The combination of venous congestion in the hypothalamic outflow and inflammatory response from the blood products results in edema, vasospasm, and ischemia of the neural tissue and subsequent hypothalamic injury. This is usually seen within 24–48 h after surgery [148].

Procedure-Related Iatrogenic Hypothalamic Injuries

Radiation Therapy

Radiation therapy (RT) is frequently used in the treatment of tumors of the suprasellar region, either directed to residual/recurrent tumors or as single modality treatment for tumors that cannot be surgically resected [149–152]. Modalities of RT include conventional external beam fractionated radiation therapy (usually delivered in a conformal fashion via intensity-modulated radiation therapy (IMRT)) and stereotactic radiosurgery, which involves higher doses in one to five fractions with a much more precise delivery system.

The goal of the treatment plan is maximum impact on tumor tissue with minimum impact on

“normal” adjacent structures, including the hypothalamus [152]. Complications from RT can occur acutely (within days), early (within weeks), or late (months to years) [152]. Hypothalamic dysfunction can occur early [152]. Hypothalamic syndrome includes obesity or cachexia, impaired thermoregulation, and disturbances of the sleep/wake cycle [152, 144]. Adequate choice of RT modality, optimization of the conformality of the plan, shielding of critical structures, and adequate dose prescription and distribution can prevent such complications.

Deep Brain Stimulation

Deep brain stimulation (DBS) surgery has also been shown to result in iatrogenic hypothalamic injury. DBS surgery is primarily indicated to treat movement disorders, such as essential tremor and Parkinson’s disease, eating disorders, behavior disorders, and pain syndromes [153]. Significant weight gain has been observed in patients with Parkinson’s disease treated with deep brain stimulation using subthalamic implants (DBS-STN) up to 20 kg [154] which have been attributed to a decrease in energy expenditure [155–157]. However, disruption of the hypothalamic centers in charge of energy regulation has not been excluded [154]. Animal studies have implicated DBS of ventromedial hypothalamic nucleus as causing decreased HPA axis function via induction of a systemic inflammatory response [158].

Special Considerations in Neurosurgical-Associated Hypothalamic Dysfunction Manifestations

Central Diabetes Insipidus (CDI)

In central DI, damage to the magnocellular neurons of the supraoptic and paraventricular hypothalamic nuclei results in inadequate production of arginine vasopressin (AVP) or antidiuretic hormone (ADH) to be able to release into the bloodstream by the posterior pituitary [159]. The primary clinical manifestation includes hypotonic polyuria with compensatory polydipsia and may occur as a complication of pituitary surgery

in up to 31% of cases [159–161]. Postoperative DI can follow one of the three courses: transient, permanent, and triphasic. Transient CDI, the most common type of CDI pattern, postoperatively occurs due to dysfunction of magnocellular neurons after surgical manipulation and may occur abruptly within 24–48 h and may last up to 5–10 days [159, 161]. Rarely, 80–90% of AVP-secreting neurons are damaged, and permanent DI may occur [160]. The third possible course is a triple-phase response [161]. The first phase, which is identical clinically to transient DI, begins within 24 h of surgery and typically lasts 4–5 days. This occurs because of absent or decreased ADH release from hypothalamic neuronal shock. Following this initial response, an interphase occurs beginning around 1 week postoperatively and lasts for approximately 1 week. As injured hypothalamic magnocellular neurons degenerate, they release their remaining ADH stores leading to water retention and decreased urine output. In some patients, hyponatremia may develop. This interphase is followed by the final phase whereby permanent DI ensues due to complete degeneration of neurons in the supraoptic and paraventricular hypothalamic nuclei [160, 162–164]. It is very rare, and only 1–3% of patients develop this triphasic response postoperatively from hypothalamic injury.

Hypothalamic Obesity Syndrome with Hyperphagia

HOS is characterized by a rapid, unrelenting weight gain that may be accompanied by severe hyperphagia due to hypothalamic damage or injury and in association with other hormonal imbalance attributed to the hypothalamic-pituitary function [154]. The pathophysiology is based around many key nuclei within the hypothalamus. Investigators found that lesions to the hypothalamic ventromedial area, paraventricular nucleus, or dorsal medial hypothalamic nuclei produced hyperphagia and obesity in rats, whereas lesions of the lateral hypothalamus resulted in hypophagia [154]. These findings gave rise to the hypothesis that the ventromedial area and the lateral hypothalamus can be categorized as the satiety and hunger centers, respec-

tively [154]. The arcuate nucleus located in the base of the hypothalamus primarily hosts the neurons expressing proopiomelanocortin (POMC) and neuropeptide-Y (NPY) [154]. POMC inhibits feeding and increases energy expenditure, whereas NPY stimulates feeding and decreases energy expenditure [154]. Both of the hormones leptin and insulin increase the expression of POMC and decrease the expression of NPY [154]. Preoperatively, imaging findings showing tumor and hypothalamic involvement have been found to be associated with postoperative obesity [145, 146]. This syndrome has been reported to potentially be associated with deep brain stimulation as well [154].

Sympathetic Dysfunction

In a case series, refractory hypotension as a manifestation of hypothalamic injury postoperatively can occur in approximately 1.5% [165]. Systemic hemodynamic response has been attributed to the paraventricular nuclei and zone AV₃V of the hypothalamus [165–169]. Although very uncommon, neurosurgeons should be aware of this phenomenon that can occur after sellar tumor surgery and be prepared to act quickly with sympathomimetic drugs [165].

Acquired Central Hypoventilation

“Acquired central hypoventilation” is exceedingly rare, and there are very few reports of this from injury to neurological structures after surgery. One case report presented a 15-year-old female who underwent a right-sided pterional craniotomy for resection of craniopharyngioma [170]. She had her first surgery complicated by permanent DI. She presented 2 months after with worsening vision and recurrence of the cystic component of the craniopharyngioma. An orbitopterional approach craniotomy was performed; however, the evening after the surgery, an attempted extubation failed even with the patient being awake and following commands. Ventilatory failure persisted until postoperative day 14. The cause of central hypoventilation in this mentioned case report was attributed to hypothalamic dysfunction. Diagnostics done included endocrinological testing, liver function

tests, chest radiographs, and repeat MRI as this was a diagnosis of exclusion. The paraventricular nucleus of the hypothalamus has been previously studied to project to medullary and spinal cord respiratory centers [170]. Leptin, a hormone produced by the adipose tissues and that acts on hypothalamic receptors, has been implicated to hypoventilation as well [170, 171].

Intraoperative Hyperthermia

A case report described an intraoperative occurrence of intraoperative hyperthermia to 40.1 °C within 20 min as the surgery for craniopharyngioma excision was ending [172]. There was no noted deviation from their standard induction or maintenance anesthetic. They managed the patient as malignant hyperthermia and discontinued inhalational anesthetic and administered paracetamol, propofol, and intravenous cold saline. Hypernatremia and an increase in urine output to 25 mL/kg/hour were noted, and DI was suspected. Diagnostic workup for possible causes was done including infection, neuroleptic malignant syndrome, thyroid dysfunction, or transfusion reaction. Due to the temporal association of the resection and tumor manipulation with the hyperthermia, this event was attributed to hypothalamic affection [172]. Anticipation of this potentially lethal complication should be in every surgical planning.

References

1. Peuskens J, Pani L, Detraux J, De Hert M. The effects of novel and newly approved antipsychotics on serum prolactin levels: a comprehensive review. *CNS Drugs*. 2014;28(5):421–53.
2. Halbreich U, Kinon BJ, Gilmore JA, Kahn LS. Elevated prolactin levels in patients with schizophrenia: mechanisms and related adverse effects. *Psychoneuroendocrinology*. 2003;28(Suppl 1):53–67.
3. La Torre D, Falorni A. Pharmacological causes of hyperprolactinemia. *Ther Clin Risk Manag*. 2007;3(5):929–51.
4. Molitch ME. Disorders of prolactin secretion. *Endocrinol Metab Clin N Amer*. 2001;30:585–610.
5. Molitch ME. Medication-induced hyperprolactinemia. *Mayo Clin Proc*. 2005;80(8):1050–7.
6. Gruen PG, Sachar EJ, Altman N, Langer G, Tabrizi MA, Halpern FS. Relation of plasma prolactin to clinical response in schizophrenic patients. *Arch Gen Psychiatry*. 1978;35(10):1222–7.
7. Green AI, Brown WA. Prolactin and neuroleptic drugs. *Endocrinol Metab Clin North Am*. 1988;17(1):213–23.
8. Wieck A, Haddad P. Hyperprolactinaemia caused by antipsychotic drugs. *BMJ*. 2002;324:250–2.
9. Maguire A. Prolactin elevation with antipsychotic medications: mechanisms of action and clinical consequences. *J Clin Psychiatry*. 2000;63(Suppl 4):56–62.
10. Schlösser R, Gründer G, Anghelescu I, Hillert A, Ewald-Gründer S, Hiemke C, Benkert O. Long-term effects of the substituted benzamide derivative amisulpride on baseline and stimulated prolactin levels. *Neuropsychobiology*. 2002;46(1):33–40.
11. Kühn KU, Meyer K, Maier W. Flupenthixol—a partial atypical neuroleptic? *Fortschr Neurol Psychiatr*. 2000;68(Suppl 1):S38–41.
12. Seeman P. Atypical antipsychotics: mechanism of action. *Can J Psychiatry*. 2002;47(1):27–38.
13. Lieberman JA. Dopamine partial agonists: a new class of antipsychotic. *CNS Drugs*. 2004;18(4):251–67.
14. McKeage K, Plosker GL. Amisulpride: a review of its use in the management of schizophrenia. *CNS Drugs*. 2004;18(13):933–56.
15. Kleinberg DL, Davis JM, deCoster R, et al. Prolactin levels and adverse events in patients treated with risperidone. *J Clin Psychopharmacol*. 1999;19:57–61.
16. Kinon BJ, Gilmore JA, Liu H, et al. Prevalence of hyperprolactinemia in schizophrenic patients treated with conventional antipsychotic medications or risperidone. *Psychoneuroendocrinology*. 2003;28:55–68.
17. Kearns AE, Goff DC, Hayden DL, et al. Risperidone associated hyperprolactinemia. *Endocr Pract*. 2003;6:425–9.
18. Volavka J, Czobor P, Cooper TB, et al. Prolactin levels in schizophrenia and schizoaffective disorder patients treated with clozapine, olanzapine, risperidone, or haloperidol. *J Clin Psychiatry*. 2004;65:57–61.
19. Breier AF, Malhotra AK, Su T-P, et al. Clozapine and risperidone in chronic schizophrenia: effects on symptoms, parkinsonian side effects, and neuroendocrine response. *Am J Psychiatry*. 1999;156:294–8.
20. Kim K-S, Pae C-U, Hae J-H, et al. Effects of olanzapine on prolactin levels of female patients treated with schizophrenia treated with risperidone. *J Clin Psychiatry*. 2002;63:408–13.
21. Goff DC, Posever T, Herz L, et al. An exploratory haloperidol-controlled dose-finding study of ziprasidone in hospitalized patients with schizophrenia or schizoaffective disorder. *J Clin Psychopharmacol*. 1998;18:296–304.
22. Casey DE, Carson WH, Saha AR, Aripiprazole Study Group, et al. Switching patients to aripiprazole from other antipsychotic agents: a multicenter randomized study. *Psychopharmacology (Berl)*. 2003;166:391–39.

23. Misra M, Papakostas GI, Klibanski A. Effects of psychiatric disorders and psychotropic medications on prolactin and bone metabolism. *J Clin Psychiatry*. 2004;65:1607–18.
24. Grunder G, Carlsson A, Wong DF. Mechanism of new antipsychotic medications. Occupancy is not just antagonism. *Arch Gen Psychiatry*. 2003;60:974–7.
25. Wieck A, Haddad PM. Antipsychotic-induced hyperprolactinaemia in women: pathophysiology, severity and consequences. Selective literature review. *Br J Psychiatry*. 2003;182:199–204.
26. Jones RB, Luscombe DK, Groom GV. Plasma prolactin concentrations in normal subjects and depressive patients following oral clomipramine. *Postgrad Med J*. 1977;53(Suppl 4):166–71.
27. Slater SL, Lipper S, Shiling DJ, et al. Elevation of plasma prolactin by monoamine-oxidase inhibitors. *Lancet*. 1977;2:275–6.
28. Peterson MC. Reversible galactorrhea and prolactin elevation related to fluoxetine use. *Mayo Clin Proc*. 2001;76:215–6.
29. Shin SH, Obonsawin MC, Van Vugt DA, Baby N, Jhamandas K. Morphine can stimulate prolactin release independent of a dopaminergic mechanism. *Can J Physiol Pharmacol*. 1988;66(11):1381–5.
30. Gudelsky GA, Porter JC. Morphine- and opioid peptide induced inhibition of the release of dopamine from tuberoinfundibular neurons. *Life Sci*. 1979;25:1697–702.
31. Lafuente A, Marco J, Esquifino AI. Opioids and the pulsatile prolactin secretory pattern: effects of hyperprolactinemia. *Vet Hum Toxicol*. 1994;36(6):524–8.
32. Bart G, Borg L, Schluger JH, Green M, Ho A, Kreek MJ. Suppressed prolactin response to dynorphin A1–13 in methadone-maintained versus control subjects. *J Pharmacol Exp Ther*. 2003;306(2):581–7.
33. Tamagna EI, Lane W, Hershman JM, Carlson HE, Sturdevant RA, Poland RE, Rubin RT. Effect of chronic metoclopramide therapy on serum pituitary hormone concentrations. *Horm Res*. 1979;11(4):161–9.
34. Fujino T, Kato H, Yamashita S, Aramaki S, Morioka H, Koresawa M, Miyauchi F, Toyoshima H, Torigoe T. Effects of domperidone on serum prolactin levels in human beings. *T Endocrinol Jpn*. 1980;27(4):521–5.
35. Steiner J, Cassar J, Mashiter K, Dawes I, Fraser TR, Breckenridge A. Effects of methyl dopa on prolactin and growth hormone. *Br Med J*. 1976;1(6019):1186–8.
36. Lee PA, Kelly MR, Wallin JD. Increased prolactin levels during reserpine treatment of hypertensive patients. *JAMA*. 1976;235(21):2316–7.
37. Kelley SR, Kamal TJ, Molitch ME. Mechanism of verapamil calcium channel blockade-induced hyperprolactinemia. *Am J Physiol*. 1996;270(1 Pt 1):E96–100.
38. Romeo JH, Dombrowski R, Kwak YS, Fuehrer S, Aron DC. Hyperprolactinaemia and verapamil: prevalence and potential association with hypogonadism in men. *Clin Endocrinol (Oxf)*. 1996;45(5):571–5.
39. Barbieri C, Larovere MT, Mariotti G, Ferrari C, Caldara R. Prolactin stimulation by intravenous labetalol is mediated inside the central nervous system. *Clin Endocrinol (Oxf)*. 1982;16(6):615–9.
40. Perret G, Hugues JN, Louchahi M, Varoquaux O, Modigliani E. Effect of a short-term oral administration of cimetidine and ranitidine on the basal and thyrotropin-releasing hormone-stimulated serum concentrations of prolactin, thyrotropin and thyroid hormones in healthy volunteers. A double-blind cross-over study. *Pharmacology*. 1986;32(2):101–8.
41. Knigge UP. Histaminergic regulation of prolactin secretion. *Dan Med Bull*. 1990;37(2):109–24.
42. Morosini PP, Campanella N, Ghirelli S, Pellegrini F, Testa I, De Martinis C. Cimetidine and hyperprolactinemia. *Boll Soc Ital Biol Sper*. 1979;55(1):14–7.
43. Sibilia V, Netti C, Guidobono F, Pagani F, Pecile A. Cimetidine-induced prolactin release: possible involvement of the GABA-ergic system. *Neuroendocrinology*. 1985;40(3):189–92.
44. Risch SC, Janowsky DS, Siever LJ, Judd LJ, Rausch JL, Huey LY, Beckman KA, Cohen RM, Murphy DL. Correlated cholinomimetic-stimulated beta-endorphin and prolactin release in humans. *Peptides*. 1982;3(3):319–22.
45. De Felice F, Pranno N, Marampon F, Musio D, Salducci M, Polimeni A, Tombolini V. Immune check-point in glioblastoma multiforme. *Crit Rev Oncol Hematol*. 2019;138:60–9.
46. De Felice F, Marchetti C, Palaia I, Musio D, Muzii L, Tombolini V, Panici PB. Immunotherapy of ovarian cancer: the role of checkpoint inhibitors. *J Immunol Res*. 2015;2015:191832.
47. De Felice F, Musio D, Cassese R, Gravina GL, Tombolini V. New approaches in glioblastoma multiforme: the potential role of immune-check point inhibitors. *Curr Cancer Drug Targets*. 2017;17(3):282–9.
48. Buchbinder EI, Desai A. CTLA-4 and PD-1 pathways: similarities, differences, and implications of their inhibition. *Am J Clin Oncol*. 2016;39(1):98–106.
49. Girotra M, Hansen A, Farooki A, Byun D, Min L, Creelan BC, Callahan MK, Atkins MB, Sharon E, Antonia SJ, West P, Gravell AE, Investigational Drug Steering Committee (IDSC) Immunotherapy Task Force Collaboration. The current understanding of the endocrine effects from immune checkpoint inhibitors and recommendations for management. *JNCI Cancer Spectr*. 2018;2(3):pky021.
50. González-Rodríguez E, Rodríguez-Abreu, on behalf of the Spanish Group for Cancer Immunotherapy (GETICA). Immune checkpoint inhibitors: review and management of endocrine adverse events. *Oncologist*. 2016;21(7):804–16.
51. Neary N, Nieman L. Adrenal insufficiency: etiology, diagnosis and treatment. *Curr Opin Endocrinol Diabetes Obes*. 2010;17:217–23.
52. de Filette J, Andreescu CE, Cools F, Bravenboer B, Velkeniers B. A systematic review and meta-analysis of endocrine-related adverse events associated with

- immune checkpoint inhibitors. *Horm Metab Res.* 2019;51(3):145–56.
53. Byun DJ, Wolchok JD, Rosenberg LM, Girotra M. Cancer immunotherapy – immune checkpoint blockade and associated endocrinopathies. *Nat Rev Endocrinol.* 2017;13(4):195–207.
 54. Iwama S, De Remigis A, Callahan MK, Slovin SF, Wolchok JD, Caturegli P. Pituitary expression of CTLA-4 mediates hypophysitis secondary to administration of CTLA-4 blocking antibody. *Sci Transl Med.* 2014;6(230):230ra45.
 55. Faje AT, Sullivan R, Lawrence D, et al. Ipilimumab-induced hypophysitis: a detailed longitudinal analysis in a large cohort of patients with metastatic melanoma. *J Clin Endocrinol Metab.* 2014;99:4078–85.
 56. Albarel F, Gaudy C, Castinetti F, et al. Long-term follow-up of ipilimumab-induced hypophysitis, a common adverse event of the anti-CTLA-4 antibody in melanoma. *Eur J Endocrinol.* 2015;172:195–204.
 57. Chan TY. Drug-induced syndrome of inappropriate antidiuretic hormone secretion. Causes, diagnosis and management. *Drugs Aging.* 1997;11(1):27–44.
 58. Shepshelovich D, Schechter A, Calvarysky B, Diker-Cohen T, Rozen-Zvi B, Gafer-Gvili A. Medication-induced SIADH: distribution and characterization according to medication class. *Br J Clin Pharmacol.* 2017;83(8):1801–7.
 59. Bartter FC, Schwartz WB. The syndrome of inappropriate secretion of antidiuretic hormone. *Am J Med.* 1967;42:790–806.
 60. Liamis G, Milionis H, Elisaf M. A review of drug-induced hyponatremia. *Am J Kidney Dis.* 2008;52(1):144–53.
 61. Spital A. Diuretic-induced hyponatremia. *Am J Nephrol.* 1999;19:447–52.
 62. Ghose RR. Plasma arginine vasopressin in hyponatraemic patients receiving diuretics. *Postgrad Med J.* 1985;61:1043–6.
 63. Sonnenblick M, Rosin AI. Significance of the measurement of uric acid fractional clearance in diuretic induced hyponatraemia. *Postgrad Med J.* 1986;62:449–52.
 64. Fichman MP, Vorherr H, Kleeman CR, et al. Diuretic-induced hyponatremia. *Ann Intern Med.* 1971;75:853–63.
 65. Clark BA, Shannon RP, Rosa RM, Epstein FH. Increased susceptibility to thiazide-induced hyponatremia in the elderly. *J Am Soc Nephrol.* 1994;5:1106–11.
 66. Jose CJ, Perez-Cruet J. Incidence and morbidity of self-induced water intoxication in state mental hospital patients. *Am J Psychiatry.* 1979;136:221–2.
 67. Jacob S, Spinler SA. Hyponatremia associated with selective serotonin-reuptake inhibitors in older adults. *Ann Pharmacother.* 2006;40:1618–22.
 68. Van Amelsvoort T, Bakshi R, Devaux CB, Schwabe S. Hyponatremia associated with carbamazepine and oxcarbazepine therapy: a review. *Epilepsia.* 1994;35:181–8.
 69. Nielsen OA, Johannessen AC, Bardrum B. Oxcarbazepine-induced hyponatremia, a cross-sectional study. *Epilepsy Res.* 1988;2:269–71.
 70. DeFronzo RA, Braine H, Colvin M, Davis PJ. Water intoxication in man after cyclophosphamide therapy. Time course and relation to drug activation. *Ann Intern Med.* 1973;78:861–9.
 71. Bressler RB, Huston DP. Water intoxication following moderate-dose intravenous cyclophosphamide. *Arch Intern Med.* 1985;145:548–9.
 72. Robertson GL, Bhoopalam N, Zelkowitz LJ. Vincristine neurotoxicity and abnormal secretion of antidiuretic hormone. *Arch Intern Med.* 1973;132:717–20.
 73. RaviKumar TS, Grage TB. The syndrome of inappropriate antidiuretic hormone secretion secondary to vinblastine-bleomycin therapy. *J Surg Oncol.* 1983;24:242–5.
 74. Giaccone G, Donadio M, Ferrati P, et al. Disorders of serum electrolytes and renal function in patients treated with cis-platinum on an outpatient basis. *Eur J Cancer Clin Oncol.* 1985;21:433–7.
 75. Moses AM, Numann P, Miller M. Mechanism of chlorpropamide-induced anti-diuresis in man: evidence for release of ADH and enhancement of peripheral action. *Metabolism.* 1973;22:59–66.
 76. Noble EE, Wang Z, Liu CM, Davis EA, Suarez AN, Stein LM, Tsan L, Terrill SJ, Hsu TM, Jung AH, Raycraft LM, Hahn JD, Darvas M, Cortella AM, Schier LA, Johnson AW, Hayes MR, Holschneider DP, Kanoski SE. Hypothalamus-hippocampus circuitry regulates impulsivity via melanin-concentrating hormone. *Nat Commun.* 2019;10(1):4923.
 77. Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P. Functional identification of an aggression locus in the mouse hypothalamus. *Nature.* 2011;470:221–6.
 78. Kelley BJ, Duker AP, Chiu P. Dopamine agonists and pathologic behaviors. *Parkinsons Dis.* 2012;2012:603631.
 79. Moore TJ, Glenmullen J, Mattison DR. Reports of pathological gambling, hypersexuality, and compulsive shopping associated with dopamine receptor agonist drugs. *JAMA Intern Med.* 2014;174(12):1930–3.
 80. Kurz A. Physiology of thermoregulation. *Best Pract Res Clin Anaesthesiol.* 2008;22:627–44.
 81. Buggy DJ, Crossley AW. Thermoregulation, mild perioperative hypothermia and postanaesthetic shivering. *Br J Anaesth.* 2000;84:615–28.
 82. Pileggi DJ, Cook AM. Neuroleptic malignant syndrome. *Ann Pharmacother.* 2016;50(11):973–81.
 83. Pesola GR, Quinto C. Prochlorperazine- induced neuroleptic malignant syndrome. *J Emerg Med.* 1996;14:727–9.
 84. Nonino F, Campomori A. Neuroleptic malignant syndrome associated with metoclopramide. *Ann Pharmacother.* 1999;33:644–5.

85. Ratan DA, Smith AH. Neuroleptic malignant syndrome secondary to droperidol. *Biol Psychiatry*. 1993;34:421–2.
86. Levenson JL. Neuroleptic malignant syndrome. *Am J Psychiatry*. 1985;142:1137–45.
87. Gurrera RJ. Sympathoadrenal hyperactivity and the etiology of neuroleptic malignant syndrome. *Am J Psychiatry*. 1999;156:169–8.
88. Musselman ME, Saely S. Diagnosis and treatment of drug-induced hyperthermia. *Am J Health Syst Pharm*. 2013;70(1):34–42.
89. Hadad E, Weinbroum AA, Ben-Abraham R. Drug-induced hyperthermia and muscle rigidity: a practical approach. *Eur J Emerg Med*. 2003;10(2):149–54.
90. Eyer F, Zilker T. Bench-to-bedside review: mechanisms and management of hyperthermia due to toxicity. *Crit Care*. 2007;11(6):236.
91. Jamshidi N, Dawson A. The hot patient: acute drug-induced hyperthermia. *Aust Prescr*. 2019;42(1):24–8.
92. Wener KM, Morales CR, Brawer JR. The effect of estradiol-induced hypothalamic pathology on sulfated glycoprotein-2 (clusterin) expression in the hypothalamus. *Brain Res*. 1997;745(1–2):37–45.
93. Hernandez L, Parada M, Hoebel BG. Amphetamine-induced hyperphagia and obesity caused by intraventricular or lateral hypothalamic injections in rats. *J Pharmacol Exp Ther*. 1983;227(2):524–30.
94. Yano S, Hide T, Shinojima N. Surgical outcomes of endoscopic endonasal skull base surgery of craniopharyngiomas evaluated according to the degree of hypothalamic extension. *World Neurosurg*. 2017;100:288–96.
95. Theodros D, Patel M, Ruzevick J, Lim M, Bettgeowda C. Pituitary adenomas: historical perspective, surgical management and future directions. *CNS Oncol*. 2015;4(6):411–29.
96. Ciric I, Ragin A, Baumgartner C, Pierce D. Complications of transsphenoidal surgery: results of a national survey, review of the literature, and personal experience. *Neurosurgery*. 1997;40(2):225–36; discussion 236–227.
97. Barbosa DN, de Oliveira-Souza R, Monte Santo F, de Oliveira Faria A, Gorgulho AA, De Salles AF. The hypothalamus at the crossroads of psychopathology and neurosurgery. *Neurosurg Focus*. 2017;43(3):E15. Retrieved Jan 7, 2020, from <https://thejns.org/focus/view/journals/neurosurg-focus/43/3/article-pE15.xml>.
98. Lemaire JJ, Frew AJ, McArthur D, Gorgulho AA, Alger JR, Salomon N, Chen C, Behnke EJ, De Saalles AA. White matter connectivity of human hypothalamus. *Brain Res*. 2011;1371:43–64.
99. Lemaire JJ, Nezzar H, Sakka L, Boirie Y, Fontaine D, Coste A, Coll G, Sontheimer A, Sarret C, Gabrillargues J, De Salles A. Maps of the adult human hypothalamus. *Surg Neurol Int*. 2013;4(Suppl 3):S156–63.
100. Pop MG, Crivii C, Opincariu I. Anatomy and function of the hypothalamus, hypothalamus in health and diseases. In: Baloyannis SJ, Gordeladze JO, editors. *IntechOpen*. November 5th 2018. <https://doi.org/10.5772/intechopen.80728>. Available from: <https://www.intechopen.com/books/hypothalamus-in-health-and-diseases/anatomy-and-function-of-the-hypothalamus>.
101. Goiveia FV. Amygdala and hypothalamus. The human hypothalamus. Basic and clinical aspects. Part II: neuropathology of the hypothalamus and adjacent brain structures. In: Dick S, editor. *Handbook of clinical neurology*, vol. 80. Amsterdam: Elsevier; 2004. p. 597.
102. Swaab DF. The human hypothalamus. Basic and clinical aspects. Part II: neuropathology of the hypothalamus and adjacent brain structures. In: Dick S, editor. *Handbook of clinical neurology*, vol. 80. Amsterdam: Elsevier; 2004. p. 597.
103. Wilson CB, Rand RW, Grollmus JM, Heurser G, Levin S, Goldfield E, Schneider V, Linfoot J, Hosobuchi Y. Surgical experience with a microscopic transsphenoidal approach to pituitary tumors and non-neoplastic parasellar conditions. *Calif Med*. 1972;117(5):1–9.
104. Grosvenor AE, Laws ER. The evolution of extracranial approaches to the pituitary and anterior skull base. *Pituitary*. 2008;11(4):337–45.
105. Lindholm J. A century of pituitary surgery: Schloffer's legacy. *Neurosurgery*. 2007;61(4):865–7; discussion 867–868.
106. Cavallo LM, Somma T, Solari D, Iannuzzo G, Frio F, Baiano C, Cappabianca P. Endoscopic endonasal transsphenoidal surgery: history and evolution. *World Neurosurg*. 2019;127:686–94.
107. Jane JA Jr, Han J, Prevedello DM, Jagannathan J, Dumont AS, Laws ER Jr. Perspectives on endoscopic transsphenoidal surgery. *Neurosurg Focus*. 2005;19(6):E2.
108. Joshi SM, Cudlip S. Transsphenoidal surgery. *Pituitary*. 2008;11(4):353–60.
109. Liu JK, Cohen-Gadol AA, Laws ER Jr, Cole CD, Kan P, Couldwell WT. Harvey Cushing and Oskar Hirsch: early forefathers of modern transsphenoidal surgery. *J Neurosurg*. 2005;103(6):1096–104.
110. Cappabianca P, de Divitiis E. Back to the Egyptians: neurosurgery via the nose. A five-thousand year history and the recent contribution of the endoscope. *Neurosurg Rev*. 2007;30(1):1–7; discussion 7.
111. Hardy J, Wigser SM. Trans-sphenoidal surgery of pituitary fossa tumors with televised radiofluoroscopic control. *J Neurosurg*. 1965;23(6):612–9.
112. Reuter M. The historical development of endophotography. *World J Urol*. 2000;18(4):299–302.
113. Sgouros S. Man-to-machine interface in neuroendoscopy: the importance of human interface in the development of neuroendoscopy. In: Sgouros S, editor. *Neuroendoscopy*. Berlin, Heidelberg: Springer; 2014.
114. Dubourg J, Jouanneau E, Messerer M. Pituitary surgery: legacies from the past. *Acta Neurochir*. 2011;153(12):2397–402.

115. Jankowski R, Auque J, Simon C, Marchal JC, Hepner H, Wayoff M. Endoscopic pituitary tumor surgery. *Laryngoscope*. 1992;102(2):198–202.
116. Rodziewicz GS, Chuang WC. Endoscopic removal of organized chronic subdural hematoma. *Surg Neurol*. 1995;43(6):569–72; discussion 572–563.
117. Sethi DS, Pillay PK. Endoscopic management of lesions of the sella turcica. *J Laryngol Otol*. 1995;109(10):956–62.
118. Li K, Nelson C, Suk I, Jallo G. Neuroendoscopy: past, present, and future. *Neurosurg Focus*. 2005;19(6):1092–0684.
119. Azab WA. Neuroendoscopy in Kuwait: evolution, current status, and future directions. *World Neurosurg*. Elsevier. 2016;82:1878–8750.
120. Conrad JJ. Binostril versus mononostril approaches in endoscopic transsphenoidal pituitary surgery: clinical evaluation and cadaver study. *J Neurosurg*. 2016;125(2):0022–3085.
121. Carrau RL, Jho HD, Ko Y. Transnasal-transsphenoidal endoscopic surgery of the pituitary gland. *Laryngoscope*. 1996;106:914–8.
122. Cappabianca P, Decq P, Schroeder HW. Future of endoscopy in neurosurgery. *Surg Neurol*. 2007;67:496–8.
123. Zada G, Du R, Laws ER Jr. Defining the “edge of the envelope”: patient selection in treating complex sellar-based neoplasms via transsphenoidal versus open craniotomy. *J Neurosurg*. 2011;114(2):286–300.
124. Decker RE, Chalif DJ. Progressive coma after the transsphenoidal decompression of a pituitary adenoma with marked suprasellar extension: report of two cases. *Neurosurgery*. 1991;28(1):154–7; discussion 157–158.
125. Puget S, Garnett M, Wray A, et al. Pediatric craniopharyngiomas: classification and treatment according to the degree of hypothalamic involvement. *J Neurosurg*. 2007;106(1 Suppl):3–12.
126. Müller HL, Gebhardt U, Etavard-Gorris N, Korenke E, Warmuth-Metz M, Kolb R, Sorensen N, Calaminus G. Prognosis and sequela in patients with childhood craniopharyngioma—results of HIT-ENDO and update on KRANIOPHARYNGEOM 2000. *Klin Padiatr*. 2004;216(6):343–8. <https://doi.org/10.1055/s-2004-832339>.
127. de Vile CJ, Grant DB, Hayward RD, Kendall BE, Neville BG, Stanhope R. Obesity in childhood craniopharyngioma: relation to post-operative hypothalamic damage shown by magnetic resonance imaging. *J Clin Endocrinol Metab*. 1996;81(7):2734–7.
128. Merchant TE, Kiehna EN, Sanford RA, Mulhern RK, Thompson SJ, Wilson MW, Lustig RH, Kun LE. Craniopharyngioma: the St. Jude Children’s Research Hospital experience 1984–2001. *Int J Radiat Oncol Biol Phys*. 2002;53(3):533–42.
129. Poretti A, Grotzer MA, Ribi K, Schonle E, Boltshauser EA. Outcome of craniopharyngioma in children: long-term complications and quality of life. *Dev Med Child Neurol*. 2004;46(4):220–9.
130. Gerganov V, Metwali H, Samii A, Fahlbusch R, Samii M. Microsurgical resection of extensive craniopharyngiomas using a frontolateral approach: operative technique and outcome. *J Neurosurg*. 2014;120:559–70.
131. Hori T, Kawamata T, Amano K, Aihara Y, Ono M, Miki N. Anterior interhemispheric approach for 100 tumors in and around the anterior third ventricle. *Neurosurgery*. 2010;66:65–74.
132. Kunihiro N, Goto T, Ishibashi K, Ohata K. Surgical outcomes of the minimum anterior and posterior combined transpetrosal approach for resection of retrochiasmatic craniopharyngiomas with complicated conditions. *J Neurosurg*. 2014;120:1–11.
133. Matsuo T, Kamada K, Izumo T, Nagata I. Unilateral basal interhemispheric approach through the sphenoid sinus to retrochiasmatic and intrasellar craniopharyngiomas: surgical technique and results. *World Neurosurg*. 2014;82:799–805.
134. David SX, Chen T, Hlubek RJ, Bristol RE, Smith KA, Ponce FA, Kerrigan JF, Nakaji P. Magnetic resonance imaging-guided laser interstitial thermal therapy for the treatment of hypothalamic hamartomas: a retrospective review. *Neurosurgery*. 2018;83(6):1183–92. <https://doi.org/10.1093/neuros/nyx604>.
135. Hardy J. Transsphenoidal microsurgery of the normal and pathological pituitary. *Clin Neurosurg*. 1969;16:185–217.
136. Hardy J. Transsphenoidal hypophysectomy. 1971. *J Neurosurg*. 2007;107(2):458–71.
137. Louis RG, Eisenberg A, Barkhoudarian G, Griffiths C, Kelly DF. Evolution of minimally invasive approaches to the sella and parasellar region. *Int Arch Otorhinolaryngol*. 2014;18(Suppl. 2):136–48. <https://doi.org/10.1055/s-0034-1395265>.
138. Singh H, Essayed WI, Cohen-Gadol A, Zada G, Schwartz TH. Resection of pituitary tumors: endoscopic versus microscopic. *J Neurooncol*. 2016;130(2):309–17.
139. Er U, Gurses L, Saka C, et al. Sublabial transseptal approach to pituitary adenomas with special emphasis on rhinological complications. *Turk Neurosurg*. 2008;18(4):425–30.
140. Rolston JD, Han SJ, Aghi MK. Nationwide shift from microscopic to endoscopic transsphenoidal pituitary surgery. *Pituitary*. 2016;19(3):248–50.
141. Wardas P, Tymowski M, Piotrowska-Seweryn A, Markowski J, Ładziński P. Hadad-Bassagasteguy flap in skull base reconstruction - current reconstructive techniques and evaluation of criteria used for qualification for harvesting the flap. *Wideochirurgia i inne techniki maloinwazyjne/Videosurg Other Miniinvasive Tech*. 2019;14(2):340–7. <https://doi.org/10.5114/wiitm.2018.79633>.

142. Taylor DG, Jane JA, Oldfield EH. Resection of pituitary macroadenomas via the pseudocapsule along the posterior tumor margin: a cohort study and technical note. *J Neurosurg.* 2018;128(2):422–8.
143. Garcia-Navarro V, Anand VK, Schwartz TH. Gasket seal closure for extended endonasal endoscopic skull base surgery: efficacy in a large case series. *World Neurosurg.* 2013;80(5):563–8.
144. Müller HL, Gebhardt U, Teske C, Faldum A, Zwiener I, Warmuth-Metz M, Pietsch T, Pohl F, Sörensen N, Calaminus G, Study Committee of Kraniopharyngeom 2000. Post-operative hypothalamic lesions and obesity in childhood craniopharyngioma: results of the multinational prospective trial Kraniopharyngeom 2000 after 3-year follow-up. *Eur J Endocrinol.* 2011;165(1):17–24.
145. Müller HL. Craniopharyngioma: long-term consequences of a chronic disease. *Expert Rev Neurother.* 2015;15(11):1241–4.
146. Mortini P, Gagliardi F, Bailo M, Spina A, Palangeli A, Falini A, Losa M. Magnetic resonance imaging as predictor of functional outcome in craniopharyngiomas. *Endocrine.* 2016;51(1):148–62.
147. Rhoton AL Jr. The sellar region. *Neurosurgery.* 2002;51(4 Suppl):S335–74. Review. PubMed PMID: 12234453.
148. Kurwale NS, Ahmad F, Suri A, Kale SS, Sharma BS, Mahapatra AK, Suri V, Sharma MC. Postoperative pituitary apoplexy: preoperative considerations toward preventing nightmare. *Br J Neurosurg.* 2012;26(1):59–63.
149. Hetelekidis S, Barnes PD, Tao ML, Fischer EG, Schneider L, Scott RM, Tarbell NJ. 20-year experience in childhood craniopharyngioma. *Int J Radiat Oncol Biol Phys.* 1993;27(2):189–95.
150. Habrand JL, Ganry O, Couanet D, Rouxel V, Levy-Piedbois C, PierreKahn A, Kalifa C. The role of radiation therapy in the management of craniopharyngioma: a 25-year experience and review of the literature. *Int J Radiat Oncol Biol Phys.* 1999;44(2):255–63.
151. Minniti G, Esposito V, Amichetti M, Enrici RM. The role of fractionated radiotherapy and radiosurgery in the management of patients with craniopharyngioma. *Neurosurg Rev.* 2009;32(2):125–32; discussion 132.
152. Savateev AN, Trunin YY, Mazerkina NA. Radiotherapy and radiosurgery in treatment of craniopharyngiomas. *Zh Vopr Neirokhir Im N N Burdenko.* 2017;81(3):94–106.
153. Ramirez-Zamora A, Giordano JJ, Gunduz A, Brown P, Sanchez JC, et al. Evolving applications, technological challenges and future opportunities in neuromodulation: proceedings of the fifth annual deep brain stimulation think tank. *Front Neurosci.* 2018;11:734.
154. Lee M, Korner J. Review of physiology, clinical manifestations, and management of hypothalamic obesity in humans. *Pituitary.* 2008;12(2):87–95.
155. Perlemonne C, Macia F, Tison F, et al. Effects of subthalamic nucleus deep brain stimulation and levodopa on energy production rate and substrate oxidation in Parkinson's disease. *Br J Nutr.* 2005;93:191–8.
156. Macia F, Perlemonne C, Coman I, et al. Parkinson's disease patients with bilateral subthalamic deep brain stimulation gain weight. *Mov Disord.* 2004;19:206–12.
157. Montaurier C, Morio B, Bannier S, et al. Mechanisms of body weight gain in patients with Parkinson's disease after subthalamic stimulation. *Brain.* 2007;130:1808.
158. Calleja-Castillo JM, Cruz-Aguilera DL, Manjarrez J, Velasco-Velázquez MA, Morales-Espinoza G, Moreno-Aguilar J, Hernández ME, Aguirre-Cruz L, Pavón L. Chronic deep brain stimulation of the hypothalamic nucleus in wistar rats alters circulatory levels of corticosterone and proinflammatory cytokines. *Clin Dev Immunol.* 2013;2013:698634.
159. Berton AM, Gatti F, Penner F, Varaldo E, Prencipe N, Rumbolo F, Settanni F, et al. Early copeptin determination allows prompt diagnosis of post-neurosurgical central diabetes insipidus. *Neuroendocrinology.* 2020;110:525–34.
160. Lamas C, del Pozo C, Villabona C, Neuroendocrinology Group of the SEEN. Clinical guidelines for management of diabetes insipidus and syndrome of inappropriate antidiuretic hormone secretion after pituitary surgery. *Endocrinol Nutr.* 2014;61(4):e15–24.
161. Schreckinger M, Szerlip N, Mittal S. Diabetes insipidus following resection of pituitary tumors. *Clin Neurol Neurosurg.* 2013;115(2):121–6.
162. Chirardello F, Hopper N, Albanese A, Maghnie M. Diabetes insipidus after craniopharyngioma: postoperative management of water and electrolyte disorders. *J Pediatr Endocrinol Metab.* 2006;19(suppl 1):413–21.
163. Hensen J, Henig A, Fahlbusch R, Meyer M, Boehnert M, Buchfelder M. Prevalence, predictors and patterns of postoperative polyuria and hyponatremia in the immediate course after transsphenoidal surgery for pituitary adenomas. *Clin Endocrinol (Oxf).* 1999;50:431–9.
164. Robinson AG, Verbalis JG. Posterior pituitary. In: Kronenberg HM, Melmed S, Polonsky KS, Larsen PR, editors. *Williams textbook of endocrinology*, vol. 11. Philadelphia: Saunders Elsevier; 2008. p. 273.
165. Popugaev KA, Savin IA, Goriachev AS, Kadashev BA. Hypothalamic injury as a cause of refractory hypotension after sellar region tumor surgery. *Neurocrit Care.* 2008;8(3):366–73.
166. Bealer SL. Vascular capacitance following preoptic recess lesions. *Am J Physiol Heart Circ Physiol.* 1993;264:H560–6.
167. Brody MJ, Johnson AK. Role of the anteroventral third ventricle region in fluid and electrolyte bal-

- ance, arterial pressure regulation, and hypertension. In: Martini L, Ganong WF, editors. *Frontiers, neuroendocrinology*. New York: Raven; 1980. p. 249–92.
168. Lind RW. Angiotensin and the lamina terminalis: illustrations of a complex unity. *Clin Exp Hypertens A*. 1988;10:79–105.
169. Smith PM, Ferguson AV. Vasopressin acts in the subfornical organ to decrease blood pressure. *Neuroendocrinology*. 1997;66:130–5.
170. Lew SM. Temporary central hypoventilation following craniopharyngioma resection. *J Pediatr Neurol*. 2004;2(3):167–70.
171. Harper RM, Gozal D, Bandler R, Spriggs D, Lee J, Alger J. Regional brain activation in humans during respiratory and blood pressure challenges. *Clin Exp Pharmacol Physiol*. 1998;25:483–6.
172. Hrish AP, Lionel KR. Intraoperative hyperthermia: a harbinger of hypothalamic injury. *J Neurosurg Anesthesiol*. 2017;29(3):379–80.

Index

A

Acquired central hypoventilation, 511, 512
Acquired immunodeficiency syndrome (AIDS), 429
Acquired pituitary deficiency, 216–217
ACTH-dependent Cushing's syndrome, 185
ACTH-independent Cushing's syndrome, 185
Adamantinomatous, 354
Adenohypophysis, 125
Adipocytokines, 236
Adiposogenital dystrophy, 245
Adrenocorticotrophic hormone (ACTH), 13, 95–97, 158, 216
Agouti gene-related peptide (AGRP), 237
Alpha-methyl dopa acts, 498
Alstrom's syndrome, 254–256
Alzheimer's disease (AD), 150, 484–486
Amphetamines, 501
Amylin, 239
Amyotrophic lateral sclerosis (ALS), 151, 488, 489
Aneurysms, 135
Angelman syndrome, 262, 324
Anti-caspr2 (contactin-associated protein-like 2), 449
Antidepressants, 498, 500
Antidiuretic hormone (ADH) deficiency, 99, 225–227, 499, 510
Anti-DPPX (dipeptidyl-peptidase-like protein-6), 449
Antiemetics/prokinetics, 498
Antihypertensives, 498
Anti-IgLON 5 disease
 haplotypes, 452, 454
 hypocretin-1 levels, 455
 IgG4, 454
 MRI, 451
 neuronal development, 449
 pathology, 450, 454
 sleep disorder, 449–452
 treatment, 452
 types of syndromes, 451
Anti-LGI1 (Leucine-rich, glioma-inactivated 1)
 encephalitis, 448, 449
Antineoplastic agents, 500
Antipsychotics, 497, 498, 500
Aquaporin-4 (AQP4)
 astrocytes, 440

 hormonal deficiencies, 442
 hypothalamic lesions, 441
 secondary narcolepsy, 442
 SIADH, 441, 442

Arachnoid cysts, 128, 388, 389
Arginine vasopressin (AVP), 44, 159, 203, 510
Autoimmune hypothalamitis, 396, 397
Autonomic control, 9–11
Autonomic nervous system (ANS), 238
Autosomal dominant NDI, 323
Autosomal recessive NDI (ARNDI), 323
AVP receptors, 47

B

Babinski-Froehlich's syndrome, 245
Bardet-Biedl syndrome, 254, 256, 301, 327
Beckwith-Wiedemann syndrome (BWS), 255
Bipolar disorder, 160, 161
Borrelia burgdorferi, 425
Brain edema, 480, 481
Brown adipose tissue (BAT), 238, 241

C

Calcitonin gene-related peptide (CGRP), 238
Carbamazepine, 500
Cataplexy, 459, 461, 462
Cavernomas
 clinical course, 391
 earlier intervention, 392
 high-flow state of, 391
 imaging, 391
 incidence, 390
 low-flow state of, 391
 malformations, 391
 sporadic cases, 391
 treatment, 392
 visual symptoms, 392
Cavernous malformations (CM), 174
Cavernous sinus thrombosis, 135
Central diabetes insipidus (CDI), 225, 425, 438, 510, 511
Central hypothyroidism, 325
Central precocious puberty (CPP), 319–320

Cerebral ischemia, 481, 482
 Childhood narcolepsy, 463
 Chlorpropamide, 500
 Cholinomimetic medications, 499
 Chordoid glioma, 368
 Choristoma, 371
 Circadian timing, 12
 Classic kallmann syndrome, 314
 CNS germ cell tumors (GCTs), 355
 Colloid cysts, 390
 Computed tomography (CT), 127
 Contiguous gene syndrome, 251
 Continuous positive airway pressure (CPAP), 324
 Corticotrophin-releasing factor (CRH), 345
 Corticotrophin-releasing hormone (CRH) gene, 310
 Corticotrophin-releasing factor (CRF), 166
 Corticotrophin-releasing hormone (CRH), 486
 Corticotrophin-releasing hormone (CRH) production, 217
 Cortisol hypersensitivity, 429
 Craniopharyngioma, 21, 24, 132, 369
 Cushing's disease, 127
 Cyclophosphamide, 500
 Cystic lesions
 arachnoid cysts, 388, 389
 colloid cysts, 390
 epidermoid and dermoid cysts, 387, 388
 RCCs, 389, 390
 Cytoplasmic glucocorticoid receptor (GCCR), 312

D

Deep brain stimulation (DBS), 510
 Delta-opioid receptor (DOR), 165
 Dementia, 486
 Dementia with lewy bodies (DLB), 487
 Demyelinating disorders, 142, 143
 Dermoid cysts, 21, 352, 387, 388
 Desmopressin (DDAVP), 189
 Dexamethasone-CRH (DEX-CRH) test, 444
 Diabetes insipidus (DI), 322
 Diabetes mellitus (DM), 326
 Diencephalic syndrome, 263
 Disrupted nighttime sleep, 462
 Diuretics, 500
 Dopamine, 243
 Dorsal medial hypothalamus (DMH), 41
 Dorsal vagal nuclear complex (DVNC), 239
 Drug induced hyperthermia, 500, 501
 Dynorphin (DYN), 165

E

Ectopic posterior lobe, 128
 Ectopic posterior pituitary, 399
 Encephalitis, 397
 Endoscopic endonasal approach, 177
 Enterocytokines, 236
 Epidermoid cysts, 17, 352, 387, 388
 Epidermoid tumors, 128
 Estradiol treatment, 501

Excessive daytime sleepiness (EDS), 459, 460
 Exogenous Cushing's syndrome, 185
 Extracellular fluid (ECF), 100

F

Fibroblast growth factor receptor 1 (FGFR1), 314
 Fluid-attenuated inversion recovery (FLAIR), 15
 Follicle-stimulating hormone (FSH), 130, 222
 Fragile X syndrome (FXS), 256, 257
 Froelich's syndrome, 245
 Frontotemporal dementia (FTD), 151
 Frontotemporal lobar degeneration (FTLD), 489

G

GABA receptors, 161
 Gangliocytomas, 348
 Ganglioglioma, 349–350
 Genetic syndromes
 BBS
 clinical features and diagnosis, 328
 pathogenesis, 327–328
 clinical disorders, 296
 familial neurohypophyseal diabetes insipidus
 ADNDI, 323
 ARNDI, 323
 genetic forms of, 323
 physiology of arginine vasopressin secretion, 323
 x-linked NDI, 323
 genes and genetic mutations, 302, 306
 genetics of hypothalamic programming, 329–331
 HHS
 clinical features of, 321
 CPP and surgery/antiepileptic drugs, 320
 diagnosis, 322
 epidemiology, 320
 genetics, 320
 pathophysiology, 320–321
 treatment, 322
 hypothalamic development anomalies, 295
 hypothalamic nutrition/appetite/weight control, 318
 leptin/leptin receptor syndromes, 328, 329
 midline anomalies, 298–299
 neurodegenerative dysfunction, 304
 neuroendocrine puberty and sexual maturation, 310, 311
 neuroendocrine thyroid, 309
 PHS
 clinical features, 325–326
 diagnosis, 326
 management, 326
 PWS
 adrenal insufficiency, 325
 central hypothyroidism, 325
 clinical features, 324
 genetics, 324
 growth hormone deficiency, 324–325
 hypogonadism, 325
 sleep disorders, 324

- wolfram syndrome
 - clinical features and management, 326–327
 - diagnosis, 327
 - genetics, 327
 - Genome-wide association scans (GWAS), 247
 - Germ cell tumors (GCT), 133, 352
 - Germinomas, 24, 376, 377
 - GH deficiency (GHD), 221
 - Ghrelin, 240
 - Glial fibrillary acidic proteins (GFAP), 368
 - Gliomas, 350–352, 367
 - Gonadotropin-releasing hormones (GnRH), 58, 222, 321, 345
 - Granulomatous disease, 28
 - Granulomatous hypophysitis, 439
 - Granulomatous infundibulohypophysitis, 396
 - Granulovacuolar degeneration (GVD), 487
 - Growth hormone (GH) deficiency, 324–325, 347
 - Growth-hormone-releasing hormone (GHRH), 12, 94, 312, 345
- H**
- Hallucinations, 462
 - Hand-Schüller-Christian disease, 357
 - Hantaviruses, 426
 - Hartsfield syndrome, 303
 - Hemangiopericytoma, 379
 - Hemorrhagic fever with renal syndrome (HFRS), 426
 - Hemorrhagic stroke, 480, 481
 - Histamine, 283
 - Histoplasmosis capsulatum*, 174, 175
 - H1N1 vaccination, 463
 - Homeostasis, 33, 35, 36
 - Hormone deficiency syndromes
 - ACTH deficiency
 - clinical manifestations, 217
 - diagnosis, 218
 - etiology, 217
 - physiology, 216
 - treatment of, 218, 219
 - ADH, 225–227
 - FSH and LH
 - clinical manifestations, 223
 - diagnosis, 223, 224
 - etiology, 222, 223
 - management, 224
 - physiology, 222
 - GHD
 - clinical manifestations, 221
 - diagnosis, 221
 - etiology, 221
 - treatment, 222
 - hypopituitarism, 215
 - prolactin deficiency, 225
 - TSH deficiency
 - clinical manifestations, 220
 - diagnosis of, 219, 220
 - etiology, 219
 - physiology, 219
 - treatment, 220
 - Hormone excess syndromes
 - acromegaly
 - clinical manifestations, 193–194
 - diagnosis, 195
 - differential diagnosis, 195
 - management, 195–198
 - medical management of, 197
 - recommendations for screening, 197
 - Cushing's syndrome
 - causes of, 185
 - clinical manifestations, 185–187
 - diagnosis of, 187–189
 - differential diagnosis, 190
 - management, 191–193
 - postoperative management, 191–193
 - pseudo-Cushing's syndrome, 189, 190
 - recurrence of, 191–193
 - remission, 191
 - gonadotroph adenomas
 - clinical manifestations of, 200–202
 - diagnosis, 200–202
 - differential diagnosis, 202
 - management, 203
 - inappropriate antidiuretic hormone
 - clinical manifestations, 203–204
 - diagnosis, 204
 - differential diagnosis, 204–205
 - management, 205–206
 - prolactinomas
 - clinical presentation of, 181, 182
 - diagnosis, 182, 183
 - differential diagnosis, 183–184
 - management, 184–185
 - TSHomas
 - clinical manifestations, 198
 - diagnosis, 198–199
 - differential diagnosis, 199–200
 - management, 200
 - H2-receptor antagonists, 498
 - Human African trypanosomiasis (HAT), 426–428
 - Huntington's disease, 305, 488
 - HWS, hyperphagia, 235
 - Hypothalamic disease
 - amenorrhea, 178
 - anatomy of, 171
 - Cushing syndrome, 178
 - deep subcortical AVMs, 173, 174
 - diabetes insipidus, 178
 - endoscopic endonasal approach, 177
 - expansive craniopharyngioma, 172–173
 - growth abnormalities, 178
 - histoplasmosis capsulatum*, 174, 175
 - hypothalamic hamartoma, 171–172
 - intra-cranial surgery, 177
 - non-operative considerations, 178
 - numerous bodily functions, 177
 - optic pathway and hypothalamic gliomas, 173
 - pilocytic astrocytomas, 172
 - pterional approach, 175

- Hypothalamic disease (*cont.*)
 sexual dysfunction, 178
 subfrontal translaminal terminalis approach, 176
 supraorbital craniotomy, 175–176
 supratentorial CMs, 174
 thyroid abnormalities, 178
 transcallosal approach, 176
 trans-sylvian approach, 175
 transventricular approach, 175
 visual pathway, 177
- Hypercortisolemia, 429
- Hyperprolactinemia
 antidepressants, 498
 antiemetics/prokinetics, 498
 antihypertensives, 498
 antipsychotics, 497, 498
 causes of, 183
 cholinomimetic medications, 499
 H2-receptor antagonists, 498
 opiates, 498
 prolactin, 497
- Hypersomnias, *see* Narcolepsy Type-1
- Hypocretin orexin system, 462, 463
- Hypocretins (Hcrts), 160
- Hypogonadism, 325
- Hypophysitis
 clinical features, 437
 CTLA-4 blockers, 440
 granulomatous hypophysitis, 439
 IgG4, 440
 lymphocytic hypophysitis, 437–439
 necrotizing hypophysitis, 439
 treatment, 440
 xanthomatous hypophysitis, 439
- Hypopituitarism, 215, 499
 immune checkpoint inhibitors, 499
- Hypothalamic craniopharyngioma, 369
- Hypothalamic dysfunction, 483, 486
 anterior pituitary hormone control, 297
 appetite control, 296
 central autonomic transmission control, 297
 cranial facial malformations, 294
 emotions, behavior, and memory control, 297
 genetic syndromes of (*see* Genetic syndromes)
 hypothalamic behavior and memory modulation, 306–308
 hypothalamic development/midline anomalies, 297–301
 hypothalamic neurodevelopment, 303–306
 hypothalamic neurogenesis, 295
 hypothalamic tumors, 301–303
 hypothalamic water/metabolic balance control, 316–317
 neuroendocrine secretion control, 313
 neurodegenerative dysfunction, 303–306
 neuroendocrine secretion control, 308–316
 nutrition/appetite/body weight control, 317–319
 plasma water electrolyte (sodium)-osmotic balance, 296
 sleep/circadian rhythm control, 297
 thermoregulation, 296
 Wnt and SHH signaling pathways, 294
- Hypothalamic function, 33
- Hypothalamic glioma, 131
- Hypothalamic hamartoma, 172, 392, 393
 clinical course, 392
 gelastic seizures, 393
 histopathologic analysis, 393
 imaging, 393
 incidence, 392
 morphologic classification, 393
 treatment, 393
- Hypothalamic hamartoma syndrome (HHS), 319–322
- Hypothalamic hamartomas (HH), 17, 129, 302, 356, 357
- Hypothalamic obesity syndrome (HOS), 235, 511
 Alstrom's syndrome, 255
 Bardet-Biedl syndrome, 254
 CART gene mutations, 249
 causes of, 244–247
 clinical presentation of, 258
 FXS, 256, 257
 genetic syndromes of, 247
 leptin gene mutations, 248
 leptin receptor mutations, 248–249
 microdeletion syndrome, 256
 monogenic, 248–251
 polygenic HOS, 257–258
 POMC mutations, 249–250
 prohormone convertase 1 gene mutations, 250
 PWLS, 254
 PWS, 251–254
 syndromic HOS, 251
 tub gene mutations, 250
 WAGR syndrome, 255
- Hypothalamic pathways assisting regulation, 320
- Hypothalamic pituitary prolactin (HPP), 73
- Hypothalamic tumors, 301–303
- Hypothalamic wasting syndrome (HWS), 235
 anatomic location/topography, 260
 causes of, 261–263
 clinical phenotype, 261
 diencephalic syndrome, 263
 hypothalamic dysfunction, 259
 melanocortin central signaling pathways, 260
 Simmonds' cachexia, 262, 263
- Hypothalamic-chiasmatic gliomas, 24
- Hypothalamic-pituitary-adrenal (HPA), 144, 159, 215
- Hypothalamic-pituitary axis (HPA), 482
- Hypothalamic-pituitary gonadal axis, 38, 488
- Hypothalamic-pituitary-gonadal axis-reproductive age, 39
- Hypothalamitis, 396, 397
- Hypothalamo-pituitary axis
 AIDS, 429
 fungal infections, 429
 hematogenous dissemination, 415
 HIV infection, 429
 neuroborreliosis, 425
 non-mycobacterial meningitis, 421
 parasitic infections
 HAT, 426–428

- toxoplasmosis, 426
 - physical and endocrine dysfunction, 415
 - pituitary abscesses
 - clinical outcome, 424
 - coronal and sagittal MRI, 423
 - hematogenous dissemination, 422
 - isolated hypogonadism, 422
 - neurological signs and symptoms, 422
 - organisms, 424
 - treatment, 424
 - risk factors, 415
 - treponemas pallidum infection, 421, 422
 - tuberculosis
 - central nervous system, 416, 417
 - epidemiology, 416
 - tuberculomas, 417–420
 - viral infections
 - CDI, 425
 - hantaviruses, 426
 - meningitis and encephalitis, 425
 - von Economo's encephalitis lethargica, 426
 - Hypothalamo-pituitary-adrenal (HPA), 13
 - Hypothalamus, 15, 79, 141–143, 151, 152, 157–161, 163–166, 236, 294
 - anatomy, MR imaging, 16
 - autonomic control, 9, 10
 - AVP functions, 46, 47
 - AVP peptide, 44
 - AVP plasma levels, 46
 - AVP receptors, 47
 - AVP secretion, 45
 - AVP synthesis, 45
 - blood pressure, 47, 48
 - blood pressure regulation, 49
 - blood supply, 8, 9
 - body temperature, 40
 - body weight, 13
 - body's homeostasis, 3
 - brainstem reticular activating system, 283
 - circadian timing, 12
 - craniopharyngiomas, 21, 24
 - Dermoid cysts, 21
 - disk-over-water paradigm, 288
 - endocrine control, 10, 11
 - epidermoid cysts, 17
 - expansive intrahypothalamic connectivity, 281
 - extrahypothalamic influences, 281
 - extra-hypothalamic projections, 36
 - extrahypothalamic systems, 288
 - feeding circuitry, 33, 34
 - food intake, 4, 33
 - food intake, hypothalamic regulation, 34
 - GABA and glutamate, 48
 - GABA neurons, 283
 - GABA/galanin neurons, 286
 - gastrointestinal hormones, 36
 - germinomas, 24
 - GnRH neurons, 37
 - granulomatous disease, 28
 - gross anatomy, 7, 8
 - HPG, 37
 - hypothalamic and basal forebrain systems, 282
 - hypothalamic sleep disorders, 288
 - hypothalamic topography, 9
 - inferior portion, 3
 - insulin, 36
 - KNDy neurons, 38
 - langerhans cell histiocytosis, 28
 - leptin, 36
 - lipoma, 17
 - lymphocytic hypophysitis, 28
 - memorization, 13, 14
 - metastatic tumors, 28
 - MR imaging, 15
 - neuronal activity, 49
 - neuropeptide Y/agouti-related peptide, 35
 - nitric oxide, 49
 - orexin/hypocretin neurons, 285
 - orexin/hypocretin system, 288
 - osmoreceptors, 43, 44
 - POA, 40
 - proopiomelanocortin, 35
 - pubertal maturation, 38, 39
 - punishment, 6
 - rathke cleft cysts, 21
 - reproduction, hypothalamic control, 37
 - reward, 6
 - sarcoidosis, 28, 29
 - SCN efferents, 284
 - SCN neurons, 284
 - SCN rhythms, 284
 - selective neuron populations, 283
 - sleep–wake cycle, 5
 - surrounding structures, 29
 - synaptic output, 40, 41
 - temperature, 41
 - thermosensitive neurons, 40
 - tuberomammillary hypothalamic neurons, 285
 - tuberomammillary nucleus, 282
 - vasopressin, 46
 - water balance, 5
 - water/fluid balance, 41
 - Hypothalamus-mediated aggressive behavior, 486
- ## I
- ### Iatrogenic hypothalamic disorders
- neurosurgical approaches
 - acquired central hypoventilation, 511, 512
 - anatomy, 501, 502
 - CDI, 510, 511
 - complications, 501
 - DBS surgery, 510
 - history, 502, 503
 - HOS, 511
 - intraoperative hyperthermia, 512
 - mechanisms of hypothalamic injury, 509, 510
 - preoperative considerations, 508–509
 - radiation therapy, 510
 - sympathetic dysfunction, 511

neurosurgical approaches (*cont.*)
 transcranial surgery, 503, 504
 transsphenoidal endoscopic surgery, 505–508
 transsphenoidal microscopic surgery, 504, 505
 transsphenoidal surgery, 504
 non-surgical
 amphetamines, 501
 drug induced hyperthermia, 500, 501
 estradiol treatment, 501
 hyperprolactinemia (*see* Hyperprolactinemia)
 hypopituitarism, 499
 impulse control disorders, 500
 medications, 497
 SIADH, 499, 500
 Idiopathic hypogonadotropic hypogonadism (iHH), 313
 Immune checkpoint inhibitors, 499
 Impulse control disorders, 500
 Infectious lesions
 abscesses, 394
 syphilis, 395
 toxoplasmosis, 394
 tuberculosis, 394, 395
 Inflammatory conditions of hypothalamus
 anti-IgLON 5 disease
 haplotypes, 452, 454
 hypocretin-1 levels, 455
 IgG4, 454
 MRI, 451
 neuronal development, 449
 pathology, 450, 453, 454
 sleep disorder, 449–452
 treatment, 452
 types of syndromes, 450–451
 anti-Ma antibodies, 446–448
 anti-MOG antibodies hypothalamic lesions, 442, 443
 AQP4 and neuromyelitis spectrum disorder
 astrocytes, 440
 hormonal deficiencies, 442
 hypothalamic lesions, 441
 secondary narcolepsy, 442
 SIADH, 441, 442
 hypersomnias (*see* Narcolepsy)
 hypophysitis
 clinical features, 437
 CTLA-4 blockers, 440
 granulomatous hypophysitis, 439
 IgG4, 440
 lymphocytic hypophysitis, 437–439
 necrotizing hypophysitis, 439
 treatment, 440
 xanthomatous hypophysitis, 439
 multiple sclerosis
 fatigue and hypothalamus in people, 445
 hypothalamic lesions, 443
 hypothalamic neurodegeneration, 443, 444
 hypothalamic-pituitary axis overactivity, 444
 thermoregulation, 445
 sarcoidosis (*see* Neurosarcoidosis)
 VGKC-complex
 anti-caspr2 encephalitis, 449

anti-DPPX encephalitis, 449
 anti-LGI1 encephalitis, 448, 449
 autoimmune encephalitis, 448
 identification, 448
 Inflammatory lesions
 ectopic posterior pituitary, 399
 encephalitis and hypothalamitis, 396, 397
 lymphocytic and granulomatous
 infundibuloneurohypophysitis, 396
 multiple sclerosis, 398, 399
 sarcoidosis, 397, 398
 Wegner's granulomatosis/granulomatosis with
 polyangiitis, 395, 396
 Inflammatory lesions Wegner's granulomatosis/
 granulomatosis with polyangiitis, 396
 Infundibular stalk, 16, 24, 28
 Infundibuloneurohypophysitis (INH), 438, 439
 Inherited pituitary deficiency, 216
 International classification of sleep disorders (ICSD-3),
 463
 Intracranial lipomas, 378
 Ischemic stroke, 480, 481
 Isolated adrenocorticotrophic hormone (ACTH)
 deficiency, 325

K

Kleine-Levin syndrome, 153, 154

L

Langerhans cell histiocytosis (LCH), 357, 358, 374, 375
 Launois-Cleret syndrome, 245
 Leptin, 237
 Leptin/leptin receptor syndromes, 328–329
 Leptomeningeal disease, 397
 Letterer-Siwe disease, 357
 Leukemia, 375–376
 Levothyroxine, 220
 Limbic encephalitis, 143, 144
 Longitudinal extensive myelopathy (LEM), 398
 Luteinizing hormone (LH), 222
 Luteinizing hormone-releasing hormone (LHRH), 61, 345
 Lyme disease, 425
 Lymphocytic hypophysitis, 28
 Lymphocytic infundibulohypophysitis, 396
 Lymphocytic infundibuloneurohypophysitis (LINH),
 437–439
 Lymphocytic pan-hypophysitis (LPH), 437

M

Macroadenomas, 130
 Macroglia, 483
 Major depressive disorder (MDD), 159
 MC4R heterozygous mutations, 319
 McKusick-Kaufman syndrome, 254
 Melancholic, 160–162
 Melanin-concentrating hormone (MCH), 5
 Melarsoprol, 428

- Meningiomas, 132, 372, 373
- Metastatic disease, 377
- Micro RNAs (miRNAs), 294
- Microdeletion syndrome, 256
- Microglia, 483
- Migraine, 146, 147
- Morvan syndrome, 449
- Müller classification, 508
- Multi-organ CHARGE syndrome, 315
- Multiple sclerosis (MS), 398, 399
 - fatigue and hypothalamus in people, 445
 - hypothalamic lesions, 443
 - hypothalamic neurodegeneration, 443, 444
 - hypothalamic-pituitary axis overactivity, 444
 - thermoregulation, 445
- Multiple sleep latency test (MSLT), 152, 463
- Multi-system atrophy (MSA), 487
- Mycobacterium tuberculosis
 - central nervous system, 416, 417
 - epidemiology, 416
 - tuberculomas, 417–420
- Myocutaneous flap, 503

- N**
- N- arachidonoyl dopamine (NADA), 241
- Narcolepsy, 152
- Narcolepsy type I
 - autoimmunity
 - H1N1 vaccination and childhood narcolepsy, 463
 - HLA-haplotype, 463
 - immunotherapy, 464, 465
 - pathogenic antibodies, 463, 464
 - symptomatic treatment, 465
 - T cells, 464
 - cataplexy, 459, 461, 462
 - diagnosis of, 463
 - disrupted nighttime sleep, 462
 - EDS, 459, 460
 - hallucination, 462
 - and hypocretin orexin system, 462, 463
 - sleep attacks, 459
 - sleep paralysis, 462
- Nelson's syndrome, 206
- Neuroborreliosis, 425
- Neurocytomas, 349
- Neurodegenerative disease
 - components, 483
 - dementia, 483
 - functional connectivity, 484
 - Huntington's disease, 488
 - microglia and macroglia, 483
 - Parkinson's disease, 487, 488
 - stress, 486, 487
- Neurodegenerative disorders, 150
- Neuroendocrine neoplasms and lesions
 - angiotensin II, 345
 - cholecystokinin, 345
 - clinical presentation, 346
 - dopamine, 345
 - gangliocytomas, 348
 - ganglioglioma, 349–350
 - glial neoplasms
 - gliomas, 350–352
 - pituicytomas, 350
 - histamine, 345
 - hypothalamic hormone production, 345
 - hypothalamic hormones, 345
 - infiltrating neoplasms
 - CNS GCTs, 355
 - craniopharyngiomas, 353–354
 - dermoid and epidermoid cysts, 355, 356
 - pituitary adenomas, 352, 353
 - Rathke's cleft cysts, 356
 - neurocytomas, 349
 - neuroendocrine disturbances, 347–348
 - neurological symptoms, 346
 - neurotensin, 345
 - nonneoplastic lesions
 - hypothalamic hamartomas, 356, 357
 - LCH, 357, 358
 - neurosarcoidosis, 358
 - sellar region neoplasms, 346
 - serotonin, 345
 - substance P, 345
 - visual disturbances, 347
- Neuroendocrinology
 - adjunctive non endocrine tissues, 53
 - anterior pituitary hormone, physiology, 92, 93
 - APr, 64
 - behavioral responses, 89
 - circadian rhythms, 86, 87, 89
 - circumventricular organs, 62
 - food intake, 79
 - growth hormone, 93, 94
 - HPA axis, 75
 - HPG axis, 76, 77, 79
 - HPGhI axis, 74
 - HPT axis, 75, 76
 - hypophysis cerebri, 102, 103
 - hypothalamic axis, 54–56, 58
 - hypothalamic functional connections, 68, 69
 - hypothalamic neuronal connections, 60–62, 69, 70
 - hypothalamus, 70–72
 - median eminence, 63
 - neurosecretory neurons, 53
 - osmotic regulation, 100, 101
 - OVLT, 63
 - pineal gland, 65–68
 - pituitary axis
 - anatomy, 90, 91
 - blood supply, 90
 - posterior pituitary, 98, 99
 - prominent hypotheses, 81–83
 - scientific discipline, 53
 - SCO, 64
 - SFO, 63, 64
 - sleep-wake cycle, 87, 89
 - tela choroidea, 64
 - thermoregulation, 84, 85

- Neuroendocrinology (*cont.*)
 thyroid-stimulating hormone, 97, 98
 unique endocrine glands, 53
- Neurohypophysis, 42
- Neuroleptic malignant syndrome (NMS), 501
- Neuromyelitis optica (NMO), 398
- Neuromyelitis optica spectrum disorder (NMOSD), 143
- Neuromyelitis spectrum disorder
 astrocytes, 440
 hormonal deficiencies, 442
 hypothalamic lesions, 441
 secondary narcolepsy, 442
 SIADH, 441, 442
- Neuromyotonia, 449
- Neuropeptide-Y (NPY), 511
- Neurosarcoid, 397
- Neurosarcoidosis, 358
 clinical manifestations, 456
 diagnosis of, 458
 epidemiology, 455, 456
 hypothalamic involvement
 endocrine manifestations, 457
 endocrinopathies, 456
 homeostatic functions, 457
 obesity, 457
 water balance impairment, 456, 457
 noncaseating granuloma, 455
 treatment of, 458, 459
- Neuro-structural syndromes, 141
- Nicotinamide adenine dinucleotide (NADH), 317
- Nitric oxide (NO), 49
- N-methyl-D-aspartate (NMDA), 145
- Noladin ether, virodhamine, 241
- Non-endocrine neoplasms
 extrinsic tumors
 germinomas, 376, 377
 lipomas and osteolipomas, 378
 metastatic disease, 377
 hematologic malignancies langerhans cell
 histiocytosis, 374, 375
 hematologic malignancies leukemia, 375–376
 intrinsic CNS lesions
 chordoid glioma, 368
 choristoma, 371
 choroid plexus papilloma/carcinoma, 374
 craniopharyngiomas, 369
 gliomas, 367
 meningiomas, 372, 373
 PCNSL, 370
 pilocytic astrocytoma, 368, 369
 pituicytomas, 370, 371
 solitary fibrous tumors, 373, 374
 vascular tumors
 hemangioblastoma, 378–379
 hemangiopericytoma, 379
- Nonfunctioning pituitary adenomas (NFPAs), 200
- Non-germinomatous germ cell tumors (NGGCT), 355
- Non-hodgkin large B-cell lymphoma, 370
- Non-mycobacterial meningitis, 421
- Nonneoplastic mass lesions
 cystic lesions
 arachnoid cysts, 388, 389
 colloid cysts, 390
 epidermoid and dermoid cysts, 387, 388
 RCCs, 389, 390
 infectious lesions
 abscesses, 394
 syphilis, 395
 toxoplasmosis, 394
 tuberculosis, 394, 395
 inflammatory lesions
 ectopic posterior pituitary, 399
 encephalitis and hypothalamitis, 396, 397
 lymphocytic and granulomatous
 infundibuloneurohypophysitis, 396
 multiple sclerosis, 398, 399
 sarcoidosis, 397, 398
 Wegner's granulomatosis/granulomatosis with
 polyangiitis, 395, 396
 vascular lesions
 cavernomas, 390–392
 hypothalamic hamartoma, 392, 393
- NOTCH signaling pathway, 295
- Nucleus accumbens (NA), 238
- Nutrient sensing, 243
- O**
- Obestatin, 240
- Online mendelian inheritance in man (OMIM), 297
- Opiates, 498, 500
- Orexin (OX), 165
- Osmoreceptors, 42, 43
- Oxyntomodulin, 239
- Oxytocin, 101, 102, 316
- P**
- Pallister-Hall syndrome (PHS), 325–326, 357
- Pancreatic polypeptide (PP), 240
- Paraneoplastic limbic encephalitis, 397
- Paraventricular nucleus (PVN), 3, 4, 47, 48, 236
- Parkinson's disease (PD), 306, 487, 488
- Pentamidine, 428
- Pilocytic astrocytomas, 172, 368, 369
- Pilomyxoid astrocytoma (PMA), 351, 369
- Pituicytomas, 350, 370, 371
- Pituitary abscesses, 135
 clinical outcome, 424
 coronal and sagittal MRI, 423
 hematogenous dissemination, 422
 isolated hypogonadism, 422
 neurological signs and symptoms, 422
 organisms, 424
 treatment, 424
- Pituitary adenomas, 129
- Pituitary gland, 91
- Plasmapheresis, 465
- Posterior pituitary hormones, 99
- Postpartum depression (PPD), 160

Prada-Willi-like syndromes (PWLS), 254
 Prader-Willi syndrome (PWS), 251–254, 304, 323–325, 411
 Primary central nervous system lymphoma (PCNSL), 370
 Profound anorexia, 235
 Prolactin, 497
 Prolactin deficiency, 225
 Proopiomelanocortin (POMC), 35, 36, 166, 511
 Pseudo-Cushing's syndrome, 189, 190
 Pterional approach, 175

R

Rapid-onset obesity with hypothalamic dysfunction, hypoventilation, and auto-nomic dysregulation (ROHHAD) syndromes
 clinical manifestations, 408
 pathogenesis, 406
 Rathke's cleft cysts (RCCs), 21, 127, 128, 352, 356, 389, 390
 Rathke's pouch (RP), 172, 294, 354, 369
 Renin-angiotensin-aldosterone system (RAAS), 100, 217
 Rhodesiense form, 427
 Russell-Silver syndromes, 262, 263

S

Sarcoidosis, 28, 135, 358, 397, 398
 Schaaf-Yang syndrome, 304
 Schizophrenia, 161, 162
 Schwannoma/neurinomas, 133
 Sellar region, 125–127
 Serotonin (5HT), 242
 Simmonds' cachexia, 262, 263
 Simmonds' disease, 262
 Sleep paralysis, 462
 Solitary fibrous tumors, 373, 374
 Somatostatinoma, 302
 Sonic hedgehog (SHH) signaling pathways, 294
 Sporadic single nucleotide polymorphisms (SNPs), 293
 Stress, 486, 487
 Strokes, 425
 Subarachnoid hemorrhage (SAH), 480, 481
 Subfrontal translaminal terminalis approach, 176
 Sulphonylureas, 500
 Sundowning effect, 486
 Sympathetic dysfunction, 511
 Sympathomimetic syndrome, 501
 Syndrome of inappropriate antidiuretic hormone secretion (SIADH), 349, 499, 500
 antidepressants, 500
 antineoplastic agents, 500
 antipsychotics, 500
 carbamazepine and oxcarbazepine, 500
 clinical manifestation, 500
 diuretics, 500
 opiates, 500
 sulphonylureas, 500
 Syphilis, 395, 421, 422

T

Takeda G protein-coupled receptor-5 (TGR5), 241
 Testosterone therapy, 224
 Thermosensitive neurons, 40
 Thiazide diuretics, 500
 Thyroid hormone, 241
 Thyroid stimulating hormone (TSH) deficiency, 97, 98, 219, 347
 Thyroid transcription factor-1 (TTF-1), 350
 Thyrotropin (TSH)-secreting pituitary adenomas (TSHomas), 198
 Thyrotropin-releasing factor (TRH), 345
 Thyrotropin-releasing hormone (TRH), 11
 Thyrotropin-releasing hormone (TRH) stimulation test, 219
 Tolosa-Hunt syndrome, 137
 Toxoplasma gondii, 426
 Toxoplasmosis, 394, 426
 Transcallosal approach, 176
 Transcranial surgery, 503, 504
 Transient receptor potential channel, vanilloid subfamily (TRPV4), 441
 Transsphenoidal endoscopic surgery, 506
 advantages and disadvantages, 505
 bimanual technique, 505
 craniopharyngiomas, 508
 CSF circulation, 507, 508
 dissection strategy, 506, 507
 dura, 506
 fascia lata graft, 508
 illumination and visualization, 505
 improvement, 508
 intraoperative navigation, 505
 limitations, 508
 sellar region, 506
 sphenoid sinus, 505–506
 Transsphenoidal microscopic surgery, 504, 505
 Trans-sphenoidal microscopic surgery, 505
 Transsphenoidal surgery, 504
 Trans-sylvian approach, 175
 Traumatic brain injury (TBI), 479, 480
 Treponemas pallidum infection, 421, 422
 Tribbles homolog 2 (Trib2), 464
 Trigeminal autonomic cephalgias (TACs), 147, 148
 Trypanosoma brucei, 426
 Tuberculomas, 417–420
 Tuberculosis, 394, 395
 Tuberculous bacilli, 417
 Type 3 diabetes, 485

U

Urine free cortisol (UFC) test, 187

V

Vascular lesions
 cavernomas, 390–392
 hypothalamic hamartoma, 392, 393
 Vasoactive intestinal peptide (VIP), 345

Vasopressin, 499
Ventral tegmental area (VTA), 83
Ventrolateral preoptic anterior (VPA), 297
Ventromedial nucleus (VMN), 235
Verapamil acts, 498
Voltage gated potassium channels (VGKCs), 145
 anti-caspr2 encephalitis, 449
 anti-DPPX encephalitis, 449
 anti-LGI1 encephalitis, 448, 449
 autoimmune encephalitis, 448
 identification, 448
Von Economo's encephalitis lethargica, 426
von Recklinghausen's disease, 263

W

WAGR syndrome, 255
Wegner's granulomatosis, 395, 396
Wernicke's encephalopathy (WE), 149
Wolfram syndrome, 326

X

Xanthomatous hypophysitis, 439
X-linked NDI, 323