

Twenty-fourth International Olympiad, 1983

- Find all functions f defined on the set of positive real numbers which take positive real values and satisfy the conditions:
(i) $f(xf(y)) = yf(x)$ for all positive x, y ;
(ii) $f(x) \rightarrow 0$ as $x \rightarrow \infty$.
- Let A be one of the two distinct points of intersection of two unequal coplanar tangents to the circles C_1 and C_2 with centers O_1 and O_2 , respectively. One of the common tangents to the circles touches C_1 at P_1 and C_2 at P_2 , while the other touches C_1 at Q_1 and C_2 at Q_2 . Let M_1 be the midpoint of P_1Q_1 , M_2 be the midpoint of P_2Q_2 prove that $\angle O_1AO_2 = \angle M_1AM_2$.
- Let a, b and c be positive integers, no two of which have a common divisor greater than 1. Show that $2abc - ab - bc - ca$ is the largest integer which cannot be expressed in the form $xbc + yca + zab$, where x, y and z are non-negative integers.
- Let ABC be an equilateral triangle and ϵ the set of all points contained in the three segments AB , BC , and CA (including A , B , and C). Determine whether for every partition of ϵ into two disjoint subsets, at least one of the two subsets that contains the vertices of a right-angled triangle. Justify your answer.
- Is it possible to choose 1983 distinct positive integers, all less than or equal to 10^5 , no three of which are consecutive terms of an arithmetic progression ? justify your answer.
- Let a, b and c be the lengths of the sides of a triangle. Prove that.

$$a^2b(a-b) + b^2c(b-c) + c^2a(c-a) \geq 0$$

Determine when equality occurs.