



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 16:50
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 01-Jan-2023 19:50
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP022	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	24.8	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 16:49
Patient Name	: anonymous	Age/Sex	: 16 Yr/M	Acceptance Date : 01-Jan-2023 19:50
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP020	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	23.2	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 18:06
Patient Name	: anonymous	Age/Sex	: 18 Yr/F	Acceptance Date : 01-Jan-2023 19:47
Sample Type	: Edta Blood	Lab No.	: 230101CB124	Reporting Date : 02-Jan-2023 07:58
Referred Doctor	:	Patient Type	: OPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood

METHOD: Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	10.2	12 - 15 g/dl
HCT	32.6	36 - 46 %
RBC	3.01	3.8 - 4.8 $\times 10^6$ / μ L
MCV	108.3	83 - 101 fL
MCH	33.9	27 - 32 pg
MCHC	31.3	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	15.8	11.6 - 14.0 %
Total Leukocyte Count(TLC)	1.50	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	12.7	40 - 80 %
Lymphocytes	82.0	20 - 40 %
Monocytes	4.0	2 - 10 %
Eosinophils	1.3	01 - 06 %
Basophil	0.0	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	0.19	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	1.23	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.06	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.02	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.00	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	111	150 - 410 $\times 10^3$ / μ L

Comments on PBS : Leukocytopenia.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:50
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 01-Jan-2023 19:53
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT034	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	9.3	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.0	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 49 Yr/F	Acceptance Date : 02-Jan-2023 02:22
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102PT008	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: OPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	10.1	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.0	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 16:50
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 01-Jan-2023 19:50
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT030	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	9.4	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.0	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	39 Yr/M
Sample Type	:	Edta Blood	Lab No.	:	230101CB133
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	01-Jan-2023 16:50
			Acceptance Date	:	01-Jan-2023 19:49
			Reporting Date	:	10-Jan-2023 16:12

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	8.0	13 - 17 g/dl
HCT	24.8	40 - 50 %
RBC	3.03	4.5 - 5.5 $\times 10^6$ / μ L
MCV	81.8	83 - 101 fL
MCH	26.4	27 - 32 pg
MCHC	32.3	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	15.0	11.6 - 14.0 %
Total Leukocyte Count(TLC)	6.77	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	59.7	40 - 80 %
Lymphocytes	29.5	20 - 40 %
Monocytes	7.5	2 - 10 %
Eosinophils	3.0	01 - 06 %
Basophil	0.3	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	4.04	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	2.00	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.51	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.20	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.02	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	121	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:21
Patient Name	: anonymous	Age/Sex	: 25 Yr/F	Acceptance Date : 01-Jan-2023 19:49
Sample Type	: Edta Blood	Lab No.	: 230101CB132	Reporting Date : 02-Jan-2023 07:58
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	10.6	12 - 15 g/dl
HCT	31.8	36 - 46 %
RBC	3.51	3.8 - 4.8 $\times 10^6$ / μ L
MCV	90.6	83 - 101 fL
MCH	30.2	27 - 32 pg
MCHC	33.3	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	17.9	11.6 - 14.0 %
Total Leukocyte Count(TLC)	5.95	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	88.4	40 - 80 %
Lymphocytes	4.5	20 - 40 %
Monocytes	6.4	2 - 10 %
Eosinophils	0.5	01 - 06 %
Basophil	0.2	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	5.26	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	0.27	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.38	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.03	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.01	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	35	150 - 410 $\times 10^3$ / μ L

Comments on PBS : Severe thrombocytopenia.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 49 Yr/F	Acceptance Date : 02-Jan-2023 02:14
Sample Type	: Edta Blood	Lab No.	: 230102CB025	Reporting Date : 10-Jan-2023 15:37
Referred Doctor	:	Patient Type	: OPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	12.5	12 - 15 g/dl
HCT	39.9	36 - 46 %
RBC	4.16	3.8 - 4.8 $\times 10^6$ / μ L
MCV	95.9	83 - 101 fL
MCH	30.0	27 - 32 pg
MCHC	31.3	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	12.9	11.6 - 14.0 %
Total Leukocyte Count(TLC)	16.10	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	87.9	40 - 80 %
Lymphocytes	8.1	20 - 40 %
Monocytes	3.9	2 - 10 %
Eosinophils	0.0	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	14.15	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	1.30	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.63	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.00	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.02	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	296	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 16:49
Patient Name	: anonymous	Age/Sex	: 16 Yr/M	Acceptance Date : 01-Jan-2023 19:50
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT030	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	11.0	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.1	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 02:18
Patient Name	: anonymous	Age/Sex	: 30 Yr/M	Acceptance Date : 01-Jan-2023 02:20
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP005	Reporting Date : 10-Jan-2023 15:47
Referred Doctor	:	Patient Type	: OPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	81.0	23.2 - 32.4 sec

Test done twice. Suggested clinical correlation.

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:50
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 01-Jan-2023 19:53
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP028	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	25.7	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 02:18
Patient Name	: anonymous	Age/Sex	: 30 Yr/M	Acceptance Date : 01-Jan-2023 02:19
Sample Type	: Edta Blood	Lab No.	: 230101CB041	Reporting Date : 10-Jan-2023 16:19
Referred Doctor	:	Patient Type	: OPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood

METHOD: Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	8.7	13 - 17 g/dl
HCT	29.0	40 - 50 %
RBC	3.09	4.5 - 5.5 $\times 10^6$ / μ L
MCV	93.9	83 - 101 fL
MCH	28.2	27 - 32 pg
MCHC	30.0	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	14.6	11.6 - 14.0 %
Total Leukocyte Count(TLC)	30.33	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	66.8	40 - 80 %
Lymphocytes	31.7	20 - 40 %
Monocytes	1.2	2 - 10 %
Eosinophils	0.1	01 - 06 %
Basophil	0.2	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	20.26	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	9.62	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.36	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.03	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.06	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	259	150 - 410 $\times 10^3$ / μ L

Comments on PBS : Leucocytosis. No atypical cells seen.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	2 Wk/F
Sample Type	:	Edta Blood	Lab No.	:	230101CB102
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	01-Jan-2023 12:12
			Acceptance Date	:	01-Jan-2023 16:59
			Reporting Date	:	05-Jan-2023 23:42

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	8.1	11.5 - 21.5 g/dl
HCT	27.2	33 - 66 %
RBC	2.78	3 - 6.3 x10 ⁶ /μL
MCV	97.8	86 - 126 fL
MCH	29.1	30 - 37 pg
MCHC	29.8	28 - 38 g/dL
Red Cell Distribution Width (RDW CV)	23.9	11.6 - 14.0 %
Total Leukocyte Count(TLC)	38.20	5 - 22 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	58.7	20 - 60 %
Lymphocytes	25.7	31 - 71 %
Monocytes	15.2	2 - 10 %
Eosinophils	0.3	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	22.40	3 - 9 x10 ³ /μL
Absolute Lymphocyte Count	9.80	2 - 16 x10 ³ /μL
Absolute Monocyte Count	5.82	0.1 - 1.2 x10 ³ /μL
Absolute Eosinophil Count	0.13	0.1 - 2 x10 ³ /μL
Absolute Basophil Count	0.05	0.02 - 0.10 x10 ³ /μL
Platelet Count	260	160 - 500 x10 ³ /μL

Comments on PBS : Nucleated RBC 3/100 WBCs. Total count raised with shift to left in myeloid series. Myelo/ metamyelocytes 12%. Neutrophils/ band form 59%, Lymphocytes 25%, Monocyte 4%.Platelets adequate singly scattered as well as in small clumps.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 12:20
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 01-Jan-2023 14:49
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT027	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	11.2	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.2	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:50
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 01-Jan-2023 19:49
Sample Type	: Edta Blood	Lab No.	: 230101CB131	Reporting Date : 10-Jan-2023 16:12
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	10.8	13 - 17 g/dl
HCT	32.5	40 - 50 %
RBC	4.17	4.5 - 5.5 x10 ⁶ /μL
MCV	77.9	83 - 101 fL
MCH	25.9	27 - 32 pg
MCHC	33.2	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	16.8	11.6 - 14.0 %
Total Leukocyte Count(TLC)	9.17	4 - 10 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	71.2	40 - 80 %
Lymphocytes	17.9	20 - 40 %
Monocytes	10.0	2 - 10 %
Eosinophils	0.7	01 - 06 %
Basophil	0.2	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	6.53	2 - 7 x10 ³ /μL
Absolute Lymphocyte Count	1.64	1 - 3 x10 ³ /μL
Absolute Monocyte Count	0.92	0.2 - 1.0 x10 ³ /μL
Absolute Eosinophil Count	0.06	0.02 - 0.5 x10 ³ /μL
Absolute Basophil Count	0.02	0.02 - 0.10 x10 ³ /μL
Platelet Count	245	150 - 410 x10 ³ /μL

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 49 Yr/F	Acceptance Date : 02-Jan-2023 02:22
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102AP012	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: OPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	25.0	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 12:35
Patient Name	: anonymous	Age/Sex	: 10 Yr/M	Acceptance Date : 01-Jan-2023 14:51
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP018	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	23.4	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:09
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 01-Jan-2023 22:59
Sample Type	: Edta	Lab No.	: 2	Reporting Date : 02-Jan-2023 10:13
Referred Doctor	:	Patient Type	: IPD	

ABSOLUTE RETICULOCYTE COUNT

SPECIMEN: EDTA

METHOD: Impedence Flow Cytometry

CLINICAL DETAILS:

TEST NAME	RESULTS	UNITS	Ref Range
Absolute Reticulocyte Count (ARC)	102.8	1000/microL	50 - 100

Comments

- Absolute reticulocyte count is a good indicator of red cell production and helps in distinguishing hypo and hyper proliferative anemias.
- It is used to assess the recovery of bone marrow function.
- It is also used to check the effectiveness of treatment of anemia.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:27
Patient Name	: anonymous	Age/Sex	: 8 D/M	Acceptance Date : 01-Jan-2023 17:00
Sample Type	: Edta Blood	Lab No.	: 230101CB103	Reporting Date : 10-Jan-2023 16:15
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	12.1	11.5 - 21.5 g/dl
HCT	41.6	33 - 66 %
RBC	4.35	3 - 6.3 x10 ⁶ /μL
MCV	95.6	86 - 126 fL
MCH	27.8	30 - 37 pg
MCHC	29.1	28 - 38 g/dL
Red Cell Distribution Width (RDW CV)	24.8	11.6 - 14.0 %
Total Leukocyte Count(TLC)	10.53	5 - 22 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	30.2	20 - 60 %
Lymphocytes	59.4	31 - 71 %
Monocytes	9.8	2 - 10 %
Eosinophils	0.5	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	3.18	3 - 9 x10 ³ /μL
Absolute Lymphocyte Count	6.26	2 - 16 x10 ³ /μL
Absolute Monocyte Count	1.03	0.1 - 1.2 x10 ³ /μL
Absolute Eosinophil Count	0.05	0.1 - 2 x10 ³ /μL
Absolute Basophil Count	0.01	0.02 - 0.10 x10 ³ /μL
Platelet Count	59	160 - 500 x10 ³ /μL

Comments on PBS : Nucleated RBC 2/100 WBC. Moderate thrombocytopenia.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:24
Patient Name	: anonymous	Age/Sex	: 46 Yr/F	Acceptance Date : 01-Jan-2023 17:01
Sample Type	: Edta Blood	Lab No.	: 230101CB112	Reporting Date : 10-Jan-2023 16:15
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	7.2	12 - 15 g/dl
HCT	24.8	36 - 46 %
RBC	3.25	3.8 - 4.8 x10 ⁶ /μL
MCV	76.3	83 - 101 fL
MCH	22.2	27 - 32 pg
MCHC	29.0	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	18.8	11.6 - 14.0 %
Total Leukocyte Count(TLC)	21.76	4 - 10 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	84.9	40 - 80 %
Lymphocytes	10.0	20 - 40 %
Monocytes	4.0	2 - 10 %
Eosinophils	0.8	01 - 06 %
Basophil	0.3	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	18.47	2 - 7 x10 ³ /μL
Absolute Lymphocyte Count	2.18	1 - 3 x10 ³ /μL
Absolute Monocyte Count	0.87	0.2 - 1.0 x10 ³ /μL
Absolute Eosinophil Count	0.18	0.02 - 0.5 x10 ³ /μL
Absolute Basophil Count	0.06	0.02 - 0.10 x10 ³ /μL
Platelet Count	322	150 - 410 x10 ³ /μL

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 02:18
Patient Name	: anonymous	Age/Sex	: 30 Yr/M	Acceptance Date : 01-Jan-2023 02:20
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT009	Reporting Date : 10-Jan-2023 15:49
Referred Doctor	:	Patient Type	: OPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	29.4	09.6 - 12.2 sec
International Normalised Ratio (INR)	3.0	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 12:20
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 01-Jan-2023 14:49
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP016	Reporting Date : 10-Jan-2023 15:47
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	25.1	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 16:49
Patient Name	: anonymous	Age/Sex	: 16 Yr/M	Acceptance Date : 01-Jan-2023 19:50
Sample Type	: Edta Blood	Lab No.	: 230101CB136	Reporting Date : 10-Jan-2023 16:15
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	13.3	13 - 17 g/dl
HCT	39.8	40 - 50 %
RBC	4.80	4.5 - 5.5 $\times 10^6$ / μ L
MCV	82.9	83 - 101 fL
MCH	27.7	27 - 32 pg
MCHC	33.4	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	11.7	11.6 - 14.0 %
Total Leukocyte Count(TLC)	9.73	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	52.8	40 - 80 %
Lymphocytes	42.1	20 - 40 %
Monocytes	3.2	2 - 10 %
Eosinophils	1.8	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	5.13	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	4.10	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.31	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.18	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.01	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	209	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 12:35
Patient Name	: anonymous	Age/Sex	: 10 Yr/M	Acceptance Date : 01-Jan-2023 14:51
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT030	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	9.8	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.0	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required