



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 03:42
Patient Name	: anonymous	Age/Sex	: 31 Yr/M	Acceptance Date : 10-Jan-2023 17:15
Sample Type	: Bal	Lab No.	: MB-2023-187	Reporting Date : 03-Feb-2023 16:22
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 15.10 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	:	-	ABHA ADDRESS	:	-	
CR No	:	-	Sample No.	:	-	Collection Date : 06-Jan-2023 14:11
Patient Name	:	anonymous	Age/Sex	:	56 Yr/M	Acceptance Date : 06-Jan-2023 18:57
Sample Type	:	Bal	Lab No.	:	MB/2023/136	Reporting Date : 23-Feb-2023 13:11
Referred Doctor	:		Patient Type	:	IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 07:56
Patient Name	: anonymous	Age/Sex	: 52 Yr/M	Acceptance Date : 09-Jan-2023 16:43
Sample Type	: Fluid	Lab No.	: MB-2023-145	Reporting Date : 09-Jan-2023 16:50
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Ascitic fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 15-Jan-2023 19:57
Patient Name	: anonymous	Age/Sex	: 49 Yr/M	Acceptance Date : 17-Jan-2023 07:39
Sample Type	: Sputum	Lab No.	: MB/2023/332	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 07:46
Patient Name	: anonymous	Age/Sex	: 25 Yr/F	Acceptance Date : 09-Jan-2023 17:17
Sample Type	: Et	Lab No.	: MB/2023/178	Reporting Date : 14-Jan-2023 14:18
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 10:47
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 09-Jan-2023 17:26
Sample Type	: Bal	Lab No.	: MB/2023/163	Reporting Date : 11-Jan-2023 08:25
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 12:53
Patient Name	: anonymous	Age/Sex	: 23 Yr/M	Acceptance Date : 09-Jan-2023 16:45
Sample Type	: Fluid	Lab No.	: MB-2023-148	Reporting Date : 09-Jan-2023 16:51
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 11:39
Patient Name	: anonymous	Age/Sex	: 53 Yr/F	Acceptance Date : 02-Jan-2023 18:37
Sample Type	: Urine	Lab No.	: MB/2023/06	Reporting Date : 04-Jan-2023 09:04
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens URINE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 07:56
Patient Name	: anonymous	Age/Sex	: 25 Yr/F	Acceptance Date : 08-Jan-2023 11:09
Sample Type	: Et	Lab No.	: MB-2023-159	Reporting Date : 09-Jan-2023 16:50
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens ET

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 10:43
Patient Name	: anonymous	Age/Sex	: 62 Yr/M	Acceptance Date : 10-Jan-2023 11:57
Sample Type	: Sputum	Lab No.	: MB-2023-153	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 08:36
Patient Name	: anonymous	Age/Sex	: 50 Yr/F	Acceptance Date : 10-Jan-2023 11:58
Sample Type	: Sputum	Lab No.	: MB-2023-158	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 06:27
Patient Name	: anonymous	Age/Sex	: 69 Yr/M	Acceptance Date : 09-Jan-2023 16:42
Sample Type	: Urine	Lab No.	: MB-2023-122	Reporting Date : 09-Jan-2023 16:50
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Urine
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 11:55
Patient Name	: anonymous	Age/Sex	: 42 Yr/M	Acceptance Date : 16-Jan-2023 19:18
Sample Type	: Sputum	Lab No.	: MB /2023/341	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 10:34
Patient Name	: anonymous	Age/Sex	: 44 Yr/M	Acceptance Date : 02-Jan-2023 18:21
Sample Type	: Sputum	Lab No.	: MB/2023/30	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 19-Jan-2023 11:08
Patient Name	: anonymous	Age/Sex	: 12 Yr/M	Acceptance Date : 19-Jan-2023 17:30
Sample Type	: Tissue	Lab No.	: MB /2023/441	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 10:52
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 02-Jan-2023 18:28
Sample Type	: Sputum	Lab No.	: MB/2023/17	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 10:05
Patient Name	: anonymous	Age/Sex	: 64 Yr/M	Acceptance Date : 02-Jan-2023 18:29
Sample Type	: Sputum	Lab No.	: MB/2023/15	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 23:37
Patient Name	: anonymous	Age/Sex	: 47 Yr/F	Acceptance Date : 09-Jan-2023 17:18
Sample Type	: Csf	Lab No.	: MB/2023/176	Reporting Date : 11-Jan-2023 13:05
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (LOW)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 15-Jan-2023 12:21
Patient Name	: anonymous	Age/Sex	: 53 Yr/F	Acceptance Date : 16-Jan-2023 19:14
Sample Type	: Fluid	Lab No.	: MB /2023/346	Reporting Date : 18-Jan-2023 09:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 12:10
Patient Name	: anonymous	Age/Sex	: 2 Yr/M	Acceptance Date : 17-Jan-2023 07:41
Sample Type	: Pus	Lab No.	: MB/2023/327	Reporting Date : 18-Jan-2023 13:17
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 12:10
Patient Name	: anonymous	Age/Sex	: 2 Yr/M	Acceptance Date : 17-Jan-2023 07:41
Sample Type	: Pus	Lab No.	: MB/2023/326	Reporting Date : 18-Jan-2023 09:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens PUS
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 11:35
Patient Name	: anonymous	Age/Sex	: 32 Yr/M	Acceptance Date : 23-Jan-2023 10:05
Sample Type	: Aspirate	Lab No.	: MB-2023-457	Reporting Date : 23-Jan-2023 11:37
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks specimen- sputum

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 13:05
Patient Name	: anonymous	Age/Sex	: 25 Yr/F	Acceptance Date : 09-Jan-2023 17:24
Sample Type	: Body Fluid	Lab No.	: MB/2023/166	Reporting Date : 11-Jan-2023 13:05
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis	NOT DETECTED
Rifampicin Resistance	NOT APPLICABLE
Test Performance	VALID
Remarks	specimen-menstrual blood.

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 03:42
Patient Name	: anonymous	Age/Sex	: 31 Yr/M	Acceptance Date : 10-Jan-2023 17:15
Sample Type	: Bal	Lab No.	: MB-2023-187	Reporting Date : 11-Jan-2023 08:25
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 12:37
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 10-Jan-2023 11:55
Sample Type	: Tissue	Lab No.	: mb-2023-150	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 12:53
Patient Name	: anonymous	Age/Sex	: 23 Yr/M	Acceptance Date : 11-Jan-2023 11:10
Sample Type	: Pleural Fluid	Lab No.	: MB-2023-149	Reporting Date : 11-Jan-2023 13:05
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 14:11
Patient Name	: anonymous	Age/Sex	: 56 Yr/M	Acceptance Date : 06-Jan-2023 18:57
Sample Type	: Bal	Lab No.	: MB/2023/136	Reporting Date : 09-Jan-2023 14:36
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 11:01
Patient Name	: anonymous	Age/Sex	: 55 Yr/M	Acceptance Date : 14-Jan-2023 17:02
Sample Type	: Sputum	Lab No.	: MB /2023/357	Reporting Date : 27-Jan-2023 16:42
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 11:01
Patient Name	: anonymous	Age/Sex	: 55 Yr/M	Acceptance Date : 14-Jan-2023 17:02
Sample Type	: Sputum	Lab No.	: MB /2023/296	Reporting Date : 07-Mar-2023 15:52
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 10:25
Patient Name	: anonymous	Age/Sex	: 34 Yr/M	Acceptance Date : 02-Jan-2023 18:22
Sample Type	: Sputum	Lab No.	: MB/2023/26	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 15-Jan-2023 15:13
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 16-Jan-2023 19:17
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/342	Reporting Date : 07-Mar-2023 15:52
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 15-Jan-2023 08:34
Patient Name	: anonymous	Age/Sex	: 59 Yr/F	Acceptance Date : 16-Jan-2023 16:46
Sample Type	: Csf	Lab No.	: MB-2022-310	Reporting Date : 16-Jan-2023 20:33
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 10:47
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 11-Jan-2023 11:09
Sample Type	: Bal	Lab No.	: MB-2023-152	Reporting Date : 11-Jan-2023 13:05
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 03:45
Patient Name	: anonymous	Age/Sex	: 31 Yr/M	Acceptance Date : 09-Jan-2023 17:12
Sample Type	: Bal	Lab No.	: MB/2023/184	Reporting Date : 14-Jan-2023 14:18
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 12:38
Patient Name	: anonymous	Age/Sex	: 45 Yr/M	Acceptance Date : 09-Jan-2023 16:46
Sample Type	: Fluid	Lab No.	: MB-2023-160	Reporting Date : 09-Jan-2023 16:51
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 12:13
Patient Name	: anonymous	Age/Sex	: 61 Yr/F	Acceptance Date : 14-Jan-2023 17:00
Sample Type	: Sputum	Lab No.	: MB /2023/300	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 07:19
Patient Name	: anonymous	Age/Sex	: 29 Yr/M	Acceptance Date : 17-Jan-2023 07:47
Sample Type	: Sputum	Lab No.	: MB/2023/314	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 10:47
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 09-Jan-2023 17:26
Sample Type	: Bal	Lab No.	: MB/2023/163	Reporting Date : 23-Feb-2023 13:11
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 09:44
Patient Name	: anonymous	Age/Sex	: 17 Yr/M	Acceptance Date : 23-Jan-2023 18:01
Sample Type	: Sputum	Lab No.	: MB /2023/503	Reporting Date : 30-Jan-2023 16:22
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 16:02
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 16-Jan-2023 19:17
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/0343	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 10:14
Patient Name	: anonymous	Age/Sex	: 37 Yr/M	Acceptance Date : 02-Jan-2023 18:27
Sample Type	: Sputum	Lab No.	: MB/2023/19	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 11:32
Patient Name	: anonymous	Age/Sex	: 28 Yr/F	Acceptance Date : 02-Jan-2023 18:24
Sample Type	: Sputum	Lab No.	: MB/2023/23	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 11:20
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 11-Jan-2023 10:31
Sample Type	: Aspirate	Lab No.	: MB /2023/188	Reporting Date : 28-Feb-2023 15:12
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-
Comments	Specimen - Bone marrow

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 08-Jan-2023 19:27
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 11-Jan-2023 10:10
Sample Type	: Sputum	Lab No.	: MB /2023/197	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 19-Jan-2023 11:03
Patient Name	: anonymous	Age/Sex	: 12 Yr/M	Acceptance Date : 19-Jan-2023 17:31
Sample Type	: Tissue	Lab No.	: MB /2023/440	Reporting Date : 21-Jan-2023 11:31
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens TISSUE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 11:47
Patient Name	: anonymous	Age/Sex	: 34 Yr/M	Acceptance Date : 14-Jan-2023 16:59
Sample Type	: Urine	Lab No.	: MB /2023/301	Reporting Date : 16-Jan-2023 10:33
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 12:24
Patient Name	: anonymous	Age/Sex	: 62 Yr/M	Acceptance Date : 14-Jan-2023 17:00
Sample Type	: Ascitic Fluid	Lab No.	: MB /2023/299	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 16:08
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 16-Jan-2023 16:47
Sample Type	: Fluid	Lab No.	: MB-2022-311	Reporting Date : 16-Jan-2023 20:33
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens PLEURAL FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 14-Jan-2023 12:07
Patient Name	: anonymous	Age/Sex	: 55 Yr/M	Acceptance Date : 17-Jan-2023 07:39
Sample Type	: Csf	Lab No.	: MB/2023/331	Reporting Date : 18-Jan-2023 13:17
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 14:24
Patient Name	: anonymous	Age/Sex	: 56 Yr/M	Acceptance Date : 06-Jan-2023 18:56
Sample Type	: Bal	Lab No.	: MB/2023/137	Reporting Date : 09-Jan-2023 14:12
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required