



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 08:56
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 17-Jan-2023 19:00
Sample Type	: Urine	Lab No.	: MB/2023/353	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens URINE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 14:14
Patient Name	: anonymous	Age/Sex	: 59 Yr/F	Acceptance Date : 11-Jan-2023 17:11
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/234	Reporting Date : 28-Feb-2023 15:19
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 19:25
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 17-Jan-2023 09:17
Sample Type	: Csf	Lab No.	: MB /2023/351	Reporting Date : 18-Jan-2023 09:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 13:00
Patient Name	: anonymous	Age/Sex	: 29 Yr/M	Acceptance Date : 17-Jan-2023 07:35
Sample Type	: Sputum	Lab No.	: MB/2023/336	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 08:35
Patient Name	: anonymous	Age/Sex	: 69 Yr/M	Acceptance Date : 17-Jan-2023 07:45
Sample Type	: Sputum	Lab No.	: MB/2023/318	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 15:06
Patient Name	: anonymous	Age/Sex	: 47 Yr/F	Acceptance Date : 04-Jan-2023 19:15
Sample Type	: Body Fluid	Lab No.	: mb-2023-96	Reporting Date : 06-Jan-2023 13:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 09:37
Patient Name	: anonymous	Age/Sex	: 74 Yr/M	Acceptance Date : 03-Mar-2023 12:49
Sample Type	: Pleural Fluid	Lab No.	: mb-2023-356	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 08:32
Patient Name	: anonymous	Age/Sex	: 55 Yr/M	Acceptance Date : 17-Jan-2023 07:42
Sample Type	: Sputum	Lab No.	: MB/2023/325	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 14:14
Patient Name	: anonymous	Age/Sex	: 59 Yr/F	Acceptance Date : 11-Jan-2023 17:11
Sample Type	: Fluid	Lab No.	: MB/2023/234	Reporting Date : 12-Jan-2023 17:08
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:33
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 09-Jan-2023 17:11
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/185	Reporting Date : 23-Feb-2023 13:12
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 13:03
Patient Name	: anonymous	Age/Sex	: 26 Yr/F	Acceptance Date : 04-Jan-2023 21:22
Sample Type	: Bal	Lab No.	: MB/2023/083	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 10:39
Patient Name	: anonymous	Age/Sex	: 52 Yr/F	Acceptance Date : 27-Jan-2023 11:52
Sample Type	: Sputum	Lab No.	: MB-2023-368	Reporting Date : 27-Jan-2023 16:46
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 10:31
Patient Name	: anonymous	Age/Sex	: 33 Yr/F	Acceptance Date : 17-Jan-2023 07:44
Sample Type	: Sputum	Lab No.	: MB/2023/321	Reporting Date : 07-Mar-2023 15:52
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 08:54
Patient Name	: anonymous	Age/Sex	: 39 Yr/F	Acceptance Date : 12-Jan-2023 17:41
Sample Type	: Sputum	Lab No.	: MB/2023/247	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:33
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 09-Jan-2023 17:12
Sample Type	: Fluid	Lab No.	: MB /2023/186	Reporting Date : 11-Jan-2023 08:25
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 12:52
Patient Name	: anonymous	Age/Sex	: 40 Yr/F	Acceptance Date : 04-Jan-2023 21:43
Sample Type	: Fluid	Lab No.	: MB/2023/76	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens ASCITIC FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 04:52
Patient Name	: anonymous	Age/Sex	: 38 Yr/F	Acceptance Date : 17-Jan-2023 18:49
Sample Type	: Ascitic Fluid	Lab No.	: MB/2023/358	Reporting Date : 19-Jan-2023 12:41
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 09:05
Patient Name	: anonymous	Age/Sex	: 2 Yr/M	Acceptance Date : 17-Jan-2023 18:57
Sample Type	: Aspirate	Lab No.	: MB/2023/354	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens GASTRIC ASPIRATE
ZN Stain Positive for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 21:27
Patient Name	: anonymous	Age/Sex	: 14 Yr/M	Acceptance Date : 17-Jan-2023 18:31
Sample Type	: Sputum	Lab No.	: MB/2023/366	Reporting Date : 01-Feb-2023 16:31
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 11:25
Patient Name	: anonymous	Age/Sex	: 27 Yr/M	Acceptance Date : 16-Jan-2023 19:16
Sample Type	: Urine	Lab No.	: MB /2023/335	Reporting Date : 18-Jan-2023 13:17
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:33
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 09-Jan-2023 17:11
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/186	Reporting Date : 14-Jan-2023 14:18
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:40
Patient Name	: anonymous	Age/Sex	: 74 Yr/M	Acceptance Date : 09-Jan-2023 17:26
Sample Type	: Fluid	Lab No.	: MB/2023/162	Reporting Date : 11-Jan-2023 08:25
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 12:54
Patient Name	: anonymous	Age/Sex	: 24 Yr/F	Acceptance Date : 11-Jan-2023 13:50
Sample Type	: Urine	Lab No.	: MB-2023-211	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (LOW)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 07:31
Patient Name	: anonymous	Age/Sex	: 74 Yr/M	Acceptance Date : 17-Jan-2023 18:55
Sample Type	: Fluid	Lab No.	: MB/2023/356	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens PLEURAL FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:42
Patient Name	: anonymous	Age/Sex	: 74 Yr/M	Acceptance Date : 11-Jan-2023 10:12
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/202	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 10:31
Patient Name	: anonymous	Age/Sex	: 33 Yr/F	Acceptance Date : 17-Jan-2023 07:43
Sample Type	: Sputum	Lab No.	: MB/2023/322	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 11:34
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 17-Jan-2023 07:39
Sample Type	: Bal	Lab No.	: MB/2023/330	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 07:43
Patient Name	: anonymous	Age/Sex	: 23 Yr/M	Acceptance Date : 17-Jan-2023 18:12
Sample Type	: Csf	Lab No.	: MB/2023/386	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 07:41
Patient Name	: anonymous	Age/Sex	: 23 Yr/M	Acceptance Date : 17-Jan-2023 18:11
Sample Type	: Csf	Lab No.	: MB/2023/387	Reporting Date : 19-Jan-2023 12:41
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 11:36
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 17-Jan-2023 07:40
Sample Type	: Sputum	Lab No.	: MB/2023/328	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 15:26
Patient Name	: anonymous	Age/Sex	: 26 Yr/F	Acceptance Date : 04-Jan-2023 21:21
Sample Type	: Bal	Lab No.	: MB/2023/83	Reporting Date : 23-Feb-2023 13:11
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 12:37
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 11-Jan-2023 17:13
Sample Type	: Ascitic Fluid	Lab No.	: MB/2023/230	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 09:50
Patient Name	: anonymous	Age/Sex	: 27 Yr/F	Acceptance Date : 16-Jan-2023 19:06
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/349	Reporting Date : 07-Mar-2023 15:52
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 00:08
Patient Name	: anonymous	Age/Sex	: 18 Yr/M	Acceptance Date : 17-Jan-2023 18:46
Sample Type	: Et	Lab No.	: MB/2023/359	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens ET

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 23:09
Patient Name	: anonymous	Age/Sex	: 42 Yr/F	Acceptance Date : 16-Jan-2023 16:50
Sample Type	: Sputum	Lab No.	: MB-2023-297	Reporting Date : 16-Jan-2023 20:33
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 13:09
Patient Name	: anonymous	Age/Sex	: 26 Yr/F	Acceptance Date : 04-Jan-2023 21:20
Sample Type	: Bal	Lab No.	: MB/2023/84	Reporting Date : 10-Jan-2023 15:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 15:40
Patient Name	: anonymous	Age/Sex	: 74 Yr/M	Acceptance Date : 09-Jan-2023 17:26
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/162	Reporting Date : 23-Feb-2023 13:12
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 11:04
Patient Name	: anonymous	Age/Sex	: 31 Yr/M	Acceptance Date : 17-Jan-2023 07:42
Sample Type	: Sputum	Lab No.	: MB/2023/324	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 13:35
Patient Name	: anonymous	Age/Sex	: 59 Yr/F	Acceptance Date : 11-Jan-2023 17:11
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/235	Reporting Date : 16-Jan-2023 17:31
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 04-Jan-2023 14:52
Patient Name	: anonymous	Age/Sex	: 34 Yr/M	Acceptance Date : 04-Jan-2023 19:16
Sample Type	: Aspirate	Lab No.	: mb/2023/95	Reporting Date : 06-Jan-2023 13:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 02:06
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 17-Jan-2023 18:44
Sample Type	: Csf	Lab No.	: MB/2023/361	Reporting Date : 19-Jan-2023 12:41
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 23:22
Patient Name	: anonymous	Age/Sex	: 68 Yr/F	Acceptance Date : 17-Jan-2023 18:42
Sample Type	: Sputum	Lab No.	: MB/2023/363	Reporting Date : 27-Jan-2023 16:43
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 09-Jan-2023 13:17
Patient Name	: anonymous	Age/Sex	: 35 Yr/F	Acceptance Date : 11-Jan-2023 10:24
Sample Type	: Sputum	Lab No.	: MB /2023/192	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 14:44
Patient Name	: anonymous	Age/Sex	: 21 Yr/M	Acceptance Date : 11-Jan-2023 17:12
Sample Type	: Pus	Lab No.	: MB/2023/232	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 00:08
Patient Name	: anonymous	Age/Sex	: 18 Yr/M	Acceptance Date : 17-Jan-2023 18:45
Sample Type	: Et	Lab No.	: MB/2023/360	Reporting Date : 19-Jan-2023 12:41
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 01:08
Patient Name	: anonymous	Age/Sex	: 58 Yr/F	Acceptance Date : 17-Jan-2023 18:39
Sample Type	: Sputum	Lab No.	: MB/2023/364	Reporting Date : 27-Jan-2023 16:44
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 09:10
Patient Name	: anonymous	Age/Sex	: 55 Yr/M	Acceptance Date : 17-Jan-2023 09:18
Sample Type	: Csf	Lab No.	: MB /2023/350	Reporting Date : 18-Jan-2023 09:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 22:42
Patient Name	: anonymous	Age/Sex	: 38 Yr/F	Acceptance Date : 20-Jan-2023 12:57
Sample Type	: Fluid	Lab No.	: MB-2023-406	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Ascitic fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 10:18
Patient Name	: anonymous	Age/Sex	: 32 Yr/F	Acceptance Date : 17-Jan-2023 07:43
Sample Type	: Sputum	Lab No.	: MB/2023/323	Reporting Date : 20-Jan-2023 12:57
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 16-Jan-2023 21:31
Patient Name	: anonymous	Age/Sex	: 14 Yr/M	Acceptance Date : 17-Jan-2023 18:33
Sample Type	: Sputum	Lab No.	: MB/2023/365	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens SPUTUM

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required