



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 13:04
Patient Name	: anonymous	Age/Sex	: 44 Yr/F	Acceptance Date : 12-Jan-2023 17:34
Sample Type	: Bal	Lab No.	: MB/2023/259	Reporting Date : 23-Jan-2023 07:02
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL

ZN Stain Positive for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 08:49
Patient Name	: anonymous	Age/Sex	: 51 Yr/F	Acceptance Date : 20-Jan-2023 10:22
Sample Type	: Sputum	Lab No.	: MB-2023-412	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 11:17
Patient Name	: anonymous	Age/Sex	: 36 Yr/F	Acceptance Date : 12-Jan-2023 17:52
Sample Type	: Sputum	Lab No.	: MB/2023/241	Reporting Date : 13-Jan-2023 17:04
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens SPUTUM

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 13:34
Patient Name	: anonymous	Age/Sex	: 11 Yr/M	Acceptance Date : 17-Jan-2023 18:17
Sample Type	: Aspirate	Lab No.	: MB/2023/382	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens ASPIRATE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:38
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 12-Jan-2023 17:32
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/262	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 09:39
Patient Name	: anonymous	Age/Sex	: 55 Yr/F	Acceptance Date : 12-Jan-2023 09:43
Sample Type	: Fluid	Lab No.	: MB /2023/238	Reporting Date : 12-Jan-2023 17:08
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Ascitic fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 10:26
Patient Name	: anonymous	Age/Sex	: 34 Yr/F	Acceptance Date : 20-Jan-2023 10:24
Sample Type	: Sputum	Lab No.	: MB-2023-414	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 12:46
Patient Name	: anonymous	Age/Sex	: 40 Yr/M	Acceptance Date : 12-Jan-2023 17:52
Sample Type	: Fluid	Lab No.	: MB/2023/240	Reporting Date : 13-Jan-2023 17:04
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens ASCITIC FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:35
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 12-Jan-2023 17:32
Sample Type	: Fluid	Lab No.	: MB/2023/262	Reporting Date : 16-Feb-2023 13:32
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens PLEURAL FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 13:04
Patient Name	: anonymous	Age/Sex	: 44 Yr/F	Acceptance Date : 12-Jan-2023 17:34
Sample Type	: Bal	Lab No.	: MB/2023/260	Reporting Date : 16-Jan-2023 17:18
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 14:10
Patient Name	: anonymous	Age/Sex	: 19 Yr/M	Acceptance Date : 18-Jan-2023 18:33
Sample Type	: Sputum	Lab No.	: 419	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 13:30
Patient Name	: anonymous	Age/Sex	: 53 Yr/M	Acceptance Date : 20-Jan-2023 12:59
Sample Type	: Urine	Lab No.	: MB-2023-417	Reporting Date : 15-Feb-2023 12:11
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 19 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 11:11
Patient Name	: anonymous	Age/Sex	: 56 Yr/F	Acceptance Date : 14-Jan-2023 16:54
Sample Type	: Sputum	Lab No.	: MB /2023/308	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 14:32
Patient Name	: anonymous	Age/Sex	: 30 Yr/F	Acceptance Date : 19-Jan-2023 17:05
Sample Type	: Urine	Lab No.	: MB /2023/425	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 08:49
Patient Name	: anonymous	Age/Sex	: 51 Yr/F	Acceptance Date : 20-Jan-2023 12:58
Sample Type	: Sputum	Lab No.	: MB-2023-411	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Sputum
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 11:09
Patient Name	: anonymous	Age/Sex	: 11 Yr/M	Acceptance Date : 12-Jan-2023 17:39
Sample Type	: Pus	Lab No.	: MB/2023/250	Reporting Date : 13-Jan-2023 17:04
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens PUS
ZN Stain Positive for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 12:08
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 19-Jan-2023 09:21
Sample Type	: Bal	Lab No.	: MB /2023/421	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 21.1 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Resistant
Comments	Drug Susceptibility to second line Drugs: - Quinolone " Resistant (gyrA - MUT3A present). SLID - Susceptible

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 12:43
Patient Name	: anonymous	Age/Sex	: 53 Yr/M	Acceptance Date : 18-Jan-2023 18:35
Sample Type	: Urine	Lab No.	: 418	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 13:04
Patient Name	: anonymous	Age/Sex	: 44 Yr/F	Acceptance Date : 12-Jan-2023 17:34
Sample Type	: Bal	Lab No.	: MB/2023/259	Reporting Date : 03-Feb-2023 16:22
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Positive for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 4.18 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 14:50
Patient Name	: anonymous	Age/Sex	: 36 Yr/F	Acceptance Date : 17-Jan-2023 18:05
Sample Type	: Bal	Lab No.	: MB/2023/393	Reporting Date : 27-Jan-2023 16:50
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:56
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 12-Jan-2023 17:29
Sample Type	: Bal	Lab No.	: MB/2023/267	Reporting Date : 16-Jan-2023 17:18
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance INDETERMINATE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 23:37
Patient Name	: anonymous	Age/Sex	: 24 Yr/F	Acceptance Date : 20-Jan-2023 10:26
Sample Type	: Pleural Fluid	Lab No.	: MB-2023-407	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 14:04
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 20-Jan-2023 10:19
Sample Type	: Sputum	Lab No.	: MB-2023-416	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 14:47
Patient Name	: anonymous	Age/Sex	: 21 Yr/M	Acceptance Date : 11-Jan-2023 17:13
Sample Type	: Pus	Lab No.	: MB/2023/231	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 13:56
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 19-Jan-2023 17:35
Sample Type	: Biopsy	Lab No.	: MB /2023/437	Reporting Date : 23-Jan-2023 11:37
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 08:08
Patient Name	: anonymous	Age/Sex	: 2 Yr/F	Acceptance Date : 19-Jan-2023 17:46
Sample Type	: Aspirate	Lab No.	: MB /2023/427	Reporting Date : 21-Jan-2023 12:47
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis	NOT DETECTED
Rifampicin Resistance	NOT APPLICABLE
Test Performance	VALID
Remarks	Specimen - Gastric aspirate

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 12:08
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 21-Jan-2023 10:07
Sample Type	: Bal	Lab No.	: MB-2023-415	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 15:05
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 12-Jan-2023 17:30
Sample Type	: Bal	Lab No.	: MB/2023/266	Reporting Date : 28-Feb-2023 15:25
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 14:43
Patient Name	: anonymous	Age/Sex	: 45 Yr/M	Acceptance Date : 17-Jan-2023 18:17
Sample Type	: Sputum	Lab No.	: MB/2023/381	Reporting Date : 27-Jan-2023 16:48
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:49
Patient Name	: anonymous	Age/Sex	: 17 Yr/M	Acceptance Date : 12-Jan-2023 17:31
Sample Type	: Bal	Lab No.	: MB/2023/265	Reporting Date : 16-Jan-2023 17:18
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 10:16
Patient Name	: anonymous	Age/Sex	: 28 Yr/M	Acceptance Date : 12-Jan-2023 17:38
Sample Type	: Biopsy	Lab No.	: MB/2023/253	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	No growth of Mycobacterium species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:35
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 12-Jan-2023 17:32
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/263	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis

Rifampicin Resistance

Test Performance INVALID

Remarks Please send repeat sample after aseptic collection.

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:49
Patient Name	: anonymous	Age/Sex	: 17 Yr/M	Acceptance Date : 14-Jan-2023 09:28
Sample Type	: Bal	Lab No.	: MB /2023/275	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 11:08
Patient Name	: anonymous	Age/Sex	: 11 Yr/M	Acceptance Date : 12-Jan-2023 17:39
Sample Type	: Pus	Lab No.	: MB/2023/251	Reporting Date : 23-Jan-2023 11:37
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 23:21
Patient Name	: anonymous	Age/Sex	: 14 Yr/F	Acceptance Date : 20-Jan-2023 10:33
Sample Type	: Csf	Lab No.	: MB-2023-409	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:12
Patient Name	: anonymous	Age/Sex	: 60 Yr/F	Acceptance Date : 12-Jan-2023 17:35
Sample Type	: Sputum	Lab No.	: MB/2023/258	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 15:05
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 12-Jan-2023 17:30
Sample Type	: Bal	Lab No.	: MB/2023/266	Reporting Date : 23-Jan-2023 07:02
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 12:08
Patient Name	: anonymous	Age/Sex	: 20 Yr/F	Acceptance Date : 19-Jan-2023 09:21
Sample Type	: Bal	Lab No.	: MB /2023/421	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 13:25
Patient Name	: anonymous	Age/Sex	: 53 Yr/F	Acceptance Date : 17-Jan-2023 18:03
Sample Type	: Bal	Lab No.	: MB/2023/395	Reporting Date : 27-Jan-2023 16:51
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 23:42
Patient Name	: anonymous	Age/Sex	: 54 Yr/F	Acceptance Date : 12-Jan-2023 17:36
Sample Type	: Csf	Lab No.	: MB/2023/256	Reporting Date : 13-Jan-2023 17:04
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 23:21
Patient Name	: anonymous	Age/Sex	: 14 Yr/F	Acceptance Date : 20-Jan-2023 12:57
Sample Type	: Csf	Lab No.	: MB-2023-408	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:41
Patient Name	: anonymous	Age/Sex	: 14 Yr/M	Acceptance Date : 12-Jan-2023 17:31
Sample Type	: Bal	Lab No.	: MB/2023/264	Reporting Date : 16-Jan-2023 17:31
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 10:37
Patient Name	: anonymous	Age/Sex	: 33 Yr/F	Acceptance Date : 12-Jan-2023 17:51
Sample Type	: Sputum	Lab No.	: MB/2023/242	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:41
Patient Name	: anonymous	Age/Sex	: 14 Yr/M	Acceptance Date : 13-Jan-2023 09:37
Sample Type	: Bal	Lab No.	: MB /2023/269	Reporting Date : 23-Jan-2023 07:02
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 08:49
Patient Name	: anonymous	Age/Sex	: 51 Yr/F	Acceptance Date : 20-Jan-2023 12:58
Sample Type	: Sputum	Lab No.	: MB-2023-411	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	- No growth of Mycobacterium species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 14:49
Patient Name	: anonymous	Age/Sex	: 17 Yr/M	Acceptance Date : 14-Jan-2023 09:28
Sample Type	: Bal	Lab No.	: MB /2023/274	Reporting Date : 14-Jan-2023 14:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 08:45
Patient Name	: anonymous	Age/Sex	: 51 Yr/F	Acceptance Date : 14-Jan-2023 16:56
Sample Type	: Sputum	Lab No.	: MB /2023/306	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 10:16
Patient Name	: anonymous	Age/Sex	: 28 Yr/M	Acceptance Date : 12-Jan-2023 17:38
Sample Type	: Tissue	Lab No.	: MB/2023/253	Reporting Date : 13-Jan-2023 17:04
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens TISSUE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 13:15
Patient Name	: anonymous	Age/Sex	: 9 Mth/F	Acceptance Date : 17-Jan-2023 18:09
Sample Type	: Tissue	Lab No.	: MB/2023/389	Reporting Date : 19-Jan-2023 12:41
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	:	-	ABHA ADDRESS	:	-	
CR No	:	-	Sample No.	:	-	Collection Date : 12-Jan-2023 11:17
Patient Name	:	anonymous	Age/Sex	:	36 Yr/F	Acceptance Date : 12-Jan-2023 17:52
Sample Type	:	Sputum	Lab No.	:	MB/2023/241	Reporting Date : 28-Feb-2023 15:22
Referred Doctor	:		Patient Type	:	IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required