



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 15:40
Patient Name	: anonymous	Age/Sex	: 53 Yr/F	Acceptance Date : 17-Jan-2023 18:04
Sample Type	: Bal	Lab No.	: MB/2023/394	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	- No growth of Mycobacterium species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:03
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 19-Jan-2023 17:13
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/452	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 15:33
Patient Name	: anonymous	Age/Sex	: 36 Yr/F	Acceptance Date : 17-Jan-2023 18:06
Sample Type	: Bal	Lab No.	: MB/2023/392	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	- No growth of Mycobacterium species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 07:42
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 03-Jan-2023 16:47
Sample Type	: Fluid	Lab No.	: MB/2023/46	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens PLEURAL FLUID
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 12:58
Patient Name	: anonymous	Age/Sex	: 35 Yr/M	Acceptance Date : 16-Jan-2023 19:13
Sample Type	: Sputum	Lab No.	: MB /2023/348	Reporting Date : 15-Feb-2023 12:12
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Positive for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 14.5 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 13:16
Patient Name	: anonymous	Age/Sex	: 35 Yr/F	Acceptance Date : 11-Jan-2023 10:00
Sample Type	: Body Fluid	Lab No.	: MB /2023/204	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:05
Patient Name	: anonymous	Age/Sex	: 86 Yr/M	Acceptance Date : 19-Jan-2023 17:06
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/426	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (LOW)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 09:48
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 14-Jan-2023 09:25
Sample Type	: Csf	Lab No.	: MB /2023/278	Reporting Date : 16-Jan-2023 12:32
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:03
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 19-Jan-2023 09:23
Sample Type	: Fluid	Lab No.	: MB /2023/422	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:06
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 03-Jan-2023 16:42
Sample Type	: Ascitic Fluid	Lab No.	: MB/2023/56	Reporting Date : 05-Jan-2023 13:30
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:06
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 03-Jan-2023 16:42
Sample Type	: Fluid	Lab No.	: MB/2023/55	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens FLUID

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 09:25
Patient Name	: anonymous	Age/Sex	: 48 Yr/M	Acceptance Date : 03-Jan-2023 16:47
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/47	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 11:54
Patient Name	: anonymous	Age/Sex	: 27 Yr/F	Acceptance Date : 11-Jan-2023 10:03
Sample Type	: Sputum	Lab No.	: MB /2023/200	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:20
Patient Name	: anonymous	Age/Sex	: 23 Yr/M	Acceptance Date : 03-Jan-2023 16:39
Sample Type	: Sputum	Lab No.	: MB/2023/62	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 13:35
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 03-Jan-2023 16:40
Sample Type	: Urine	Lab No.	: MB/2023/60	Reporting Date : 05-Jan-2023 13:30
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 11:06
Patient Name	: anonymous	Age/Sex	: 72 Yr/F	Acceptance Date : 14-Jan-2023 16:58
Sample Type	: Csf	Lab No.	: MB /2023/303	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 10:23
Patient Name	: anonymous	Age/Sex	: 52 Yr/F	Acceptance Date : 14-Jan-2023 09:26
Sample Type	: Et	Lab No.	: MB /2023/277	Reporting Date : 16-Jan-2023 12:32
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 13:39
Patient Name	: anonymous	Age/Sex	: 32 Yr/F	Acceptance Date : 11-Jan-2023 09:57
Sample Type	: Tissue	Lab No.	: MB /2023/206	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (VERY LOW)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 11:55
Patient Name	: anonymous	Age/Sex	: 29 Yr/F	Acceptance Date : 11-Jan-2023 09:58
Sample Type	: Sputum	Lab No.	: MB /2023/205	Reporting Date : 12-Jan-2023 17:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:42
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 03-Jan-2023 16:43
Sample Type	: Sputum	Lab No.	: MB/2023/54	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 09:15
Patient Name	: anonymous	Age/Sex	: 64 Yr/M	Acceptance Date : 03-Jan-2023 16:53
Sample Type	: Sputum	Lab No.	: MB/2023/44	Reporting Date : 23-Feb-2023 13:03
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 19:05
Patient Name	: anonymous	Age/Sex	: 80 Yr/M	Acceptance Date : 19-Jan-2023 09:29
Sample Type	: Csf	Lab No.	: MB /2023/424	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 08:24
Patient Name	: anonymous	Age/Sex	: 5 Yr/M	Acceptance Date : 06-Jan-2023 19:17
Sample Type	: Aspirate	Lab No.	: MB/2023/119	Reporting Date : 09-Jan-2023 14:36
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Gastric aspirate
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 15:40
Patient Name	: anonymous	Age/Sex	: 53 Yr/F	Acceptance Date : 17-Jan-2023 18:04
Sample Type	: Bal	Lab No.	: MB/2023/394	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	42 Yr/M
Sample Type	:	Bal	Lab No.	:	MB/2023/33
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	02-Jan-2023 14:31
			Acceptance Date	:	02-Jan-2023 18:19
			Reporting Date	:	23-Feb-2023 13:03

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	:	-	ABHA ADDRESS	:	-	
CR No	:	-	Sample No.	:	-	Collection Date : 10-Jan-2023 13:15
Patient Name	:	anonymous	Age/Sex	:	35 Yr/F	Acceptance Date : 11-Jan-2023 10:00
Sample Type	:	Body Fluid	Lab No.	:	MB /2023/203	Reporting Date : 28-Feb-2023 15:19
Referred Doctor	:		Patient Type	:	OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 13:16
Patient Name	: anonymous	Age/Sex	: 35 Yr/F	Acceptance Date : 11-Jan-2023 10:00
Sample Type	: Aspirate	Lab No.	: MB /2023/203	Reporting Date : 11-Jan-2023 17:05
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens body fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 12-Jan-2023 13:44
Patient Name	: anonymous	Age/Sex	: 39 Yr/M	Acceptance Date : 14-Jan-2023 09:24
Sample Type	: Csf	Lab No.	: MB /2023/279	Reporting Date : 16-Jan-2023 12:32
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 13:36
Patient Name	: anonymous	Age/Sex	: 42 Yr/M	Acceptance Date : 02-Jan-2023 18:18
Sample Type	: Bal	Lab No.	: MB/2023/34	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 14:27
Patient Name	: anonymous	Age/Sex	: 52 Yr/M	Acceptance Date : 18-Jan-2023 18:32
Sample Type	: Sputum	Lab No.	: 420	Reporting Date : 20-Jan-2023 12:51
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 08:24
Patient Name	: anonymous	Age/Sex	: 5 Yr/M	Acceptance Date : 06-Jan-2023 19:01
Sample Type	: Aspirate	Lab No.	: MB/2023/128	Reporting Date : 09-Jan-2023 14:11
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks Specimen - Gastric Aspirate

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 13:57
Patient Name	: anonymous	Age/Sex	: 42 Yr/M	Acceptance Date : 02-Jan-2023 18:19
Sample Type	: Bal	Lab No.	: MB /2023/33	Reporting Date : 04-Jan-2023 09:04
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:05
Patient Name	: anonymous	Age/Sex	: 86 Yr/M	Acceptance Date : 19-Jan-2023 09:27
Sample Type	: Fluid	Lab No.	: MB /2023/423	Reporting Date : 20-Jan-2023 13:28
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 09:15
Patient Name	: anonymous	Age/Sex	: 64 Yr/M	Acceptance Date : 03-Jan-2023 16:53
Sample Type	: Sputum	Lab No.	: MB/2023/44	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens SPUTUM

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:03
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 19-Jan-2023 09:23
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/422	Reporting Date : 15-Feb-2023 12:11
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 18.14 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 19:05
Patient Name	: anonymous	Age/Sex	: 80 Yr/M	Acceptance Date : 19-Jan-2023 17:45
Sample Type	: Csf	Lab No.	: MB /2023/429	Reporting Date : 23-Jan-2023 11:42
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 11:44
Patient Name	: anonymous	Age/Sex	: 44 Yr/F	Acceptance Date : 11-Jan-2023 17:19
Sample Type	: Sputum	Lab No.	: MB/2023/221	Reporting Date : 16-Jan-2023 17:18
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (MEDIUM)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 11:06
Patient Name	: anonymous	Age/Sex	: 34 Yr/M	Acceptance Date : 14-Jan-2023 16:53
Sample Type	: Sputum	Lab No.	: MB /2023/309	Reporting Date : 17-Jan-2023 13:13
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:20
Patient Name	: anonymous	Age/Sex	: 65 Yr/F	Acceptance Date : 03-Jan-2023 16:45
Sample Type	: Ascitic Fluid	Lab No.	: MB/2023/50	Reporting Date : 05-Jan-2023 13:30
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 18-Jan-2023 15:05
Patient Name	: anonymous	Age/Sex	: 86 Yr/M	Acceptance Date : 19-Jan-2023 09:27
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/423	Reporting Date : 15-Feb-2023 12:11
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Growth of Mycobacterium tuberculosis complex after 16.19 days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Susceptible Isoniazid- Susceptible

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 15:33
Patient Name	: anonymous	Age/Sex	: 36 Yr/F	Acceptance Date : 17-Jan-2023 18:06
Sample Type	: Bal	Lab No.	: MB/2023/392	Reporting Date : 19-Jan-2023 16:37
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 13-Jan-2023 11:06
Patient Name	: anonymous	Age/Sex	: 72 Yr/F	Acceptance Date : 14-Jan-2023 09:19
Sample Type	: Csf	Lab No.	: MB /2023/270	Reporting Date : 14-Jan-2023 14:32
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens CSF
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 16:40
Patient Name	: anonymous	Age/Sex	: 24 Yr/F	Acceptance Date : 03-Mar-2023 12:55
Sample Type	: Pleural Fluid	Lab No.	: MB-2023-396	Reporting Date : 04-Mar-2023 09:14
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	- No growth of Mycobacterium species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	48 Yr/M
Sample Type	:	Pleural Fluid	Lab No.	:	MB/2023/46
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	03-Jan-2023 07:42
			Acceptance Date	:	03-Jan-2023 16:47
			Reporting Date	:	23-Feb-2023 13:03

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 09:00
Patient Name	: anonymous	Age/Sex	: 5 Yr/F	Acceptance Date : 03-Jan-2023 16:45
Sample Type	: Sputum	Lab No.	: MB/2023/49	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:54
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 03-Jan-2023 16:44
Sample Type	: Sputum	Lab No.	: MB/2023/53	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens SPUTUM
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:54
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 03-Jan-2023 16:44
Sample Type	: Sputum	Lab No.	: MB/2023/53	Reporting Date : 23-Feb-2023 13:03
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 16:37
Patient Name	: anonymous	Age/Sex	: 68 Yr/M	Acceptance Date : 03-Jan-2023 16:54
Sample Type	: Tt	Lab No.	: MB/2023/43	Reporting Date : 23-Feb-2023 13:03
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 10:33
Patient Name	: anonymous	Age/Sex	: 39 Yr/F	Acceptance Date : 03-Jan-2023 16:44
Sample Type	: Bal	Lab No.	: MB/2023/52	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 17-Jan-2023 14:51
Patient Name	: anonymous	Age/Sex	: 38 Yr/M	Acceptance Date : 17-Jan-2023 18:06
Sample Type	: Bal	Lab No.	: MB/2023/391	Reporting Date : 27-Jan-2023 16:50
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required