



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:59
Patient Name	: anonymous	Age/Sex	: 35 Yr/M	Acceptance Date : 11-Jan-2023 09:50
Sample Type	: Pleural Fluid	Lab No.	: MB /2023/210	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 14:28
Patient Name	: anonymous	Age/Sex	: 11 Yr/F	Acceptance Date : 05-Jan-2023 19:00
Sample Type	: Pus	Lab No.	: MB/2023/111	Reporting Date : 06-Jan-2023 20:09
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Pus
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 19:25
Patient Name	: anonymous	Age/Sex	: 7 Yr/M	Acceptance Date : 11-Jan-2023 17:34
Sample Type	: Bal	Lab No.	: MB/2023/212	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 12:11
Patient Name	: anonymous	Age/Sex	: 53 Yr/M	Acceptance Date : 09-Jan-2023 16:44
Sample Type	: Fluid	Lab No.	: MB-2023-146	Reporting Date : 09-Jan-2023 16:50
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Ascitic fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:35
Patient Name	: anonymous	Age/Sex	: 20 Yr/M	Acceptance Date : 06-Jan-2023 18:59
Sample Type	: Peritoneal Fluid	Lab No.	: MB/2023/132	Reporting Date : 09-Jan-2023 14:10
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 05:54
Patient Name	: anonymous	Age/Sex	: 25 Yr/M	Acceptance Date : 03-Jan-2023 16:46
Sample Type	: Tt	Lab No.	: MB/2023/48	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 11:01
Patient Name	: anonymous	Age/Sex	: 47 Yr/F	Acceptance Date : 06-Jan-2023 19:03
Sample Type	: Sputum	Lab No.	: MB/2023/126	Reporting Date : 09-Jan-2023 14:10
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:16
Patient Name	: anonymous	Age/Sex	: 28 Yr/F	Acceptance Date : 06-Jan-2023 19:02
Sample Type	: Sputum	Lab No.	: MB/2023/127	Reporting Date : 09-Jan-2023 14:10
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 08:38
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 11-Jan-2023 17:34
Sample Type	: Fluid	Lab No.	: MB/2023/213	Reporting Date : 12-Jan-2023 17:08
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens Ascitic fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 10:13
Patient Name	: anonymous	Age/Sex	: 24 Yr/F	Acceptance Date : 11-Jan-2023 17:19
Sample Type	: Biopsy	Lab No.	: MB/2023/219	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis	NOT DETECTED
Rifampicin Resistance	NOT APPLICABLE
Test Performance	VALID
Remarks	specimen- tissue from neck region

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 07:57
Patient Name	: anonymous	Age/Sex	: 72 Yr/F	Acceptance Date : 11-Jan-2023 17:31
Sample Type	: Sputum	Lab No.	: MB/2023/217	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:45
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 06-Jan-2023 18:58
Sample Type	: Pleural Fluid	Lab No.	: MB/2023/134	Reporting Date : 09-Jan-2023 14:10
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 10:44
Patient Name	: anonymous	Age/Sex	: 11 Yr/F	Acceptance Date : 11-Jan-2023 17:17
Sample Type	: Sputum	Lab No.	: MB/2023/223	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 12:05
Patient Name	: anonymous	Age/Sex	: 39 Yr/F	Acceptance Date : 03-Jan-2023 16:44
Sample Type	: Bal	Lab No.	: MB/2023/51	Reporting Date : 23-Feb-2023 13:11
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 21:22
Patient Name	: anonymous	Age/Sex	: 51 Yr/M	Acceptance Date : 06-Jan-2023 19:15
Sample Type	: Sputum	Lab No.	: MB-2023-121	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 11:01
Patient Name	: anonymous	Age/Sex	: 47 Yr/F	Acceptance Date : 06-Jan-2023 19:03
Sample Type	: Sputum	Lab No.	: MB/2023/125	Reporting Date : 28-Feb-2023 15:19
Referred Doctor	:	Patient Type	: OPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

Status :1 Amendment/Addendum Report

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 12:19
Patient Name	: anonymous	Age/Sex	: 27 Yr/F	Acceptance Date : 02-Jan-2023 18:37
Sample Type	: Urine	Lab No.	: MB/2023/07	Reporting Date : 07-Jan-2023 12:45
Referred Doctor	:	Patient Type	: OPD	Addendum Date : 07-jan-2023 11:11

ACID FAST STAIN REPORT

Specimens URINE

ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous Result Modified By: Hemant Kashyap

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:46
Patient Name	: anonymous	Age/Sex	: 45 Yr/F	Acceptance Date : 11-Jan-2023 09:56
Sample Type	: Bal	Lab No.	: MB /2023/208	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis DETECTED (LOW)

Rifampicin Resistance NOT DETECTED

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:56
Patient Name	: anonymous	Age/Sex	: 35 Yr/M	Acceptance Date : 11-Jan-2023 09:54
Sample Type	: Fluid	Lab No.	: MB /2023/209	Reporting Date : 11-Jan-2023 17:05
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens Pleural fluid
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:46
Patient Name	: anonymous	Age/Sex	: 45 Yr/F	Acceptance Date : 11-Jan-2023 09:56
Sample Type	: Bal	Lab No.	: MB /2023/207	Reporting Date : 28-Feb-2023 15:19
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 16:56
Patient Name	: anonymous	Age/Sex	: 41 Yr/M	Acceptance Date : 11-Jan-2023 10:29
Sample Type	: Aspirate	Lab No.	: MB /2023/190	Reporting Date : 13-Jan-2023 17:47
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis	NOT DETECTED
Rifampicin Resistance	NOT APPLICABLE
Test Performance	VALID
Remarks	sepcimen- pericordial fluid.

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 09:42
Patient Name	: anonymous	Age/Sex	: 47 Yr/M	Acceptance Date : 06-Jan-2023 19:06
Sample Type	: Sputum	Lab No.	: MB/2023/124	Reporting Date : 10-Jan-2023 15:56
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 12:12
Patient Name	: anonymous	Age/Sex	: 52 Yr/F	Acceptance Date : 02-Jan-2023 18:20
Sample Type	: Urine	Lab No.	: MB/2023/31	Reporting Date : 04-Jan-2023 10:30
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 09:37
Patient Name	: anonymous	Age/Sex	: 3 Yr/F	Acceptance Date : 10-Jan-2023 11:57
Sample Type	: Sputum	Lab No.	: MB-2023-157	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 12:05
Patient Name	: anonymous	Age/Sex	: 39 Yr/F	Acceptance Date : 03-Jan-2023 16:44
Sample Type	: Bal	Lab No.	: MB/2023/51	Reporting Date : 06-Jan-2023 10:35
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 13:39
Patient Name	: anonymous	Age/Sex	: 56 Yr/F	Acceptance Date : 05-Jan-2023 18:59
Sample Type	: Sputum	Lab No.	: MB/2023/108	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:46
Patient Name	: anonymous	Age/Sex	: 45 Yr/F	Acceptance Date : 11-Jan-2023 09:56
Sample Type	: Bal	Lab No.	: MB /2023/207	Reporting Date : 11-Jan-2023 17:05
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 19:25
Patient Name	: anonymous	Age/Sex	: 7 Yr/M	Acceptance Date : 11-Jan-2023 10:26
Sample Type	: Bal	Lab No.	: MB /2023/191	Reporting Date : 12-Jan-2023 17:08
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens BAL
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:35
Patient Name	: anonymous	Age/Sex	: 20 Yr/M	Acceptance Date : 06-Jan-2023 18:59
Sample Type	: Pus	Lab No.	: MB/2023/131	Reporting Date : 09-Jan-2023 16:50
Referred Doctor	:	Patient Type	: OPD	

ACID FAST STAIN REPORT

Specimens PUS
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 10:55
Patient Name	: anonymous	Age/Sex	: 63 Yr/M	Acceptance Date : 11-Jan-2023 17:17
Sample Type	: Sputum	Lab No.	: MB/2023/222	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 14:04
Patient Name	: anonymous	Age/Sex	: 16 Yr/F	Acceptance Date : 05-Jan-2023 18:58
Sample Type	: Sputum	Lab No.	: MB/2023/107	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 06:57
Patient Name	: anonymous	Age/Sex	: 31 Yr/M	Acceptance Date : 04-Jan-2023 19:35
Sample Type	: Sputum	Lab No.	: MB/2023/90	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:11
Patient Name	: anonymous	Age/Sex	: 12 Yr/M	Acceptance Date : 06-Jan-2023 19:00
Sample Type	: Sputum	Lab No.	: MB/2023/129	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 08:00
Patient Name	: anonymous	Age/Sex	: 8 Yr/F	Acceptance Date : 10-Jan-2023 11:57
Sample Type	: Aspirate	Lab No.	: MB-2023-155	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis	NOT DETECTED
Rifampicin Resistance	NOT APPLICABLE
Test Performance	VALID
Remarks	SPECIMEN- GASTIRC ASPIRATE.

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 13:29
Patient Name	: anonymous	Age/Sex	: 70 Yr/F	Acceptance Date : 10-Jan-2023 11:53
Sample Type	: Sputum	Lab No.	: MB-2023-164	Reporting Date : 10-Jan-2023 15:02
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 11:57
Patient Name	: anonymous	Age/Sex	: 29 Yr/F	Acceptance Date : 02-Jan-2023 18:23
Sample Type	: Sputum	Lab No.	: MB/2023/24	Reporting Date : 07-Jan-2023 14:02
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 11:58
Patient Name	: anonymous	Age/Sex	: 26 Yr/M	Acceptance Date : 02-Jan-2023 18:23
Sample Type	: Sputum	Lab No.	: MB/2023/25	Reporting Date : 09-Jan-2023 10:56
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis

Rifampicin Resistance

Test Performance INVALID

Remarks Please send repeat sample.

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 08:38
Patient Name	: anonymous	Age/Sex	: 50 Yr/M	Acceptance Date : 11-Jan-2023 17:33
Sample Type	: Ascitic Fluid	Lab No.	: MB/2023/214	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 15:44
Patient Name	: anonymous	Age/Sex	: 65 Yr/M	Acceptance Date : 05-Jan-2023 19:00
Sample Type	: Biopsy	Lab No.	: MB/2023/114	Reporting Date : 07-Jan-2023 14:12
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 10-Jan-2023 14:56
Patient Name	: anonymous	Age/Sex	: 35 Yr/M	Acceptance Date : 24-Feb-2023 15:37
Sample Type	: Pleural Fluid	Lab No.	: mb-2023-209	Reporting Date : 28-Feb-2023 15:19
Referred Doctor	:	Patient Type	: IPD	

RAPID LIQUID MYCOBACTERIAL CULTURE AND FIRST LINE DST BY LPA

ZN Stain	Negative for Acid Fast Bacilli
Culture	Culture Negative for Mycobacteria species after 42 Days of aerobic incubation.
Drug Susceptibility to first line Drugs	Rifampicin- Isoniazid-

Comments

- **Test method for culture** - Liquid culture for Mycobacteria by MGIT 960 automated system (BD diagnostics)
- **Test method for DST** - Line probe assay using Genotype MTBDRplusver 2.0 (Hain lifesciences)
- **Notes:** LPA performed using MTBDRplusver 2.0 kit can only detect resistance conferred by MTBC to Rifampicin and Isoniazid targeting mutations originating in *rpoB*, *katG* and *inhA* regions. Resistances originating from mutations of other genes or gene regions as well as resistances due to other resistance mechanisms are not detected by this test.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 14:31
Patient Name	: anonymous	Age/Sex	: 54 Yr/M	Acceptance Date : 14-Jan-2023 12:23
Sample Type	: Sputum	Lab No.	: MB-2023-156	Reporting Date : 14-Jan-2023 14:18
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 11-Jan-2023 10:17
Patient Name	: anonymous	Age/Sex	: 13 Yr/M	Acceptance Date : 11-Jan-2023 17:19
Sample Type	: Sputum	Lab No.	: MB/2023/220	Reporting Date : 16-Jan-2023 17:27
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 00:18
Patient Name	: anonymous	Age/Sex	: 68 Yr/M	Acceptance Date : 03-Jan-2023 16:48
Sample Type	: Et	Lab No.	: MB/2023/45	Reporting Date : 05-Jan-2023 13:30
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 09:22
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 06-Jan-2023 19:18
Sample Type	: Pus	Lab No.	: MB/2023/118	Reporting Date : 09-Jan-2023 14:36
Referred Doctor	:	Patient Type	: IPD	

ACID FAST STAIN REPORT

Specimens PUS
ZN Stain Negative for Acid Fast Bacilli

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 12:06
Patient Name	: anonymous	Age/Sex	: 59 Yr/M	Acceptance Date : 03-Jan-2023 16:26
Sample Type	: Sputum	Lab No.	: MB/2023/71	Reporting Date : 09-Jan-2023 14:04
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 06-Jan-2023 12:46
Patient Name	: anonymous	Age/Sex	: 27 Yr/M	Acceptance Date : 06-Jan-2023 19:00
Sample Type	: Pus	Lab No.	: MB/2023/130	Reporting Date : 09-Jan-2023 14:10
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 03-Jan-2023 11:37
Patient Name	: anonymous	Age/Sex	: 50 Yr/F	Acceptance Date : 03-Jan-2023 16:41
Sample Type	: Aspirate	Lab No.	: MB/2023/58	Reporting Date : 05-Jan-2023 13:30
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 13:05
Patient Name	: anonymous	Age/Sex	: 72 Yr/M	Acceptance Date : 02-Jan-2023 18:14
Sample Type	: Sputum	Lab No.	: MB/2023/36	Reporting Date : 04-Jan-2023 10:30
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in *rpoB* gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 07-Jan-2023 12:11
Patient Name	: anonymous	Age/Sex	: 53 Yr/M	Acceptance Date : 11-Jan-2023 11:09
Sample Type	: Ascitic Fluid	Lab No.	: MB-2023-147	Reporting Date : 11-Jan-2023 13:05
Referred Doctor	:	Patient Type	: OPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Microbiology

Mycobacteriology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 05-Jan-2023 10:00
Patient Name	: anonymous	Age/Sex	: 57 Yr/M	Acceptance Date : 05-Jan-2023 18:56
Sample Type	: Sputum	Lab No.	: MB/2023/103	Reporting Date : 07-Jan-2023 14:12
Referred Doctor	:	Patient Type	: IPD	

CBNAAT (GENEXPERT) FOR MYCOBACTERIUM TUBERCULOSIS

Mycobacterium Tuberculosis NOT DETECTED

Rifampicin Resistance NOT APPLICABLE

Test Performance VALID

Remarks

- **Test method** - Cartridge based nucleic acid amplification test (CBNAAT) performed using Xpert MTB/RIF assay (Cepheid)
- **Test interpretation-**
 1. GeneXpert is a real time nested PCR for detection of Mycobacterium tuberculosis from clinical specimens. It simultaneously detects rifampicin resistance associated with mutations in rpoB gene only. Resistance due to other mechanisms or mutations at other gene regions are not detected by this assay.
 2. Indeterminate results for Rifampicin mean resistance could not be determined due to insufficient signal as a result of low concentration of MTB. In this case, the sample should be sent for culture based drug sensitivity testing.
 3. Positive test result does not necessarily indicate the presence of viable organisms as it targets bacterial DNA. Hence, test results should not be used to determine therapeutic success/failure as the DNA might persist following antimicrobial therapy.

***** END OF THE REPORT *****

All reports need clinical correlation. Kindly discuss if necessary. No part of the report can be reproduced without written permission of the department.

Validated By: anonymous

This is computer generated report. Signature not required