



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:16
Patient Name	: anonymous	Age/Sex	: 64 Yr/F	Acceptance Date : 01-Jan-2023 11:12
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT011	Reporting Date : 10-Jan-2023 15:49
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	17.8	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.8	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:20
Patient Name	: anonymous	Age/Sex	: 26 Yr/M	Acceptance Date : 01-Jan-2023 11:13
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT013	Reporting Date : 10-Jan-2023 15:49
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	15.4	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.6	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:08
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 01-Jan-2023 19:51
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT030	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: OPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	11.1	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.1	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 18:19
Patient Name	: anonymous	Age/Sex	: 58 Yr/M	Acceptance Date : 02-Jan-2023 00:05
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP029	Reporting Date : 02-Jan-2023 02:04
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	29.0	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 19:02
Patient Name	: anonymous	Age/Sex	: 22 Yr/M	Acceptance Date : 02-Jan-2023 02:17
Sample Type	: Edta Blood	Lab No.	: 230102CB038	Reporting Date : 10-Jan-2023 15:34
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	15.2	13 - 17 g/dl
HCT	51.6	40 - 50 %
RBC	6.94	4.5 - 5.5 $\times 10^6$ / μ L
MCV	74.4	83 - 101 fL
MCH	21.9	27 - 32 pg
MCHC	29.5	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	15.0	11.6 - 14.0 %
Total Leukocyte Count(TLC)	6.34	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	44.7	40 - 80 %
Lymphocytes	47.2	20 - 40 %
Monocytes	4.7	2 - 10 %
Eosinophils	2.8	01 - 06 %
Basophil	0.6	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	2.83	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	2.99	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.30	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.18	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.04	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	284	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:08
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 01-Jan-2023 19:47
Sample Type	: Edta Blood	Lab No.	: 230101CB120	Reporting Date : 10-Jan-2023 16:12
Referred Doctor	:	Patient Type	: OPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	8.8	12 - 15 g/dl
HCT	27.0	36 - 46 %
RBC	2.98	3.8 - 4.8 $\times 10^6$ / μ L
MCV	90.6	83 - 101 fL
MCH	29.5	27 - 32 pg
MCHC	32.6	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	16.3	11.6 - 14.0 %
Total Leukocyte Count(TLC)	7.03	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	72.6	40 - 80 %
Lymphocytes	21.5	20 - 40 %
Monocytes	5.8	2 - 10 %
Eosinophils	0.1	01 - 06 %
Basophil	0.0	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	5.10	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	1.51	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.41	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.01	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.00	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	161	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:37
Patient Name	: anonymous	Age/Sex	: 31 Yr/F	Acceptance Date : 01-Jan-2023 19:52
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT032	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	8.4	09.6 - 12.2 sec
International Normalised Ratio (INR)	0.9	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	47 Yr/F
Sample Type	:	Edta Blood	Lab No.	:	230101CB059
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	01-Jan-2023 00:14
			Acceptance Date	:	01-Jan-2023 10:10
			Reporting Date	:	10-Jan-2023 16:03

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	11.7	12 - 15 g/dl
HCT	38.4	36 - 46 %
RBC	4.45	3.8 - 4.8 $\times 10^6$ / μ L
MCV	86.3	83 - 101 fL
MCH	26.3	27 - 32 pg
MCHC	30.5	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	20.5	11.6 - 14.0 %
Total Leukocyte Count(TLC)	15.30	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	75.0	40 - 80 %
Lymphocytes	16.9	20 - 40 %
Monocytes	6.8	2 - 10 %
Eosinophils	1.2	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	11.46	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	2.59	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	1.04	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.19	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.02	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	519	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:44
Patient Name	: anonymous	Age/Sex	: 67 Yr/F	Acceptance Date : 02-Jan-2023 00:05
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102PT001	Reporting Date : 02-Jan-2023 02:03
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	23.1	09.6 - 12.2 sec
International Normalised Ratio (INR)	2.4	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 17:08
Patient Name	: anonymous	Age/Sex	: 19 Yr/F	Acceptance Date : 01-Jan-2023 19:51
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP024	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: OPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	45.5	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	31 Yr/F
Sample Type	:	Edta Blood	Lab No.	:	230101CB116
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	01-Jan-2023 13:37
			Acceptance Date	:	01-Jan-2023 17:02
			Reporting Date	:	10-Jan-2023 16:15

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	15.6	12 - 15 g/dl
HCT	50.8	36 - 46 %
RBC	6.01	3.8 - 4.8 x10 ⁶ /μL
MCV	84.5	83 - 101 fL
MCH	26.0	27 - 32 pg
MCHC	30.7	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	23.9	11.6 - 14.0 %
Total Leukocyte Count(TLC)	9.03	4 - 10 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	82.2	40 - 80 %
Lymphocytes	9.9	20 - 40 %
Monocytes	7.8	2 - 10 %
Eosinophils	0.0	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	7.43	2 - 7 x10 ³ /μL
Absolute Lymphocyte Count	0.89	1 - 3 x10 ³ /μL
Absolute Monocyte Count	0.70	0.2 - 1.0 x10 ³ /μL
Absolute Eosinophil Count	0.00	0.02 - 0.5 x10 ³ /μL
Absolute Basophil Count	0.01	0.02 - 0.10 x10 ³ /μL
Platelet Count	134	150 - 410 x10 ³ /μL

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 54 Yr/M	Acceptance Date : 02-Jan-2023 02:14
Sample Type	: Edta Blood	Lab No.	: 230102CB024	Reporting Date : 10-Jan-2023 15:37
Referred Doctor	:	Patient Type	: OPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	6.6	13 - 17 g/dl
HCT	20.7	40 - 50 %
RBC	2.48	4.5 - 5.5 $\times 10^6$ / μ L
MCV	83.5	83 - 101 fL
MCH	26.6	27 - 32 pg
MCHC	31.9	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	14.7	11.6 - 14.0 %
Total Leukocyte Count(TLC)	22.00	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	94.3	40 - 80 %
Lymphocytes	1.2	20 - 40 %
Monocytes	4.4	2 - 10 %
Eosinophils	0.0	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	20.74	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	0.27	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.97	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.00	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.02	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	184	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:37
Patient Name	: anonymous	Age/Sex	: 31 Yr/F	Acceptance Date : 01-Jan-2023 19:52
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101AP026	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	21.8	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 18:19
Patient Name	: anonymous	Age/Sex	: 58 Yr/M	Acceptance Date : 02-Jan-2023 00:04
Sample Type	: Edta Blood	Lab No.	: 230102CB005	Reporting Date : 10-Jan-2023 15:34
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	11.2	13 - 17 g/dl
HCT	37.0	40 - 50 %
RBC	4.71	4.5 - 5.5 $\times 10^6$ / μ L
MCV	78.6	83 - 101 fL
MCH	23.8	27 - 32 pg
MCHC	30.3	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	15.1	11.6 - 14.0 %
Total Leukocyte Count(TLC)	14.11	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	66.1	40 - 80 %
Lymphocytes	18.7	20 - 40 %
Monocytes	10.2	2 - 10 %
Eosinophils	4.9	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	9.33	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	2.64	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	1.44	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.69	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.01	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	233	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 54 Yr/M	Acceptance Date : 02-Jan-2023 02:22
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102AP013	Reporting Date : 10-Jan-2023 15:48
Referred Doctor	:	Patient Type	: OPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	74.2	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	2 D/F
Sample Type	:	Edta Blood	Lab No.	:	230101CB130
Referred Doctor	:		Patient Type	:	IPD
			Collection Date	:	01-Jan-2023 13:40
			Acceptance Date	:	01-Jan-2023 19:49
			Reporting Date	:	05-Jan-2023 23:42

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	19.7	14 - 22 g/dl
HCT	58.6	45 - 75 %
RBC	6.04	5 - 7 x10 ⁶ /μL
MCV	97.0	92 - 120 fL
MCH	32.6	31 - 37 pg
MCHC	33.6	30 - 36 g/dL
Red Cell Distribution Width (RDW CV)	23.6	11.6 - 14.0 %
Total Leukocyte Count(TLC)	17.76	10 - 26 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	61.2	41 - 81 %
Lymphocytes	31.6	21 - 41 %
Monocytes	6.3	2 - 10 %
Eosinophils	0.3	01 - 06 %
Basophil	0.6	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	10.88	3 - 8 x10 ³ /μL
Absolute Lymphocyte Count	5.62	3 - 8 x10 ³ /μL
Absolute Monocyte Count	1.11	0.5 - 2 x10 ³ /μL
Absolute Eosinophil Count	0.05	0.1 - 1 x10 ³ /μL
Absolute Basophil Count	0.10	0.02 - 0.10 x10 ³ /μL
Platelet Count	131	100 - 450 x10 ³ /μL

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:49
Patient Name	: anonymous	Age/Sex	: 60 Yr/M	Acceptance Date : 01-Jan-2023 02:13
Sample Type	: Edta Blood	Lab No.	: 230101CB039	Reporting Date : 05-Jan-2023 23:42
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	11.1	13 - 17 g/dl
HCT	33.1	40 - 50 %
RBC	4.10	4.5 - 5.5 x10 ⁶ /μL
MCV	80.7	83 - 101 fL
MCH	27.1	27 - 32 pg
MCHC	33.5	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	15.8	11.6 - 14.0 %
Total Leukocyte Count(TLC)	13.42	4 - 10 x10 ³ /μL
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	80.8	40 - 80 %
Lymphocytes	13.3	20 - 40 %
Monocytes	5.4	2 - 10 %
Eosinophils	0.3	01 - 06 %
Basophil	0.2	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	10.83	2 - 7 x10 ³ /μL
Absolute Lymphocyte Count	1.79	1 - 3 x10 ³ /μL
Absolute Monocyte Count	0.73	0.2 - 1.0 x10 ³ /μL
Absolute Eosinophil Count	0.04	0.02 - 0.5 x10 ³ /μL
Absolute Basophil Count	0.03	0.02 - 0.10 x10 ³ /μL
Platelet Count	277	150 - 410 x10 ³ /μL

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:54
Patient Name	: anonymous	Age/Sex	: 1 D/F	Acceptance Date : 01-Jan-2023 01:19
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT007	Reporting Date : 10-Jan-2023 15:49
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	18.5	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.9	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:26
Patient Name	: anonymous	Age/Sex	: 1 Yr/M	Acceptance Date : 01-Jan-2023 02:37
Sample Type	: Edta Blood	Lab No.	: 230101CB042	Reporting Date : 10-Jan-2023 16:19
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	5.6	11.1 - 14.1 g/dl
HCT	19.8	30 - 40 %
RBC	2.48	3.9 - 5.3 $\times 10^6$ / μ L
MCV	79.8	68 - 84 fL
MCH	22.6	24 - 30 pg
MCHC	28.3	30 - 36 g/dL
Red Cell Distribution Width (RDW CV)	24.5	30 - 36 %
Total Leukocyte Count(TLC)	6.12	6 - 18 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	65.8	15 - 45 %
Lymphocytes	31.9	47 - 77 %
Monocytes	2.3	2 - 10 %
Eosinophils	0.0	01 - 06 %
Basophil	0.0	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	4.03	1 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	1.95	3.5 - 11 $\times 10^3$ / μ L
Absolute Monocyte Count	0.14	0.1 - 1.2 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.00	0.1 - 1 $\times 10^3$ / μ L
Absolute Basophil Count	0.00	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	7	200 - 550 $\times 10^3$ / μ L

Comments on PBS : Platelets appear severely depleted on peripheral smear examination.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 18:19
Patient Name	: anonymous	Age/Sex	: 58 Yr/M	Acceptance Date : 02-Jan-2023 00:05
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102PT001	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	12.0	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.2	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 13:44
Patient Name	: anonymous	Age/Sex	: 67 Yr/F	Acceptance Date : 02-Jan-2023 00:05
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102AP001	Reporting Date : 02-Jan-2023 02:04
Referred Doctor	:	Patient Type	: IPD	

Activated Partial Thromboplastin Time (APTT)

SPECIMEN: 3.2 % citrated plasma **METHOD:** Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Activated Partial Thromboplastin Time (APTT)	57.9	23.2 - 32.4 sec

Comments:

- 1.Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
- 2.Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
- 3.APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	:	-	ABHA ADDRESS	:	-
CR No	:	-	Sample No.	:	-
Patient Name	:	anonymous	Age/Sex	:	43 Yr/M
Sample Type	:	Edta Blood	Lab No.	:	230101CB109
Referred Doctor	:		Patient Type	:	OPD
			Collection Date	:	01-Jan-2023 16:26
			Acceptance Date	:	01-Jan-2023 17:01
			Reporting Date	:	02-Jan-2023 07:58

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	10.7	13 - 17 g/dl
HCT	35.0	40 - 50 %
RBC	3.61	4.5 - 5.5 $\times 10^6$ / μ L
MCV	97.0	83 - 101 fL
MCH	29.6	27 - 32 pg
MCHC	30.6	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	14.4	11.6 - 14.0 %
Total Leukocyte Count(TLC)	6.90	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	83	40 - 80 %
Lymphocytes	7	20 - 40 %
Monocytes	10	2 - 10 %
Eosinophils		01 - 06 %
Basophil		00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count		2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count		1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count		0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count		0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count		0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	105	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:14
Patient Name	: anonymous	Age/Sex	: 47 Yr/F	Acceptance Date : 01-Jan-2023 11:11
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230101PT010	Reporting Date : 10-Jan-2023 15:49
Referred Doctor	:	Patient Type	: IPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	22.7	09.6 - 12.2 sec
International Normalised Ratio (INR)	2.3	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 01-Jan-2023 00:16
Patient Name	: anonymous	Age/Sex	: 64 Yr/F	Acceptance Date : 01-Jan-2023 10:10
Sample Type	: Edta Blood	Lab No.	: 230101CB057	Reporting Date : 10-Jan-2023 16:03
Referred Doctor	:	Patient Type	: IPD	

Complete Blood Count (CBC)/ Hemogram

SPECIMEN: EDTA anticoagulated whole blood **METHOD:** Electrical impedance and flow cytometry

Test	Result	Ref. Range and Unit
Hemoglobin	7.1	12 - 15 g/dl
HCT	22.8	36 - 46 %
RBC	2.44	3.8 - 4.8 $\times 10^6$ / μ L
MCV	93.4	83 - 101 fL
MCH	29.1	27 - 32 pg
MCHC	31.1	31.5 - 34.5 g/dL
Red Cell Distribution Width (RDW CV)	21.1	11.6 - 14.0 %
Total Leukocyte Count(TLC)	8.64	4 - 10 $\times 10^3$ / μ L
DIFFERENTIAL LEUKOCYTE COUNT (DLC)		
Neutrophils	73.9	40 - 80 %
Lymphocytes	16.0	20 - 40 %
Monocytes	7.2	2 - 10 %
Eosinophils	2.8	01 - 06 %
Basophil	0.1	00 - 01 %
ABSOLUTE LEUKOCYTE COUNT		
Absolute Neutrophil Count	6.39	2 - 7 $\times 10^3$ / μ L
Absolute Lymphocyte Count	1.38	1 - 3 $\times 10^3$ / μ L
Absolute Monocyte Count	0.62	0.2 - 1.0 $\times 10^3$ / μ L
Absolute Eosinophil Count	0.24	0.02 - 0.5 $\times 10^3$ / μ L
Absolute Basophil Count	0.01	0.02 - 0.10 $\times 10^3$ / μ L
Platelet Count	345	150 - 410 $\times 10^3$ / μ L

Comments on PBS :

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required



C-DAC: Centre for Development of Advanced Computing
C-56/1, Sector-62, Noida - 201307, Uttar Pradesh,
91-0120-3063311-13.



Department of Pathology And Lab Medicine

General Hematology

ABHA NO	: -	ABHA ADDRESS	: -	
CR No	: -	Sample No.	: -	Collection Date : 02-Jan-2023 01:51
Patient Name	: anonymous	Age/Sex	: 54 Yr/M	Acceptance Date : 02-Jan-2023 02:22
Sample Type	: 3.2 Citrated Plasma	Lab No.	: 230102PT008	Reporting Date : 10-Jan-2023 15:50
Referred Doctor	:	Patient Type	: OPD	

Prothrombin Time (PT-INR)

SPECIMEN: 3.2 % citrated plasma

METHOD: Coagulation, optical detection method

CLINICAL DETAILS:

<u>Test</u>	<u>Result</u>	<u>Ref. Range and Unit</u>
Prothrombin time (PT)	14.0	09.6 - 12.2 sec
International Normalised Ratio (INR)	1.4	

Comments:

1. Prothrombin time is measure of the integrity of the extrinsic and common pathways of the coagulation cascade. PT is used for monitoring oral anticoagulation therapy, evaluation of liver function and coagulation disorders especially involving extrinsic pathway.
2. Recommended therapeutic range: **INR 2.0-3.0** - treatment of venous thrombosis and pulmonary embolism, prevention of systemic embolism, hypercoagulable state, atrial fibrillation, etc. **INR 2.5-3.5** - mechanical prosthetic valves, systemic recurrent emboli etc.
3. APTT is measure of the intrinsic coagulation pathway as well as common pathway. APTT is widely used for monitoring unfractionated heparin therapy, however it is affected by number of variables not related to heparin.

***** END OF THE REPORT *****

Validated By: anonymous

This is computer generated report. Signature not required