

05 Hr 58 Min
13 Sec

Guidelines

Coding Area

Editor | Compile &
Run History

Submissions

Feedback Form

Result

Dashboard

Graphs

Coding Area

A

B

C

D

E

F

ONLINE EDITOR (D)

Bad Permutations

+ Problem Description

Mr. X is teaching number theory in his class. He is discussing about factors and permutations in his class. A factor of a positive integer N is a positive integer that divides n precisely (without leaving a remainder). The set of factors always includes 1 and N.

Mr. X likes combinatorics a lot. He asked his students find out all the factors of the number Y, and sort them in an ascending order. He asks them to list all permutations of the factors. They then need to cross out all permutations where two adjacent numbers are adjacent in the same order in the original list. The number of uncrossed (valid) permutations are to be given to him.

Illustration:

Integer 9 has 3 factors [1,3,9].

The permutations of these factors of number 9 are [1,3,9],[1,9,3],[3,9,1],[3,1,9],[9,1,3],[9,3,1].

Of these 6 permutations, we need to cross out [1,3,9] (1 3 adjacent in same order), [3,9,1] (3 9 in same order) and [9,1,3] (1 3 in same order)

The remaining (valid) permutations are:

[1,9,3] ,[9,3,1],[3,1,9]

Hence the number of valid permutations =3, which is the answer.

+ Constraints

$1 \leq N \leq 120000$

$1 \leq T \leq 100$

+ Input Format

The first line contains T, the number of testcases

Next T lines contain the integer N

+ Output

T lines containing number of valid permutationssatisfying conditions mentioned in the problem statement for given input.

+ Test Case

+ Explanation

Example 1

Input

1

10

Output

11

Explanation

T=1 (there is 1 test case)

N=10.

10 has 4 factors [1,2,5,10]. There are 24 permutations of these four factors. The 11 valid permutations are [1,5,2,10],[1,10,5,2], [2,1,10,5],[2,10,1,5], [2,10,5,1], [5,1,10,2], [5,2,1,10], [5,2,10,1], [10,1,5,2], [10,2,1,5], [10,5,2,1]. Hence the output is 11

Example 2

Input

2

6

9

Output

11

3

Explanation

T=2 (there are 2 test cases).

In the first test case, N=6. 6 has four factors [1,2,3,6]. As in the previous example, there are 11 valid permutations for these. Hence the output for the first test case is 11. This is the first line of the output

In the second test case, N=9. As was shown in the Illustration in the problem statement, the number of valid permutations is 3. Hence the output for the second test case (the second line of the output) is 3.

Upload Solution [Question : D]

☐ I, **chukka bharath** confirm that the answer submitted is my own.

☐ Took help from online sources (attributions)

Choose a File

...

[CodeVita FAQ's](#)

[About CodeVita](#)

[Privacy Policy](#)

[Careers](#)



© 2018 Tata Consultancy Services Limited. All Rights Reserved.