Ex.No-1 NUMPY

AIM:

To calculate the values for the mathematical formulas using NumPylibrary

INTEGRATEDDEVELOPMENTENVIRONMENT(IDE)REQUIRED:

JUPYTERNOTEBOOK

REQUIREDLIBRARIESFORPYTHON:

Numpy

PROCEDURE:

A) Euclideandistance

The mathematical formula for calculating the Euclidean distance between 2 points in 2D space:

$$d(p,q) = \sqrt[2]{(q_1-p_1)^2+(q_2-p_2)^2}$$

B) Dot Product

$$u=\begin{bmatrix}5\\12\end{bmatrix},\quad v=\begin{bmatrix}8\\6\end{bmatrix}$$
 Dot product is
$$u\cdot v=u_1\times v_1+u_2\times v_2\\ =5\times 8+12\times 6\\ =112$$

C)SolvingaSystemofLinearEquations

A system of linear equations can be represented in matrix form as AX=B, where A is the matrix of coefficients, X is the column vector of variables, and B is the column vector of solutions. To solve for X, we can use: $X=A^{-1}$ Bassuming Ais invertible.

PROGRAM:

$\underline{A) Calculating the Euclidean Distance Between Two Points}$

importnumpy as

npdefeuclidean_distance(p,q):

```
returnnp.sqrt(np.sum((q-p)**2))#
Exampleusage
p = np.array([1,2])
q = np.array([4,6])
distance=euclidean_distance(p,q)
print("OutputforCalculatingtheEuclideanDistanceBetweenTwoPointsis:",distance)
\underline{B)} \underline{Calculating the Dot Product of Two Vectors}
importnumpyasnp
A=np.array([1,3,-5])
B = np.array([4, -2, -
1])dot_product=np.dot(A,B)
print("OutputfordotproductoftwovectorsAandBis",dot product)
<u>C)SolvingaSystemofLinearEquations</u>
importnumpyasnp
#CoefficientsmatrixAandresultvectorbA=
np.array([[3, 1], [1,2]])
b=np.array([9,8])#
Solvefor x
x=np.linalg.solve(A,b)
print("OutputsolutionofSystemofLinearEquationsis",x)
Output:
A)OutputforCalculatingtheEuclideanDistancebetweenTwoPointsis:5.0.Exercise2-B)Output fordot
productof two vectorsAand Bis3
C)OutputsolutionofSystemofLinearEquationsis[2.3.]
Result:
```

Theprograms wererunsuccessfully