



SRM Institute of Science and Technology
Ramapuram Campus
Department of Mathematics
18MAB101T - Calculus And Linear Algebra

Year/Sem: I/I

Branch: Common to ALL B.Tech. except B.Tech. (Business Systems)

Unit – IV

DIFFERENTIAL CALCULUS

Part – C

1. Find the radius of curvature of the curve $y^2 = 12x$ at the point (3, 6).

Solution:

$$\frac{dy}{dx} = \frac{6}{y}$$

$$\text{At (3, 6), } \frac{dy}{dx} = 1$$

$$\frac{d^2y}{dx^2} = 6 \left(\frac{-1}{y^2} \right) \frac{dy}{dx}$$

$$\text{At (3, 6), } \frac{d^2y}{dx^2} = 6 \left(\frac{-1}{36} \right) 1 = \frac{-1}{6}$$

$$\rho = \frac{(1 + y_1^2)^{\frac{3}{2}}}{y_2}$$

$$\rho = \frac{(1+1)^{\frac{3}{2}}}{-1/6} = -12\sqrt{2}$$

2. Find the radius of curvature of the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ at the point (a/4, a/4).

Solution:

$$\frac{dy}{dx} = -\sqrt{\frac{y}{x}}$$

$$\text{At } (a/4, a/4), \frac{dy}{dx} = -1$$

$$\frac{d^2y}{dx^2} = -\left[\frac{\sqrt{x} \cdot \frac{1}{2} y^{-1/2} \frac{dy}{dx} - \sqrt{y} \cdot \frac{1}{2} x^{-1/2}}{x} \right]$$

$$\text{At } (a/4, a/4), \frac{d^2y}{dx^2} = \frac{4}{a}$$

$$\rho = \frac{(1 + y_1^2)^{\frac{3}{2}}}{y_2}$$

$$\rho = \frac{a}{\sqrt{2}}$$

3. Find the radius of curvature of the curve $xy = c^2$ at the point (c, c) .**Solution:**

$$x \cdot \frac{dy}{dx} + y \cdot 1 = 0$$

$$\frac{dy}{dx} = -\frac{y}{x}$$

$$\text{At } (c, c), \frac{dy}{dx} = -1$$

$$\frac{d^2y}{dx^2} = -\left[\frac{x \frac{dy}{dx} - y \cdot 1}{x^2} \right]$$

$$\text{At } (c, c), \frac{d^2y}{dx^2} = -\left(\frac{-2c}{c^2} \right) = \frac{2}{c}$$

$$\rho = \frac{(1 + y_1^2)^{\frac{3}{2}}}{y_2}$$

$$\rho = \frac{(1+1)^{\frac{3}{2}}}{2/c} = \sqrt{2} c$$

4. If $x = a \cos \theta$, $y = b \sin \theta$, then find $\frac{dy}{dx}$.

Solution:

$$\frac{dx}{d\theta} = -a \sin \theta, \frac{dy}{d\theta} = b \cos \theta, \frac{dy}{dx} = -\frac{b}{a} \cot \theta$$

5. Find the envelope of $x \cos \theta + y \sin \theta = 1$, θ being the parameter.

Solution:

$$x \cos \theta + y \sin \theta = 1 \quad \text{_____} (1)$$

Differentiate partially w.r.t. θ .

$$x(-\sin \theta) + y(\cos \theta) = 0 \quad \text{_____} (2)$$

Squaring and adding (1) and (2)

$$x^2 + y^2 = 1$$

6. Find the envelope of $x \cos \alpha + y \sin \alpha = a \sec \alpha$, α being the parameter.

Solution:

$$x \cos \alpha + y \sin \alpha = a \sec \alpha$$

Divide by $\cos \alpha$.

$$x + y \tan \alpha = a \sec^2 \alpha$$

$$x + y \tan \alpha = a (1 + \tan^2 \alpha)$$

$$a \tan^2 \alpha - y \tan \alpha + (a - x) = 0$$

$$\text{Here } A = a, B = -y, C = a - x$$

Envelope is given by $B^2 - 4AC = 0$.

$$y^2 = 4a(a - x)$$

7. Find $\int_0^1 x^6 (1-x)^9 dx$.

Solution:

$$m = 7, n = 10$$

$$B(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx$$

$$= \int_0^1 x^{7-1} (1-x)^{10-1} dx$$

$$= \frac{\Gamma(7)\Gamma(10)}{\Gamma(17)} = \frac{6!9!}{16!}$$

8. Prove that $\frac{B(m+1, n)}{B(m, n+1)} = \frac{m}{n}$.

Solution:

$$\frac{B(m+1, n)}{B(m, n+1)} = \frac{\frac{\Gamma(m+1)\Gamma(n)}{\Gamma(m+n+1)}}{\frac{\Gamma(m)\Gamma(n+1)}{\Gamma(m+n+1)}} = \frac{m\Gamma(m)\Gamma(n)}{n\Gamma(m)\Gamma(n)} = \frac{m}{n}$$

9. Find $\int_0^{\pi/2} \sqrt{\tan \theta} d\theta$.

Solution:

$$\begin{aligned} \int_0^{\pi/2} \sqrt{\tan \theta} d\theta &= \int_0^{\pi/2} \sqrt{\frac{\sin \theta}{\cos \theta}} d\theta = \int_0^{\pi/2} \sin^{1/2} \theta \cos^{-1/2} \theta d\theta \\ &= \frac{1}{2} B\left(\frac{3/2}{2}, \frac{1/2}{2}\right) \\ &= \frac{1}{2} B\left(\frac{3}{4}, \frac{1}{4}\right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \frac{\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{1}{4}\right)}{\Gamma(1)} \\
 &= \frac{\pi}{\sqrt{2}}
 \end{aligned}$$

Formula $\Gamma(n)\Gamma(1-n) = \frac{\pi}{\sin n\pi}$

10. Find $\int_0^{\pi/2} \sin^6 \theta \cos^6 \theta d\theta.$

Solution: m = 6, n = 6

$$\begin{aligned}
 \int_0^{\pi/2} \sin^6 \theta \cos^6 \theta d\theta &= \frac{1}{2} B\left(\frac{m+1}{2}, \frac{n+1}{2}\right) \\
 &= \frac{1}{2} B\left(\frac{7}{2}, \frac{7}{2}\right) \\
 &= \frac{1}{2} \frac{\Gamma\left(\frac{7}{2}\right)\Gamma\left(\frac{7}{2}\right)}{\Gamma(7)} \\
 &= \frac{1}{2} \frac{\left(\frac{15}{8}\sqrt{\pi}\right)\left(\frac{15}{8}\sqrt{\pi}\right)}{6!}
 \end{aligned}$$

11. Prove that $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$

Solution:

$$B(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta$$

Put m = 1/2, n = 1/2 .

$$B\left(\frac{1}{2}, \frac{1}{2}\right) = 2 \int_0^{\pi/2} d\theta$$

$$\frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}\right)}{\Gamma(1)} = 2 \cdot \frac{\pi}{2}$$

$$\left(\Gamma\left(\frac{1}{2}\right)\right)^2 = \pi$$

Hence $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}.$

* * * * *