
# Probability and Statistics: Lecture-35

Monsoon-2020

```
by Dr. Pawan Kumar (IIIT, Hyderabad)
on November 4, 2020
```



Example (Solved Example

**Example (Solved Example** 

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x, y \ge 0\\ 0 & \text{otherwise} \end{cases}$$

**Example (Solved Example** 

Let X, Y be jointly continuous RVs with joint PDF

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x,y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

• Are X and Y independent?

Example (Solved Example

Let X, Y be jointly continuous RVs with joint PDF

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x,y \ge 0 \\ 0 & \text{otherwise} \end{cases}$$

ullet Are  ${\it X}$  and  ${\it Y}$  independent?  ${\it \bullet}$  Find  ${\it E}[{\it Y} \,|\, {\it X} > 2]$ 

Let X. Y be jointly continuous RVs with joint PDF

$$f_{X,Y} = \begin{cases} 6e^{-(2x+3y)} & x, y \ge 0\\ 0 & \text{otherwise} \end{cases}$$

• Are X and Y independent? • Find E[Y | X > 2] • Find P(X > Y)



» Answer to previous problem...

32

Example (Solved example

Let  $\boldsymbol{X}$  be a continuous RV with PDF

Example (Solved example)

Let X be a continuous RV with PDF

$$f_{X}(x) = egin{cases} 2x & 0 \leq x \leq 1 \\ 0 & ext{otherwise} \end{cases}$$

Example (Solved example

Let X be a continuous RV with PDF

$$f_{\mathcal{X}}(\mathbf{x}) = \begin{cases} 2\mathbf{x} & 0 \le \mathbf{x} \le 1 \\ 0 & \text{otherwise} \end{cases}$$

We are also know that given X = x, the RV Y is uniformly distributed on [-x, x].


Let X be a continuous RV with PDF

$$f_{X}(x) = \begin{cases} 2x & 0 \le x \le 1 \\ 0 & \text{otherwise} \end{cases}$$

We are also know that given X = x, the RV Y is uniformly distributed on [-x, x].

• Find the joint PDF 
$$f_{XY}(x,y)$$
 • Find  $P_Y(y)$  • Find  $P(|Y| < X^3)$ 











Example (Solved example

Example (Solved example

$$f_{\mathcal{X},\mathcal{Y}}(\mathbf{x},\mathbf{y}) = egin{cases} 6\mathbf{x}\mathbf{y} & 0 \leq \mathbf{x} \leq 1, \ 0 \leq \mathbf{y} \leq \sqrt{\mathbf{x}} \\ 0 & \text{otherwise} \end{cases}$$

Example (Solved example

Let X, Y be two jointly continuous RVs with joint PDF

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

1. Plot  $R_{XY}$ 

#### Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot  $R_{XY}$
- 2. Find  $f_X(x)$  and  $f_Y(y)$

Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & & ext{otherwise} \end{cases}$$

- 1. Plot  $R_{XY}$
- 2. Find  $f_X(x)$  and  $f_Y(y)$
- 3. Are X and Y independent?

#### Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot  $R_{XY}$
- 2. Find  $f_X(x)$  and  $f_Y(y)$
- 3. Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y,  $f_{X|Y}(x \mid y)$

#### Example (Solved example

$$f_{X,Y}(x,y) = egin{cases} 6xy & 0 \leq x \leq 1, \ 0 \leq y \leq \sqrt{x} \\ 0 & ext{otherwise} \end{cases}$$

- 1. Plot  $R_{XY}$
- 2. Find  $f_X(x)$  and  $f_Y(y)$
- 3. Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y,  $f_{X|Y}(x \mid y)$
- 5. Find  $E[X | Y = y], 0 \le y \le 1$

#### Example (Solved example

$$f_{X,Y}(x,y) = \begin{cases} 6xy & 0 \le x \le 1, \ 0 \le y \le \sqrt{x} \\ 0 & \text{otherwise} \end{cases}$$

- ightharpoonup Plot  $R_{XY}$
- Find  $f_X(x)$  and  $f_Y(y)$
- Are X and Y independent?
- 4. Find the conditional PDF of X given Y = y,  $f_{X|Y}(x \mid y)$
- **5.** Find E[X | Y = y], 0 ≤ y ≤ 1
- 6 Find  $Var(X \mid Y = y)$  for  $0 \le y \le 1$

\*Answer to previous problem...

Frith: 
$$\int fxy(x)(x) dx$$

Frith:  $\int fxy(x)(x) dx$ 

Frith:  $\int fxy(x) dx$ 

Fr

» Answer to previous problem...

(a) 
$$\int x|y|(x|y) = \int x|y|(x|y)$$

$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$


$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$

$$= \int \frac{2}{3}x^{3}|y|^{2} = \frac{2}{1-y^{4}} \left[\frac{2}{3}\right]^{2}$$

$$= \int \frac{2}{3}x^{3} = \int \frac{2}{3}x^{3} =$$

» Answer to previous problem...

9/132



» Definition of Covariance...

Definition of Covariance

Let X and Y be two random variables.

» Definition of Covariance...

#### Definition of Covariance

Let *X* and *Y* be two random variables. The covariance between *X* and *Y* is defined as


#### » Definition of Covariance...

#### **Definition of Covariance**

Let X and Y be two random variables. The covariance between X and Y is defined as

$$Cov(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - (E[X])(E[Y])$$

#### Derivation



Solved Example

Example (Solved Example)

Suppose 
$$X \sim \text{Uniform}(1,2)$$
 and given  $X = x$ ,  $Y$  is exponential with parameter  $\lambda = x$ .

Find  $Cov(X, Y)$ .


Cov  $(X,Y) = E[XY] - E[X] = [Y]$  We have  $E[X] = \frac{3}{2}$ .

We have  $E[Y] = E[E[Y|X]]_2$  (by LoJ.t)

 $= E[XY] = E[XY|X]$  (LoJ.t)

 $= E[XY] = E[XY|X]$  (LoJ.t)

 $= E[XY|X] = E[XY|X]$  (LoJ.t)



» Properties of Covariance...

#### **Properties of Covariance**

1. 
$$Cov(X, X) = Var(X)$$

» Properties of Covariance...

#### Properties of Covariance

 $\mathsf{L}.\mathsf{Cov}(X,X) = \mathsf{Var}(X)$ 



 $\nearrow$  If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]

» Properties of Covariance...

#### **Properties of Covariance**

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- $\mathcal{S}$ .  $\mathsf{Cov}(X,Y) = \mathsf{Cov}(Y,X)$
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)


- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)
- 6. Cov(X + Y, Z) = Cov(X, Z) + Cov(Y, Z)

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- 4. Cov(aX, Y) = aCov(X, Y)
- 5. Cov(X + c, Y) = Cov(X, Y)
- 6. Cov(X + Y, Z) = Cov(X, Z) + Cov(Y, Z)
- 7. More generally, we have

- 1. Cov(X, X) = Var(X)
- 2. If X and Y are independent, then Cov(X, Y) = 0. [Note: converse is not true!]
- 3. Cov(X, Y) = Cov(Y, X)
- -4. Cov(aX, Y) = aCov(X, Y)
  - $6. \operatorname{Cov}(X + c, Y) = \operatorname{Cov}(X, Y)$
- $6. \operatorname{Cov}(X + Y, Z) = \operatorname{Cov}(X, Z) + \operatorname{Cov}(Y, Z)$
- 7. More generally, we have

$$\mathsf{Cov}\left(\sum_{i=1}^m a_i X_i, \ \sum_{j=1}^n b_j Y_j\right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \mathsf{Cov}(X_i, Y_j)$$





» Solved Example...

Let X and Y be two independent random variables following standard normal distribution, and

$$Z = 1 + X + XY^{2}$$

$$W = 1 + X$$

Find 
$$Cov(Z, W)$$
.
Solution

Find 
$$Cov(2, W)$$
.

Solution
$$Cov(2, W) = Cov(1+x+xy^2, 1+x) = Cov(x+xy, x)$$

$$= Cov(x,x) + Cov(x,x) + (Rop6)$$


$$= Vov(x) + E(xy^2) - E(xy^2) E(x) + (Sinu x,y ind x)$$

$$= 1 + E(xy) E(xy^2) - E(xy^2) E(yy^2)$$

$$= 1 + E(xy) E(xy^2) - E(xy^2) E(yy^2)$$

$$= 1 + E(xy^2) E(yy^2) = 1 + 1 = 2$$

$$= 1 + 2 \cdot 1 - 0 \cdot 1 = 1 + 1 = 2$$



Example (Variance of a Sum)

Let Z = X + Y, then

Example (Variance of a Sum)

Let Z = X + Y, then

$$\mathsf{Var}(\mathbf{Z}) = \mathsf{Var}(\mathbf{X}) + \mathsf{Var}(\mathbf{Y}) + 2\mathsf{Cov}(\mathbf{X},\mathbf{Y})$$

Example (Variance of a Sum)

Let Z = X + Y, then

$$\mathsf{Var}(\mathbf{Z}) = \mathsf{Var}(\mathbf{X}) + \mathsf{Var}(\mathbf{Y}) + 2\mathsf{Cov}(\mathbf{X},\mathbf{Y})$$

More generally, for  $a,b\in\mathbb{R},$  and Z=aX+bY, we have

Example (Variance of a Sum)

Let 
$$Z = X + Y$$
, then

$$Var(Z) = Var(X) + Var(Y) + 2Cov(X, Y)$$

More generally, for  $a,b\in\mathbb{R}$ , and Z=aX+bY, we have

$$Var(Z) = a^{2} Var(X) + b^{2} Var(Y) + 2ab Cov(X, Y)$$
Solution
$$Var(Z) = Cov(Z, Z) = Cov(X+Y, X+Y)$$

Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

#### Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{var}(X) \text{Var}(Y)}}$$

#### Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

\* Given two RVs X and Y, define standardized versions of these as follows

#### Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

\* Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

#### Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X\,\rho_Y}$$

\* Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

\* With this by computing the covariance Cov(U,V), we have

#### Correlation coefficient

The correlation coefficient  $\rho_{XY}$  or  $\rho(X, Y)$  is defined as follows

$$\rho_{XY} = \rho(X, Y) = \frac{\mathsf{Cov}(X, Y)}{\sqrt{\mathsf{Var}(X)\,\mathsf{Var}(Y)}} = \frac{\mathsf{Cov}(X, Y)}{\rho_X \,\rho_Y}$$

\* Given two RVs X and Y, define standardized versions of these as follows

$$U = \frac{X - E[X]}{\rho_X}, \quad V = \frac{Y - E[Y]}{\rho_Y}$$

 $\triangleright$  With this by computing the covariance Cov(U, V), we have

$$\begin{aligned}
\rho_{XY} &= \operatorname{Cov}(U, V) = \operatorname{Cov}\left(\frac{X - E[X]}{\rho_X}, \frac{Y - E[Y]}{\rho_Y}\right) \\
&= \operatorname{Cov}\left(\frac{X}{\rho_X}, \frac{Y}{\rho_Y}\right) = \frac{\operatorname{Cov}(X, Y)}{\rho_X \rho_Y} = 0
\end{aligned}$$



Properties of correlation coefficient

Let X, Y be two RVs.

Properties of correlation coefficient

### Properties of correlation coefficient

1. 
$$-1 \le \rho(X, Y) \le 1$$

### Properties of correlation coefficient

- 1.  $-1 \le \rho(X, Y) \le 1$
- 2. If  $\rho(X, Y) = 1$ ,

## Properties of correlation coefficient

- 1.  $-1 \le \rho(X, Y) \le 1$
- 2. If  $\rho(X, Y) = 1$ , then Y = aX + b, where a > 0

## Properties of correlation coefficient

1. 
$$-1 \le \rho(X, Y) \le 1$$

2. If 
$$\rho(X, Y) = 1$$
, then  $Y = aX + b$ , where  $a > 0$ 

3. If 
$$\rho(X, Y) = 1$$
,

## Properties of correlation coefficient

1. 
$$-1 \le \rho(X, Y) \le 1$$

2. If 
$$\rho(X, Y) = 1$$
, then  $Y = aX + b$ , where  $a > 0$ 

3. If 
$$\rho(X, Y) = 1$$
, then  $Y = aX + b$ , where  $a < 0$ 

### Properties of correlation coefficient

$$1/-1 \le \rho(X, Y) \le 1$$

If 
$$\rho(X, Y) = 1$$
, then  $Y = aX + b$ , where  $a > 0$ 

If 
$$\rho(X, Y) = 1$$
, then  $Y = aX + b$ , where  $a < 0$ 

4. 
$$\rho(aX + b, cY + d) = \rho(X, Y)$$
 for  $a, c > 0$ 





8/1

» Positive Correlation, Negative Correlation, Uncorrelation...

Definition of positive, negative correlation

Let X and Y be two RVs.

### Definition of positive, negative correlation

Let X and Y be two RVs.

1. If  $\rho(\mathbf{X},\mathbf{Y})=0$ , we say that  $\mathbf{X}$  and  $\mathbf{Y}$  are uncorrelated

### Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If  $\rho(X, Y) = 0$ , we say that X and Y are uncorrelated
- 2. If  $\rho(X, Y) > 0$ , we say that X and Y are positively correlated

### Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If  $\rho(X, Y) = 0$ , we say that X and Y are uncorrelated
- 2. If  $\rho(\mathbf{X},\mathbf{Y})>0$ , we say that  $\mathbf{X}$  and  $\mathbf{Y}$  are positively correlated
- 3. If  $\rho(\mathbf{X},\mathbf{Y})<0$ , we say that  $\mathbf{X}$  and  $\mathbf{Y}$  are negatively correlated

#### Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If  $\rho(X, Y) = 0$ , we say that X and Y are uncorrelated
- 2. If  $\rho(X, Y) > 0$ , we say that X and Y are positively correlated
- 3. If  $\rho(X,Y) < 0$ , we say that X and Y are negatively correlated

#### Pairwise uncorrelation and Variance

If X and Y are uncorrelated, then

$$Var(X + Y) = Var(X) + Var(Y)$$

More generally, if  $X_1, X_2, \dots, X_n$ , are pairwise uncorrelated, i.e.,  $\rho(X_i, X_j) = 0$  when  $i \neq j$ , then

### Definition of positive, negative correlation

Let X and Y be two RVs.

- 1. If  $\rho(X, Y) = 0$ , we say that X and Y are uncorrelated
- 2. If  $\rho(X, Y) > 0$ , we say that X and Y are positively correlated
- 3. If  $\rho(X, Y) < 0$ , we say that X and Y are negatively correlated

#### Pairwise uncorrelation and Variance

If X and Y are uncorrelated, then

$$Var(X + Y) = Var(X) + Var(Y)$$

More generally, if  $X_1, X_2, \dots, X_n$ , are pairwise uncorrelated, i.e.,  $\rho(X_i, X_j) = 0$  when  $i \neq j$ , then

$$Var(X_1 + X_2 + \cdots + X_n) = Var(X_1) + Var(X_2) + \cdots + Var(X_n)$$