

Probability and Statistics: Lecture-37

Monsoon-2020

by Dr. Pawan Kumar (IIIT, Hyderabad)
on November 9, 2020

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n discrete RVs.

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n **discrete** RVs.

* The **joint PMF** of X_1, X_2, \dots, X_n is defined as

$$P((X_1, X_2, \dots, X_n) \in A) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n **discrete** RVs.

- * The **joint PMF** of X_1, X_2, \dots, X_n is defined as

$$P((X_1, X_2, \dots, X_n) \in A) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

- * Let X_1, X_2, \dots, X_n be n **continuous** RVs,

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n **discrete** RVs.

- * The **joint PMF** of X_1, X_2, \dots, X_n is defined as

$$P(\underline{X_1, X_2, \dots, X_n} \in A) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

- * Let X_1, X_2, \dots, X_n be n **continuous** RVs, then **PDF** is denoted by

$$f_{\underline{X_1, X_2, \dots, X_n}}(x_1, x_2, \dots, x_n),$$

and the **probability** is computed as follows

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n **discrete** RVs.

- * The **joint PMF** of X_1, X_2, \dots, X_n is defined as

$$P((X_1, X_2, \dots, X_n) \in A) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

- * Let X_1, X_2, \dots, X_n be n **continuous** RVs, then **PDF** is denoted by

$$f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n),$$

and the **probability** is computed as follows

$$P((X_1, X_2, \dots, X_n) \in A) = \int \cdots \int_A \cdots \int f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n) dx_1 \cdots dx_n$$

Joint PDF, Joint CDF

Let X_1, X_2, \dots, X_n be n **discrete** RVs.

- * The **joint PMF** of X_1, X_2, \dots, X_n is defined as

$$P((X_1, X_2, \dots, X_n) \in A) = P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

- * Let X_1, X_2, \dots, X_n be n **continuous** RVs, then **PDF** is denoted by

$$f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n),$$

and the **probability** is computed as follows

$$P((X_1, X_2, \dots, X_n) \in A) = \int \dots \int_A \dots \int f_{X_1 X_2 \dots X_n}(x_1, x_2, \dots, x_n) dx_1 \dots dx_n$$

Multiple Random Variables: Marginal PDF and Joint CDF...

Multiple Random Variables: Marginal PDF and Joint CDF...

- * The **marginal PDF** of RV X_i can be obtained

Multiple Random Variables: Marginal PDF and Joint CDF...

- * The **marginal PDF** of RV X_i can be obtained by **integrating** out all other X_j 's, for example,

Multiple Random Variables: Marginal PDF and Joint CDF...

- * The **marginal PDF** of RV X_i can be obtained by **integrating** out all other X'_j s, for example,

$$\underbrace{f_{X_1}}(x_1) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1 X_2 \dots X_n} \underbrace{dx_2}_{\text{ }} \underbrace{dx_3}_{\text{ }} \cdots \underbrace{dx_n}_{\text{ }}$$

Multiple Random Variables: Marginal PDF and Joint CDF...

- * The **marginal PDF** of RV X_i can be obtained by **integrating** out all other X_j 's, for example,

$$f_{X_1}(x_1) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1 X_2 \dots X_n} dx_2 dx_3 \cdots dx_n$$

- * The **joint CDF** of n random variables X_1, X_2, \dots, X_n is defined as

Multiple Random Variables: Marginal PDF and Joint CDF...

- * The **marginal PDF** of RV X_i can be obtained by **integrating** out all other X'_j s, for example,

$$f_{X_1}(x_1) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1 X_2 \dots X_n} dx_2 dx_3 \cdots dx_n$$

- * The **joint CDF** of n random variables X_1, X_2, \dots, X_n is defined as

$$F_{\underbrace{X_1, X_2, \dots, X_n}}(\underbrace{x_1, x_2, \dots, x_n}) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n)$$

» Example

Example (Three jointly continuous RVs)

Let X , Y and Z be three jointly continuous random variables

» Example

Example (Three jointly continuous RVs)

Let X , Y and Z be three jointly continuous random variables with joint PDF

» Example

Example (Three jointly continuous RVs)

Let X , Y and Z be three **jointly continuous** random variables with **joint PDF**

$$f_{XYZ}(x, y, z) = \begin{cases} c(x + 2y + 3z) & 0 \leq x, y, z \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

» Example

Example (Three jointly continuous RVs)

Let X , Y and Z be three **jointly continuous** random variables with **joint PDF**


$$\underbrace{f_{XYZ}}(x, y, z) = \begin{cases} c(x + 2y + 3z) & 0 \leq x, y, z \leq 1 \\ 0 & \text{otherwise} \end{cases}$$


1. Find the constant c

» Example

Example (Three jointly continuous RVs)

Let X , Y and Z be three **jointly continuous** random variables with **joint PDF**

$$f_{XYZ}(x, y, z) = \begin{cases} c(x + 2y + 3z) & 0 \leq x, y, z \leq 1 \\ 0 & \text{otherwise} \end{cases}$$


1. Find the constant c
 2. Find the **marginal PDF** of X
- 

» Answer to previous problem...

① We have

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{xyz}(x, y, z) dx dy dz \\ &= \int_0^1 \int_0^1 \int_0^1 c(x + 2y + 3z) dx dy dz \\ &= \int_0^1 \int_0^1 \left[c \left(\frac{x^2}{2} + 2yx + 3zx \right) \right]_0^1 dy dz \\ &= \int_0^1 \int_0^1 c \left[\frac{1}{2} + 2y + 3z \right] dy dz \end{aligned}$$

$$= \int_0^1 c \left[\frac{1}{2}y + \frac{2y^2}{2} + 3zy \right]_0^1 dz$$

$$= \int_0^1 c \left[\frac{3}{2} + 3z \right] dz$$

$$= c \left[\frac{3}{2}z + \frac{3z^2}{2} \right]_0^1$$

$$= c \cdot 3$$

$$\Rightarrow c = \frac{1}{3}$$

$$\begin{aligned} \textcircled{2} f_X(x) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{xyz}(x, y, z) dy dz \\ &= \int_0^1 \int_0^1 \frac{1}{3} (x + 2y + 3z) dy dz = \text{---} \end{aligned}$$

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

» Independence of Multiple Random Variables...

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$\underline{F_{X_1, X_2, \dots, X_n}}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

» Independence of Multiple Random Variables...

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

* If X_1, X_2, \dots, X_n are **discrete**, then they are **independent** if

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

* If X_1, X_2, \dots, X_n are **discrete**, then they are **independent** if

$$P_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P_{X_1}(x_1)P_{X_2}(x_2) \cdots P_{X_n}(x_n)$$

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **discrete**, then they are **independent** if

$$\underline{P_{X_1, X_2, \dots, X_n}}(x_1, x_2, \dots, x_n) = P_{X_1}(x_1)P_{X_2}(x_2) \cdots P_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **continuous**, then they are **independent** if

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **discrete**, then they are **independent** if

$$P_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P_{X_1}(x_1)P_{X_2}(x_2) \cdots P_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **continuous**, then they are **independent** if

$$\underline{f_{X_1, X_2, \dots, X_n}}(x_1, x_2, \dots, x_n) = f_{X_1}(x_1)f_{X_2}(x_2) \cdots f_{X_n}(x_n)$$

Independence of Multiple Random Variables

The n random variables X_1, X_2, \dots, X_n are **independent** if for all $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$

$$F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **discrete**, then they are **independent** if

$$P_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P_{X_1}(x_1)P_{X_2}(x_2) \cdots P_{X_n}(x_n)$$

- * If X_1, X_2, \dots, X_n are **continuous**, then they are **independent** if

$$f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = f_{X_1}(x_1)f_{X_2}(x_2) \cdots f_{X_n}(x_n)$$

- * If random variables X_1, X_2, \dots, X_n are **independent**, then we have

$$E[X_1, X_2, \dots, X_n] = E[X_1]E[X_2] \cdots E[X_n]$$

» Independent and Identically Distributed Random Variables (I.I.D.)

» Independent and Identically Distributed Random Variables (I.I.D.)

Definition of I.I.D.

Random variables X_1, X_2, \dots, X_n are said to be **independent and identically distributed (i.i.d.)**

» Independent and Identically Distributed Random Variables (I.I.D.)

Definition of I.I.D.

Random variables X_1, X_2, \dots, X_n are said to be independent and identically distributed (i.i.d.) if they are **independent**,

» Independent and Identically Distributed Random Variables (I.I.D.)

Definition of I.I.D.

Random variables X_1, X_2, \dots, X_n are said to be **independent and identically distributed (i.i.d.)** if they are **independent**, and they have the **same marginal** distributions:

» Independent and Identically Distributed Random Variables (I.I.D.)

Definition of I.I.D.

Random variables X_1, X_2, \dots, X_n are said to be **independent and identically distributed (i.i.d.)** if they are **independent**, and they have the **same marginal** distributions:


$$\underline{F_{X_1}(x)} = \underline{F_{X_2}(x_2)} = \dots = \underline{F_{X_n}(x)}, \quad \text{for all } x \in \mathbb{R}$$

» Independent and Identically Distributed Random Variables (I.I.D.)

Definition of I.I.D.

Random variables X_1, X_2, \dots, X_n are said to be **independent and identically distributed (i.i.d.)** if they are **independent**, and they have the **same marginal** distributions:

$$F_{\underline{X_1}}(x) = F_{\underline{X_2}}(x_2) = \dots = F_{\underline{X_n}}(x), \quad \text{for all } x \in \mathbb{R}$$

- * If we flip the same coin N times and record the outcome, then $\underline{X_1}, \dots, \underline{X_n}$ are I.I.D. 
- * Verify that these I.I.D. variables will have **same** mean and **variances**

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

- * The expectation of Y is

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \cdots + E[X_n]$$

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \cdots E[X_n]$$

- * We recall that

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \cdots E[X_n]$$

- * We recall that

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\text{Cov}(X_1, X_2)$$

Expectation and Variance

Let $Y = X_1 + X_2 + \cdots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \cdots E[X_n]$$

- * We recall that

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\text{Cov}(X_1, X_2)$$

In general, for Y , we have

Expectation and Variance

Let $Y = X_1 + X_2 + \dots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \dots + E[X_n]$$

- * We recall that

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\text{Cov}(\underline{X_1}, \underline{X_2})$$

In general, for Y , we have

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

Expectation and Variance

Let $Y = X_1 + X_2 + \dots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \dots + E[X_n]$$

- * We recall that

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\text{Cov}(X_1, X_2)$$

In general, for Y , we have

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

- * If X_1, X_2, \dots, X_n are **independent**, then

» Expectation and Variance...

$$\text{Cov}(X, Y) = \frac{E(XY)}{n} - E[X]E[Y]$$

Expectation and Variance

Let $Y = X_1 + X_2 + \dots + X_n$. Then we have the following

- * The expectation of Y is

$$E[Y] = E[X_1] + E[X_2] + \dots + E[X_n]$$

- * We recall that

$$\text{Var}(X_1 + X_2) = \text{Var}(X_1) + \text{Var}(X_2) + 2\underline{\text{Cov}(X_1, X_2)}$$

$$\begin{aligned} \text{Var}(X_1 + X_2) &= \text{Var}(X_1) + \text{Var}(X_2) \\ &+ 2\text{Cov}(X_1, X_2) \end{aligned}$$

In general, for Y , we have

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

- * If $X_1, \underline{X_2}, \dots, X_n$ are **independent**, then

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i)$$

» Example...

» Example...

Example (Who will receive the present?)

N people sit around a round table, where $N > 5$.

» Example...

Example (Who will receive the present?)

N people sit around a round table, where $N > 5$. Each person tosses a coin.

» Example...

Example (Who will receive the present?)

N people sit around a round table, where $N > 5$. Each person tosses a coin. Anyone whose outcome is different from his/her two neighbors will receive a present.

» Example...

Example (Who will receive the present?)

N people sit around a round table, where $N > 5$. Each person tosses a coin. Anyone whose outcome is different from his/her two neighbors will receive a present. Let X be the number of people who receive presents.

» Example...

Example (Who will receive the present?)

N people sit around a round table, where $N > 5$. Each person tosses a coin. Anyone whose outcome is different from his/her two neighbors will receive a present. Let X be the number of people who receive presents. Find $E[X]$ and $\text{Var}(X)$.

Try

» Answer to previous problem...

» Answer to previous problem...

» Answer to previous problem...

PDF of the Sum of Multiple RVs

We recall that if $Y = \underbrace{X_1 + X_2}$, and X_1 and X_2 being independent, we have

» PDF of the Sum of Multiple Random Variables...

$$\bar{X} = [X_1, X_2, \dots, X_n]$$
$$\bar{Y} = [Y_1, Y_2, \dots, Y_n]$$

PDF of the Sum of Multiple RVs

We recall that if $Y = \underbrace{X_1}_{\text{RV}} + \underbrace{X_2}_{\text{RV}}$, and X_1 and X_2 being independent, we have

$$\underbrace{f_Y(y)}_{\text{PDF}} = f_{X_1}(y) * \underbrace{f_{X_2}(y)}_{\text{PDF}} = \int_{-\infty}^{\infty} \underbrace{f_{X_1}(x)}_{\text{PDF}} \underbrace{f_{X_2}(y-x)}_{\text{PDF}} dx$$

Convolution

For multiple variable case, i.e., if $Y = \underbrace{X_1}_{\text{RV}} + \underbrace{X_2}_{\text{RV}} + \dots + \underbrace{X_n}_{\text{RV}}$, we have

$$f_Y(y) = \underbrace{f_{X_1}(y) * f_{X_2}(y) * \dots * f_{X_n}(y)}_{\text{PDF}}$$

* However, it is computationally difficult!

Definition of Random Vector and Expectation

When dealing with multiple RVs, it is useful to use vector and matrix notations.

Definition of Random Vector and Expectation

When dealing with multiple RVs, it is useful to use vector and matrix notations. This helps using a compact notation and tools from linear algebra.

Definition of Random Vector and Expectation

When dealing with multiple RVs, it is useful to use vector and matrix notations. This helps using a compact notation and tools from linear algebra. Let X_1, X_2, \dots, X_n be n independent RVs, then random vector is defined as

Definition of Random Vector and Expectation

When dealing with multiple RVs, it is useful to use vector and matrix notations. This helps using a compact notation and tools from linear algebra. Let X_1, X_2, \dots, X_n be n independent RVs, then random vector is defined as

$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \quad E[X] = \begin{bmatrix} E[X_1] \\ E[X_2] \\ \vdots \\ E[X_n] \end{bmatrix}$$

» Definition of Random Vector...

Definition of Random Vector and Expectation

When dealing with multiple RVs, it is useful to use vector and matrix notations. This helps using a compact notation and tools from linear algebra. Let X_1, X_2, \dots, X_n be n independent RVs, then random vector is defined as

$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \quad E[X] = \begin{bmatrix} E[X_1] \\ E[X_2] \\ \vdots \\ E[X_n] \end{bmatrix}$$

We call X a **random vector** and $E[X]$ is the expectation of random vector.

- * CDF is $F_X(x) = F_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = P(X_1 \leq x_1, \dots, X_n \leq x_n)$
- * If X is jointly continuous, the PDF is denoted as

$$f_X(x) = f_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n)$$

Short-hand notation

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$
$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix}$$

Random matrix, Expectation

A **random matrix** is a matrix whose elements are **random** variables.

Random matrix, Expectation

A **random matrix** is a matrix whose elements are **random** variables.

$$M = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ X_{m1} & X_{m2} & \dots & X_{mn} \end{bmatrix} \quad E[M] = \begin{bmatrix} E[X_{11}] & E[X_{12}] & \dots & E[X_{1n}] \\ E[X_{21}] & E[X_{22}] & \dots & E[X_{2n}] \\ \vdots & \vdots & \ddots & \vdots \\ E[X_{m1}] & E[X_{m2}] & \dots & E[X_{mn}] \end{bmatrix}$$

Random matrix, Expectation


A **random matrix** is a matrix whose elements are **random** variables.

$$M = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ X_{m1} & X_{m2} & \dots & X_{mn} \end{bmatrix} \quad E[M] = \begin{bmatrix} E[X_{11}] & E[X_{12}] & \dots & E[X_{1n}] \\ E[X_{21}] & E[X_{22}] & \dots & E[X_{2n}] \\ \vdots & \vdots & \ddots & \vdots \\ E[X_{m1}] & E[X_{m2}] & \dots & E[X_{mn}] \end{bmatrix}$$


Here M is called the random matrix, and $E[M]$ is the expectation of random matrix.

Random matrix, Expectation

A **random matrix** is a matrix whose elements are **random** variables.

$$M = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ X_{m1} & X_{m2} & \dots & X_{mn} \end{bmatrix} \quad E[M] = \begin{bmatrix} E[X_{11}] & E[X_{12}] & \dots & E[X_{1n}] \\ E[X_{21}] & E[X_{22}] & \dots & E[X_{2n}] \\ \vdots & \vdots & \vdots & \vdots \\ E[X_{m1}] & E[X_{m2}] & \dots & E[X_{mn}] \end{bmatrix}$$


Here M is called the **random matrix**, and $E[M]$ is the **expectation** of random matrix.

- * If $Y = AX + b$, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, then $E[Y] = AE[X] + b$
 - * Also, if X_1, X_2, \dots, X_k are k n -dimensional RVs, then
- 

Random matrix, Expectation

A **random matrix** is a matrix whose elements are **random** variables.

$$M = \begin{bmatrix} X_{11} & X_{12} & \dots & X_{1n} \\ X_{21} & X_{22} & \dots & X_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ X_{m1} & X_{m2} & \dots & X_{mn} \end{bmatrix} \quad E[M] = \begin{bmatrix} E[X_{11}] & E[X_{12}] & \dots & E[X_{1n}] \\ E[X_{21}] & E[X_{22}] & \dots & E[X_{2n}] \\ \vdots & \vdots & \vdots & \vdots \\ E[X_{m1}] & E[X_{m2}] & \dots & E[X_{mn}] \end{bmatrix}$$

Here M is called the **random matrix**, and $E[M]$ is the **expectation** of random matrix.

- * If $Y = AX + b$, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, then $E[Y] = AE[X] + b$
- * Also, if $\underline{X_1}, \underline{X_2}, \dots, \underline{X_k}$ are k n -dimensional RVs, then

$$E[\underline{X_1} + \underline{X_2} + \dots + \underline{X_k}] = E[X_1] + E[X_2] + \dots + E[X_k].$$

Variance of Multiple Random Variables

Let X_1, X_2, \dots, X_n be n random variables.

Variance of Multiple Random Variables

Let X_1, X_2, \dots, X_n be n random variables. Then we have the following

» Variance of a Sum of Random Variables...

Variance of Multiple Random Variables

Let X_1, X_2, \dots, X_n be n random variables. Then we have the following

$$\text{Var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

» Variance of a Sum of Random Variables...

Variance of Multiple Random Variables

Let X_1, X_2, \dots, X_n be n random variables. Then we have the following

$$\text{Var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

If X_1, X_2, \dots, X_n are **independent**, then we have

Variance of Multiple Random Variables

Let X_1, X_2, \dots, X_n be n random variables. Then we have the following

$$\text{Var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i < j} \text{Cov}(X_i, X_j)$$

If X_1, X_2, \dots, X_n are **independent**, then we have

$$\text{Var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{Var}(X_i)$$

Definition of Correlation and Covariance Matrix

For a random vector X , the **correlation matrix** \underline{R}_X and **covariance matrix** \underline{C}_X is

» Correlation and Covariance Matrix...

Outer product $\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix} = \begin{bmatrix} x_1^2 & x_1 x_2 & \dots & x_1 x_n \\ x_2 x_1 & x_2^2 & \dots & x_2 x_n \\ \vdots & \vdots & \ddots & \vdots \\ x_n x_1 & x_n x_2 & \dots & x_n^2 \end{bmatrix}$

Definition of Correlation and Covariance Matrix

For a random vector X , the **correlation matrix** R_X and **covariance matrix** C_X is

$$R_X = E[XX^T] = E \begin{bmatrix} X_1^2 & X_1 X_2 & \dots & X_1 X_n \\ X_2 X_1 & X_2^2 & \dots & X_2 X_n \\ \vdots & \vdots & \ddots & \vdots \\ X_n X_1 & X_n X_2 & \dots & X_n^2 \end{bmatrix} = \begin{bmatrix} E[X_1^2] & E[X_1 X_2] & \dots & E[X_1 X_n] \\ E[X_2 X_1] & E[X_2^2] & \dots & E[X_2 X_n] \\ \vdots & \vdots & \ddots & \vdots \\ E[X_n X_1] & E[X_n X_2] & \dots & E[X_n^2] \end{bmatrix}$$

$$C_X = E[(X - E[X])(X - E[X])^T] = \begin{bmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \dots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Var}(X_n) \end{bmatrix}$$

$$U = \begin{bmatrix} x \\ y \end{bmatrix}$$

Definition of Correlation and Covariance Matrix

For a random vector X , the **correlation matrix** R_X and **covariance matrix** C_X is

$$R_X = E[XX^T] = \begin{bmatrix} X_1^2 & X_1X_2 & \dots & X_1X_n \\ X_2X_1 & X_2^2 & \dots & X_2X_n \\ \vdots & \vdots & \ddots & \vdots \\ X_nX_1 & X_nX_2 & \dots & X_n^2 \end{bmatrix} = \begin{bmatrix} E[X_1^2] & E[X_1X_2] & \dots & E[X_1X_n] \\ E[X_2X_1] & E[X_2^2] & \dots & E[X_2X_n] \\ \vdots & \vdots & \ddots & \vdots \\ E[X_nX_1] & E[X_nX_2] & \dots & E[X_n^2] \end{bmatrix}$$

$$C_X = E[(X - E[X])(X - E[X])^T] = \begin{bmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \dots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Var}(X_n) \end{bmatrix}$$

$$C_X = R_X - E[X]E[X]^T$$

Recall: $\text{Cov}(X,Y) = \text{Cov}(Y,X)$

Definition of Correlation and Covariance Matrix

For a random vector X , the **correlation matrix** R_X and **covariance matrix** C_X is

$$R_X = E[XX^T] = \begin{bmatrix} X_1^2 & X_1X_2 & \dots & X_1X_n \\ X_2X_1 & X_2^2 & \dots & X_2X_n \\ \vdots & \vdots & \ddots & \vdots \\ X_nX_1 & X_nX_2 & \dots & X_n^2 \end{bmatrix} = \begin{bmatrix} E[X_1^2] & E[X_1X_2] & \dots & E[X_1X_n] \\ E[X_2X_1] & E[X_2^2] & \dots & E[X_2X_n] \\ \vdots & \vdots & \ddots & \vdots \\ E[X_nX_1] & E[X_nX_2] & \dots & E[X_n^2] \end{bmatrix}$$

$$C_X = E[(X - E[X])(X - E[X])^T] = \begin{bmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \dots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \dots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \dots & \text{Var}(X_n) \end{bmatrix}$$

$(i,j) = (j,i)$

1. $C_X = R_X - E[X]E[X]^T$

2. If $Y = AX + b$, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, then $C_Y = AC_XA^T$

» Answer to previous problem...

» Example of Correlation and Covariance Matrices...

Example (Example of correlation and covariance matrices)

Let X and Y be jointly continuous random variables with joint PDF

» Example of Correlation and Covariance Matrices...

Example (Example of correlation and covariance matrices)

Let X and Y be jointly continuous random variables with joint PDF

$$f_{X,Y} = \begin{cases} \frac{3}{2}x^2 + y & 0 < x, y < 1 \\ 0 & \text{otherwise} \end{cases}$$

» Example of Correlation and Covariance Matrices...

Example (Example of correlation and covariance matrices)

Let X and Y be jointly continuous random variables with joint PDF

$$f_{X,Y} = \begin{cases} \frac{3}{2}x^2 + y & 0 < x, y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Let $U = \begin{bmatrix} X \\ Y \end{bmatrix}$ be the random vector.

» Example of Correlation and Covariance Matrices...

Example (Example of correlation and covariance matrices)

Let X and Y be jointly continuous random variables with joint PDF

$$f_{X,Y} = \begin{cases} \frac{3}{2}x^2 + y & 0 < x, y < 1 \\ 0 & \text{otherwise} \end{cases}$$

Let $U = \begin{bmatrix} X \\ Y \end{bmatrix}$ be the random vector. Find the **correlation and covariance matrices** of U .

» Answer to previous problem...

Similarly,

$$S_T(y) = \int_0^1 \left(\frac{3}{2}x^2 + y \right) dx$$

$$= y + \frac{1}{2} \quad 0 < y < 1.$$

We recall

$$R = \begin{bmatrix} \frac{E[X^2]}{\sigma^2} & \frac{E[XY]}{E[Y^2]} \\ \frac{E[YX]}{\sigma^2} & \frac{E[Y^2]}{E[Y^2]} \end{bmatrix}$$

$$E[x^2] = \int_{-\infty}^{\infty} x^2 f_x(x) dx$$

$$= \int_0^1 x^2 \cdot \left(\frac{3}{2}x^2 + \frac{1}{2} \right) dx$$

$$E[xy] = \int_0^1 \int_0^1 (xy) \left(\frac{3}{2}x^2 + y \right) dx dy$$

» Answer to previous problem...

$$\text{Cov}(X, Y) = \underline{E[XY]} - \underline{E[X]} \underline{E[Y]}$$

Covariance Matrix

$$C_u = E \left[(U - E[U]) (U - E[U])^T \right]$$
$$= \begin{bmatrix} \underline{\text{Var}(X)} & \underline{\text{Cov}(X, Y)} \\ \underline{\text{Cov}(X, Y)} & \underline{\text{Var}(Y)} \end{bmatrix}$$

Properties of Covariance

We have the following properties for **covariance matrix**:

Properties of Covariance

We have the following properties for **covariance matrix**:

1. The covariance matrix C_X is **symmetric matrix**

Properties of Covariance

We have the following properties for **covariance matrix**:

1. The covariance matrix C_X is **symmetric matrix**
2. The covariance matrix C_X is **positive semi-definite (PSD)**

» Properties of Covariance Matrix...

Recall definition of semi-positive definite (SPD), Assume A symmetric:
A matrix A is SPD if $x^T A x \geq 0 \quad \forall x \neq 0$

Properties of Covariance

We have the following properties for covariance matrix:

1. The covariance matrix C_X is symmetric matrix
2. The covariance matrix C_X is positive semi-definite (PSD)
3. The covariance matrix is positive definite if and only if all its eigenvalues are larger than zero

$$x^T E[zz^T] x \geq 0$$

$$C_X = E[(x - E[x])(x - E[x])^T] = E[\underbrace{z z^T}_A]$$

① Symmetry: $A^T = (z z^T)^T = z z^T = A \Rightarrow A$ is symmetric
 ② S.P.D.

$$\begin{aligned} x^T (z z^T) x &= (z^T x)^T (z^T x) \\ \|z^T x\|_2 &\geq 0 \end{aligned}$$

» Properties of Covariance Matrix...

a x b.

$$x^T A x > 0 \quad \forall x \neq 0$$

If λ is eig. val. of $A \Rightarrow Au = \lambda u$, u eig. vec.
 u eig. vec. $\Rightarrow u \neq 0 \Rightarrow u^T A u > 0$ (A is P.D.) $\Rightarrow u^T \lambda u > 0$
 $\Rightarrow \lambda \|u\|_2^2 > 0$

Properties of Covariance

We have the following properties for **covariance matrix**:

$$\Rightarrow \boxed{\lambda > 0}$$

1. The covariance matrix C_X is **symmetric matrix**
2. The covariance matrix C_X is **positive semi-definite (PSD)**
3. The covariance matrix is **positive definite** if and only if all its **eigenvalues are larger than zero**
4. The covariance matrix is **positive definite** if and only if $\det(C_X) > 0$

$$\det(C_X) = \prod \lambda_i$$

But $\lambda_i > 0 \Rightarrow \det(C_X) > 0$

» Answer to previous problem...

» Answer to previous problem...

» Example of Covariance Matrix...

Example (Example of covariance matrix)

Consider two independent random variables X and Y .

» Example of Covariance Matrix...

Example (Example of covariance matrix)

Consider two independent random variables X and Y . Let X and Y follow Uniform(0, 1) distribution.

» Example of Covariance Matrix...

Example (Example of covariance matrix)

Consider two independent random variables X and Y . Let X and Y follow Uniform(0, 1) distribution. Let the random vectors U and V be defined as

» Example of Covariance Matrix...

Example (Example of covariance matrix)

Consider two independent random variables X and Y . Let X and Y follow Uniform(0, 1) distribution. Let the random vectors U and V be defined as

$$U = \begin{bmatrix} X \\ X + Y \end{bmatrix}, \quad V = \begin{bmatrix} X \\ Y \\ X + Y \end{bmatrix}$$

» Example of Covariance Matrix...

Example (Example of covariance matrix)

Consider two independent random variables X and Y . Let X and Y follow Uniform(0, 1) distribution. Let the random vectors U and V be defined as

$$U = \begin{bmatrix} X \\ X + Y \end{bmatrix}, \quad V = \begin{bmatrix} X \\ Y \\ X + Y \end{bmatrix}$$

Are the matrices C_U and C_V positive definite?

» Answer to previous problem...

» Answer to previous problem...

» Denition of Cross-Correlation and Cross-Covariance Matrix...

$$R_x = E[xx^T]$$
$$C_x = E[(x - E[x])(x - E[x])^T]$$

Definition of Cross Correlation and Cross Covariance Matrices

Let X and Y be two random vectors,

Definition of Cross Correlation and Cross Covariance Matrices

Let X and Y be two random vectors, we define the **cross-correlation matrix** of X and Y as

» Denition of Cross-Correlation and Cross-Covariance Matrix...

Definition of Cross Correlation and Cross Covariance Matrices

Let X and Y be two random vectors, we define the **cross-correlation matrix** of X and Y as

$$R_{XY} = E[XY^T]$$

Definition of Cross Correlation and Cross Covariance Matrices

Let X and Y be two random vectors, we define the **cross-correlation matrix** of X and Y as

$$R_{XY} = E[\underline{XY^T}]$$

Similarly the **cross-covariance matrix** of X and Y random **vectors** is defined as

» Denition of Cross-Correlation and Cross-Covariance Matrix...

Definition of Cross Correlation and Cross Covariance Matrices

Let X and Y be two random vectors, we define the **cross-correlation matrix** of X and Y as

$$R_{XY} = E[XY^T]$$

Similarly the **cross-covariance matrix** of X and Y random **vectors** is defined as

$$C_{XY} = E[(X - E[X])(Y - E[Y])^T]$$

» Functions of Random Variables...

Vectors

$$X \in \mathbb{R}^n.$$

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$,

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$, where $G: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and invertible function with continuous partial derivatives.

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$, where $G : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **continuous and invertible** function with **continuous partial derivatives**. If $H = G^{-1}$, then $X = H(Y)$.

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$, where $G: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **continuous and invertible** function with **continuous partial derivatives**. If $\underline{H} = G^{-1}$, then $X = H(Y)$.

$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} = \begin{bmatrix} H_1(Y_1, Y_2, \dots, Y_n) \\ H_2(Y_1, Y_2, \dots, Y_n) \\ \vdots \\ H_n(Y_1, Y_2, \dots, Y_n) \end{bmatrix}$$

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$, where $G : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **continuous and invertible** function with **continuous partial derivatives**. If $H = G^{-1}$, then $X = H(Y)$.

$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} = \begin{bmatrix} H_1(Y_1, Y_2, \dots, Y_n) \\ H_2(Y_1, Y_2, \dots, Y_n) \\ \vdots \\ H_n(Y_1, Y_2, \dots, Y_n) \end{bmatrix}$$

Then the **PDF** of Y , denoted by $f_{Y_1, Y_2, \dots, Y_n}(y_1, y_2, \dots, y_n)$ is given as follows

Method of Transformation for Functions of Random Vectors

Consider a function of random vector: $Y = G(X)$, where $G: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **continuous and invertible** function with **continuous partial derivatives**. If $H = G^{-1}$, then $X = \underline{H(Y)}$.

$$X = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} = \begin{bmatrix} H_1(Y_1, Y_2, \dots, Y_n) \\ H_2(Y_1, Y_2, \dots, Y_n) \\ \vdots \\ H_n(Y_1, Y_2, \dots, Y_n) \end{bmatrix}$$

Then the **PDF** of Y , denoted by $f_{Y_1, Y_2, \dots, Y_n}(y_1, y_2, \dots, y_n)$ is given as follows

$$\underline{f_Y(y) = f_X(H(y)) |J|}, \quad \text{where } J = \det \begin{bmatrix} \frac{\partial H_1}{\partial y_1} & \frac{\partial H_1}{\partial y_2} & \cdots & \frac{\partial H_1}{\partial y_n} \\ \frac{\partial H_2}{\partial y_1} & \frac{\partial H_2}{\partial y_2} & \cdots & \frac{\partial H_2}{\partial y_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial H_n}{\partial y_1} & \frac{\partial H_n}{\partial y_2} & \cdots & \frac{\partial H_n}{\partial y_n} \end{bmatrix}$$

» Example of Method of Transform for Function of Random Vector...

» Example of Method of Transform for Function of Random Vector...

Example (Example of Method of Transform for Function of Random Vector)

Let $\underline{Y} = \underline{A}\underline{X} + \underline{b}$, where \underline{X} is a n dimensional random vector,

» Example of Method of Transform for Function of Random Vector...

Example (Example of Method of Transform for Function of Random Vector)

Let $Y = AX + b$, where X is a n dimensional random vector, A be a fixed (non-random) n by n matrix,

» Example of Method of Transform for Function of Random Vector...

Example (Example of Method of Transform for Function of Random Vector)

Let $Y = AX + b$, where X is a n dimensional random vector, A be a fixed (non-random) n by n matrix, and b be a fixed n -dimensional vector.

» Example of Method of Transform for Function of Random Vector...

$$X = A^{-1}(Y - b) = H(Y)$$
$$J = \det(A^{-1}) = \frac{1}{\det(A)}$$

Example (Example of Method of Transform for Function of Random Vector)

invertible

Let $Y = AX + b$, where X is a n dimensional random vector, A be a fixed (non-random) n by n matrix, and b be a fixed n -dimensional vector. Find the PDF of Y in terms of X .

$$f_Y(Y) = f_X(H(Y)) |J|$$
$$= f_X(A^{-1}(Y - b)) |J| = \frac{1}{\det(A)} f_X(A^{-1}(Y - b))$$

» Answer to previous problem...

» Definition of Jointly Normal or Gaussian Random Vectors...

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**,

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**,

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are jointly normal, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a normal variable

variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be normal vector, if the random vectors X_1, X_2, \dots, X_n are jointly normal
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.**

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.** Then the PDF of Z is

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.** Then the PDF of Z is

$$f_Z(z) = \frac{1}{(2\pi)^{n/2}} \exp \left\{ -\frac{1}{2} z^T z \right\}$$

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.** Then the PDF of Z is

$$f_Z(z) = \frac{1}{(2\pi)^{n/2}} \exp \left\{ -\frac{1}{2} z^T z \right\}$$

4. For a normal random vector X , with mean m and covariance C ,

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.** Then the PDF of Z is

$$f_Z(z) = \frac{1}{(2\pi)^{n/2}} \exp \left\{ -\frac{1}{2} z^T z \right\}$$

4. For a normal random vector X , with mean m and covariance C , the **PDF** is

» Definition of Jointly Normal or Gaussian Random Vectors...

Definition of Jointly Normal Random Vectors

1. Random variables X_1, X_2, \dots, X_n are **jointly normal**, if the linear combination

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n, \quad a_1, a_2, \dots, a_n \in \mathbb{R}$$

is a **normal** variable

2. A random vector $X = [X_1, X_2, \dots, X_n]$ is said to be **normal vector**, if the random vectors X_1, X_2, \dots, X_n are **jointly normal**
3. Consider a random vector Z whose components $Z_i \sim N(0, 1)$, and they are **I.I.D.** Then the PDF of Z is

$$f_Z(z) = \frac{1}{(2\pi)^{n/2}} \exp \left\{ -\frac{1}{2} z^T z \right\}$$

4. For a normal random vector X , with mean m and covariance C , the **PDF** is

$$f_X(x) = \frac{1}{(2\pi)^{n/2} \sqrt{\det C}} \exp \left\{ -\frac{1}{2} (x - m)^T C^{-1} (x - m) \right\} \leftarrow$$