

Introductory Galois Theory Cheat Sheet

Definition of a Field

A field F is a set with two binary operators $(+, \times)$ satisfying the following axioms,

- $(F, +)$ is an abelian group with identity 0.
- The non zero elements of F form an abelian group under multiplication with identity $1 \neq 0$.
- Left and right distributivity

Characteristic of Fields

A characteristic of a field F , denoted by $\text{ch}(F)$ is defined as is the smallest integer p such that $\underbrace{1 + 1 + \dots + 1}_{p \text{ times}} = 0$. If such a p does not, exist $\text{ch}(F) = 0$.

K-algebra

A K-algebra (or algebra over a field) is a ring A which is a module over field K with multiplication being K-bilinear, (i.e., $k_1 a_1 \cdot k_2 a_2 = k_1 k_2 a_1 a_2$).

Field Extensions

For fields K, L . We say L is a field extension of K if K is a subfield of L . Alternatively, L is a field extension of K , if L is a K-algebra.

Algebraic elements and Algebraic extensions

For a field extension $K \subset L$.

Algebraic element: $\alpha \in L$ is called algebraic if $\exists P \in K[x]$ s.t. $P(\alpha) = 0$.

Transcendental element: If such a P does not exist then α is transcendental.

Consider the following definitions,

- Denote the smallest subfield of L containing K and α to be $K(\alpha)$.
- Denote the smallest sub ring of L containing K and α to be $K[\alpha]$.

The following statements are equivalent,

- α is algebraic over K .
- $K[\alpha]$ is finite dimensional algebra over K .
- $K[\alpha] = K(\alpha)$.

Algebraic extension: L is called algebraic over K if all $\alpha \in L$ are algebraic over K .

- If L is algebraic over K then any K -subalgebra of L is a field.
- Consider $K \subset L \subset M$. If $\alpha \in M$ is algebraic over K , then it is algebraic over L , also its minimal polynomial over L divides its minimal polynomial over K .
- If $K \subset L \subset M$ then M is an algebraic extension over $K \iff M$ is algebraic over L and L is algebraic over K .

Algebraic closure: A subfield L' of L s.t. $L' = \{\alpha \in L \mid \alpha \text{ is algebraic over } K\}$

Minimal Polynomial

If α is an algebraic element then $\exists!$ monic polynomial P of minimal degree such that $P(\alpha) = 0$ such a polynomial is called the **minimal polynomial**.

- The minimal polynomial is irreducible
- Any other polynomial Q s.t. $Q(\alpha) = 0$ will be divisible by P .

Primitive polynomials and Gauss' lemma

Primitive polynomial: A polynomial $P \in \mathbb{Z}[X]$ is called primitive if it has a positive degree and the gcd of its coefficients is 1.

Gauss' lemma: A polynomial $P \in \mathbb{Z}[X]$ is irreducible over $\mathbb{Z}[X] \iff$ it is primitive and irreducible over $\mathbb{Q}[x]$

Eisenstein criterion for irreducibility

A polynomial $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0 \in \mathbb{Z}[x]$ is irreducible if $\exists p$ prime s.t. p divides all coefficients except a_n and p^2 does not divide a_0 .

Finite extensions

For a field extension $K \subset L$. L is called a **finite extension** of K if the vector space of L over K has a finite dimension.

Degree of finite extension: Denoted as $[L : K] = \dim_K L$

- $K \subset L \subset M$. Then M is finite over $K \iff M$ is finite over L and L is finite over K . Also in this case, $[M : K] = [M : L][L : K]$.
- Let $K(\alpha_1, \dots, \alpha_n) \subset L$ denote the smallest subfield of L containing K and $\alpha_i \in L$. This $K(\alpha_1, \dots, \alpha_n)$ is generated by $\alpha_1, \dots, \alpha_n$.
- L is finite over $K \iff L$ is generated by a finite number of algebraic elements over K .
- $[K(\alpha) : K] = \deg P_{\min}(\alpha, K)$

Stem field

Let $P \in K[X]$ be an irreducible monic polynomial. A field extension E is called a stem field of P if $\exists \alpha \in E$, s.t. α is a root of P and $E = K[\alpha]$

- If E, E' are two stem fields for $P \in K[x]$, s.t. $E = K[\alpha], E' = K[\alpha']$ where α, α' are roots of P . Then $\exists!$ isomorphism $E \cong E'$ of K-algebras which maps α to α' .
- If a stem field contains two roots of P , then $\exists!$ automorphism that maps one root to another.
- If E is a stem field, $[E : K] = \deg P$
- If $[E : K] = \deg P$ and E contains a root of P then E is a stem field.

Some irreducibility criteria,

- $P \in K[X]$ is irreducible over $K \iff$ it does not have roots in L/K of degree $\leq \deg P/2$.
- $P \in K[X]$ is irreducible over K with $\deg P = n$. If L/K with $[L : K] = m$ if $\gcd(m, n) = 1$ then P is irreducible over L .

Splitting field

Let $P \in K[X]$. The splitting field of P over K is an extension of L where P is split into linear factors and the roots of P generate L (alternatively if P cannot be factored into any intermediate field).

- Splitting field L exists and its degree is $\leq d!$, where $d = \deg P$. And it is unique up to isomorphism.

Algebraic closure

- A field K is algebraically closed if any non-constant polynomial $P \in K[X]$ has a root in K .
- L is called an **algebraic closure** of K if it is algebraically closed and a field extension over K .
- Every field has an algebraic closure.
- Algebraic closures of K are unique up to isomorphism as K -algebras.

Properties of finite fields

Let p be a prime integer and let $q = p^r$ for some positive integer r . Then the following statements hold.

- There exists a field of order q .
- Any two fields of order q are isomorphic.
- Let K be a field of order q . The multiplicative group K^\times of non-zero elements of K is a cyclic group of order $q - 1$.
- Let K be a field of order q . The elements of K are the roots of $x^q - x \in \mathbb{F}_p[x]$.
- A field of order p^r contains a field of order $p^k \iff k|r$
- The irreducible factors of $x^q - x$ over \mathbb{F}_p are the irreducible polynomials in $\mathbb{F}_p[x]$ whose degree divides r .
- The splitting field of $x^q - x$ has q elements.
- \mathbb{F}_q is a stem field and a splitting field of any irreducible polynomial $P \in \mathbb{F}_p$ of degree n .

Frobenius homomorphism

Let K be a field, $\text{ch}(K) = p > 0$. There exists a homomorphism $\varphi : K \rightarrow K$, s.t. $\varphi(x) = x^p$. This is Frobenius homomorphism.

- The group of automorphisms over \mathbb{F}_{p^r} over \mathbb{F}_p is cyclic and is generated by the Frobenius map.

Separability

- **Separable polynomial:** A irreducible polynomial $P \in K[X]$ is called separable if $\gcd(P, P') = 1$.
- **Degree of separability:** $\deg_{\text{sep}} P = \deg Q$ for some $P(X) = Q(X^p)$
- **Degree of inseparability:** $\deg_i P = \frac{\deg P}{\deg Q}$
- **Purely inseparable polynomial:** P is purely inseparable if $\deg_i P = \deg P$. Also if P is purely inseparable $P = X^{p^r} - a$
- **Separable element:** If L/K is an algebraic extension, then $\alpha \in L$ is called separable if its minimal polynomial over K is separable. And vice versa.
- If $\alpha \in K$ is separable then $|\text{Hom}(K(\alpha), \overline{K})| = \deg P_{\min}(\alpha, K)$
- **Separable degree:** For L/K , we have $[L : K]_{\text{sep}} = |\text{Hom}_K(K(\alpha), \overline{K})|$. Inseparable degree is degree of extension divided by separable degree.
- **Separable extension:** L is separable over K if $[L : K]_{\text{sep}} = [L : K]$.
 - If $\text{ch}(K) = 0$ then any extension of K is separable.
 - If $\text{ch}(K) = p$ then pure inseparable extension has degree p^r with degree of inseparability p^r
- Separable degrees obey the multiplicative property.
- TFAE
 - L is separable over K
 - Any element of L is separable over K
 - $L = K(\alpha_1, \alpha_2, \dots, \alpha_n)$, where each α_i is separable over K .
 - $L = K(\alpha_1, \alpha_2, \dots, \alpha_n)$, then α_i is separable over $K(\alpha_1, \dots, \alpha_{i-1})$.
- **Separable closure:** $L^{\text{sep}} = \{x \mid x \text{ separable over } K\}$

Multilinear map

For a module M over ring A . A function L from $M^r = \underbrace{M \times M \times \dots \times M}_{r \text{ times}}$ into

A is called multilinear if $L(\alpha_1, \dots, \alpha_r)$ is linear as a function of each α_i when the other α_j are fixed.

Tensor product

Consider a ring A and two A -modules, M, N . The tensor product is denoted as $M \otimes_A N$ is an A -module along with a A -bilinear map, $\varphi : M \times N \rightarrow M \otimes_A N$ which a “universal property”.

Universal property of tensor product:

For a A -module P , if for an A -bilinear map, $f : M \times N \rightarrow P$, then $\exists!$ homomorphism \tilde{f} of A -modules s.t. $f = \tilde{f} \circ \varphi$

$$\begin{array}{ccc} M \times N & \xrightarrow{\varphi} & M \otimes_A N \\ & \searrow f & \downarrow \tilde{f} \\ & & P \end{array}$$

- Commutativity of tensor product $M \otimes_A N \cong N \otimes_A M$
- $A \otimes_A M \cong M$
- The basis for the tensor product of free modules is the tensor product of their individual basis elements.
- The tensor product is associative.

Base change theorem: For a ring A, B an A -algebra, M an A -module and N a B -module. Then we have the following bijection

$$\text{Hom}_A(M, N) \leftrightarrow \text{Hom}_B(B \otimes_A M, N)$$

- For I an ideal of a ring A and M an A -module we have, $A/I \otimes_A M \cong M/IM$

Chinese remainder theorem

Comaximal ideals: Two ideals of a ring are called comaximal (or coprime) if their sum gives the ring itself.

- If I, J are comaximal then $IJ = I \cap J$
- If I_1, \dots, I_k comaximal w.r.t J then $\prod_{i=1}^k I_i$ is also relatively prime with J .
- If I, J are comaximal then so are I^m, J^n for any m, n .

Chinese remainder theorem: For a ring A , consider two comaximal ideals I, J , then $\forall a, b \in R, \exists x \in A$ s.t. $x \equiv a \pmod{I}$ and $x \equiv b \pmod{J}$

Generalized Chinese remainder theorem: For a ring A , let I_1, \dots, I_n be ideals of the ring A . Consider the map $\pi : A \rightarrow A/I_1 \times \dots \times A/I_n$ defined as $\pi(a) = (a \pmod{I_1}, \dots, a \pmod{I_n})$. Then $\ker \pi = I_1 \cap \dots \cap I_n$, i.e. it is surjective iff I_1, \dots, I_n are pairwise comaximal. If π is a surjection we have,

$$A / \bigcap I_k \cong A / \prod I_k \cong \prod (A / I_k)$$

Structure of finite algebras

Let A be a finite K -algebra then,

- There are only finitely many maximal ideals in A .
- For finitely many maximal ideals m_i . Let $J = m_1 \cap \dots \cap m_r$. Then $J^n = 0$ for some n .
- $A \cong A/m_1^{n_1} \times \dots \times A/m_r^{n_r}$ for some (not necessarily unique) n_1, \dots, n_r .

Reduced K-Algebra: If it has no nilpotent elements.

Local ring: If it has only one maximal ideal. A non zero ring in which every element is either a unit or nilpotent is local.

Further results on separability

Let L be a finite extension over K then the following hold,

- L is separable $\iff L \otimes_K \bar{K}$ is reduced.
- L is purely inseparable $\iff L \otimes_K \bar{K}$ is local.
- L is separable $\iff \forall$ algebraic extensions $\Omega, L \otimes_K \Omega$ is reduced.
- L is purely inseparable $\iff \forall$ algebraic extensions $\Omega, L \otimes_K \Omega$ is local.
- If L is separable then the map $\varphi : L \otimes_K \bar{K} \rightarrow \bar{K}^n$ defined as $\varphi(l \otimes k) = (k\varphi_1(l), \dots, k\varphi_n(l))$ (where φ_i are distinct homomorphisms from L to \bar{K}), is an isomorphism.
- Let L be a finite separable extension of K then it has only finitely many sub extensions.

Primitive element theorem

There exists $\alpha \in L$ s.t. $L = K(\alpha)$ whenever L is finite and separable.

Normal extensions

A normal extension of K is a splitting field of a family of polynomials in $K[X]$.

TFAE for an extension L of K ,

- $\forall x \in L, P_{\min}(x, K)$ splits in L .
- L is a normal extension.
- All homomorphisms from L to \bar{K} have the same image.
- The group of automorphisms, $\text{Aut}(L/K)$ acts transitively on $\text{Hom}_K(L, \bar{K})$.

Some properties of normal extensions,

- $K \subset L \subset M$, if M is normal over K then it is normal over L , but L need not be normal over K .
- Extensions with degree 2 are normal.

Galois extensions

A field extension that is both normal and separable is called a Galois extension.

- For a finite extension L over K the number of automorphisms $|\text{Aut}(L/K)| \leq [L : K]$. Equality holds iff L is a Galois' extension.

If L is normal over K then,

- Isomorphism of sub extensions extend to automorphisms of L .
- $\text{Aut}(L/K)$ acts transitively on the roots of any irreducible polynomial in $K[X]$.
- If $\text{Aut}(L/K)$ fixes $x \notin K$. Then x is purely inseparable.

Galois groups

If L is a Galois extension, $G = \text{Gal}(L/K) = \text{Aut}(L/K)$ is called the Galois group of the extension.

- $L^{\text{Gal}(L/K)} = K$, (i.e. the set of invariants in L with the action of the Galois group is equal to K).
- Let L be a field and G a subgroup of $\text{Aut}(L)$, then
 - If all orbits of G are finite, then L is a Galois extension of L^G .
 - If order of G is finite then, $[L, L^G] = n$ and G is a Galois group.

The Fundamental theorem of Galois theory

Let L/K be a Galois extension, and $\text{Aut}(L/K) = \text{Gal}(L/K)$ is its Galois group.

- If L is finite over K , then for a intermediate field F and a subgroup $H \subset \text{Gal}(L/K)$ we have the following correspondence,
 - $F \rightarrow \text{Gal}(L/F)$
 - $H \rightarrow L^H$
- F is Galois over $K \iff g(F) = F, \forall g \in \text{Gal}(L/K) \iff \text{Gal}(L/F) \trianglelefteq \text{Gal}(L/K)$

Discriminant

For a polynomial with roots x_i , the discriminant is $\Delta = \prod_{i < j} (x_i - x_j)^2$. For $\text{Gal}(P) \subset S_n$,

- Δ is preserved by any permutation.
- $\sqrt{\Delta}$ is preserved only by even permutations
- $G \subset A_n \iff \sqrt{\Delta} \in K$

Cyclotomic polynomials

Let $P_n = X^n - 1$ where n is coprime to the characteristic of a finite field K . P_n has n distinct roots which form a cyclic multiplicative subgroup $\mu_n \subset \bar{K}^\times$. Let μ_n^* denote the set of **primitive roots of unity** (no roots of degree $< n$).

- $|\mu_n^*| = \varphi(n)$

Cyclotomic polynomials: $\Phi_n = \prod_{\alpha \in \mu_n^*} (X - \alpha) \in \bar{K}[X]$.

- $P_n = \prod_{d|n} \Phi_d$.
- Φ_n has coefficients in prime fields.
- If $\text{ch}(K) = 0$ then $\Phi_n \in \mathbb{Z}[X]$, else if $\text{ch}(K) = p$, we have Φ_n is the reduction mod p of the n^{th} cyclotomic polynomial over \mathbb{Z} .
- If $\text{ch}(K) = 0$, then Φ_n is irreducible over $\mathbb{Z}[X]$.

Consider L , splitting field of K

- The splitting field of P_n over K is $K(\zeta)$ where ζ is a root of Φ_n .
- All $g \in \text{Gal} L/K$ acts as $\zeta \rightarrow \zeta^{a^g}, (a^g, n) = 1$.
- $\text{Gal}(L/K)$ injects into $\mathbb{Z}/n\mathbb{Z}^\times$ and this is an isomorphism when Φ_n is irreducible over K .