

Darcy's law

The water flux in a porous medium is proportional to the gradient of **hydraulic head (total water potential)**, with the proportionality given by the **hydraulic conductivity (depending on soil moisture)**

how tightly water is held by:
(1) the soil (matric potential) +
(2) elevation head (gravitational potential)

how easily water moves through the soil;
in unsaturated conditions shrinks rapidly
as the soil dries

Neglecting lateral fluxes....

$$q = -K(\theta) \left(\frac{\partial H}{\partial z} \right)$$

specific discharge (m/s)

hydraulic conductivity (m/s)

hydraulic gradient (adimensional)

Darcy's law

The water flux in a porous medium is proportional to the gradient of **hydraulic head (total water potential)**, with the proportionality given by the **hydraulic conductivity (depending on soil moisture)**

how tightly water is held by the soil
(matric potential) + elevation head
(gravitational potential)

how easily water moves through the soil;
in unsaturated conditions shrinks rapidly
as the soil dries

Neglecting lateral fluxes....

by discretizing the soil column, the vertical head
gradient at interface between two adjacent
layers is approximated by a finite difference

$$q = -K(\theta) \left(\frac{\partial H}{\partial z} \right) \approx -K(\theta) \left(\frac{\Delta H}{\Delta z} \right)$$

specific discharge (m/s)

hydraulic conductivity (m/s)

soil moisture (m³/m³)

hydraulic gradient

$$q_{l+1/2} = -K_{l+1/2}^{effective} \left(\frac{H_{l+1} - H_l}{z_{l+1} - z_l} \right), \quad H_l = \frac{\psi_l}{\rho_w g} + z_l$$

the conductivity at the interface could be approximated using the harmonic mean of Ks values from the two adjacent layers

soil water potential ($MPa = kgm^{-1}s^{-2}$)

gravitational potential (m)

water density (kgm^{-3})

gravitational acceleration (ms^{-2})

One possible simplification, based on Liu et al. 2017 (PNAS) and, is to calculate the upward flux by the simplified darcy's law equation and the downward flux by using the bucket model logic of percolation.

[(**SPAC**: Liu et al., 2017 - PNAS); (**JULES**: Best et al., 2011 - Geosci. Model Dev)]

$$q_{l+1/2}^{cap} = -K_{l+1/2}^{effective} \left(\frac{\psi_{l+1} - \psi_l}{p_w g \Delta z} + 1 \right)$$

To guarantee only upward flux:

$$q_{l+1/2}^{cap} = \max(q_{l+1/2}^{cap}, 0)$$



