

# Modeled changes on functional diversity and carbon storage driven by drought in the Amazon forest: a plant-trait vs. PFT-based comparison

**Authors:** Bianca Fazio Rius<sup>1,2</sup>, João Paulo Darella Filho<sup>1,3</sup>, David Montenegro Lapola<sup>1</sup>

## **Institutional affiliations:**

<sup>1</sup>Earth System Science Lab, Center for Meteorological and Climatic Research Applied to Agriculture, University of Campinas – UNICAMP, Campinas - SP, Brazil

<sup>2</sup>Post-Graduate Programme in Ecology, Biology Institute, University of Campinas – UNICAMP, Campinas - SP, Brazil

<sup>3</sup>Post-Graduate Programme in Ecology, Evolution and Biodiversity, Biology Institute, São Paulo State University – UNESP, Rio Claro - SP, Brazil

## **Contact Information:**

Bianca Fazio Rius  
Earth System Science Lab, Center for Meteorological and Climatic Research Applied to Agriculture, University of Campinas – UNICAMP, Campinas, 1308-886 - SP, Brazil  
Email: [biancafaziorius@gmail.com](mailto:biancafaziorius@gmail.com)  
Tel: +55 11 99729 2917

**Keywords:** trait-based modeling, climate change, carbon allocation, functional composition, functional richness, functional evenness, tropical forest, biomass

## **Introduction**

The possible modification in Amazon forest's ability to absorb and store carbon due to climate change is permeated with uncertainties (Finegan et al., 2015), and the role of functional diversity on this ecosystem process is poorly explored (Poorter et al., 2015; Sakschewski et al., 2016; Sitch et al., 2008). Dynamic global vegetation models (DGVMs) have been widely used to explore the question from a biogeochemical perspective (Cramer et al., 2001; Scheiter, Langan, & Higgins, 2013) providing substantial contribution to our current knowledge of the Amazon forest ecology and resilience (Díaz & Cabido, 1997; Prentice et al., 2007; Scheiter et al., 2013). For instance, the projected possibility of a large-scale degradation