

# Copyright Notice

These slides are distributed under the Creative Commons License.

[DeepLearning.ai](#) makes these slides available for educational purposes. You may not use or distribute these slides for commercial purposes. You may make copies of these slides and use or distribute them for educational purposes as long as you cite [DeepLearning.AI](#) as the source of the slides.

For the rest of the details of the license, see <https://creativecommons.org/licenses/by-sa/2.0/legalcode>

# W2 Lesson 1



DeepLearning.AI

# Describing Distributions

---

## Expected value

# Mean: Example


Age:

0

1

2

3



# Mean: Example


Age:

0

1


2

3



# Mean: Example

Age: 0      1      2      3



$$\begin{array}{r} 0 + 0 + 0 + 1 + 1 + 2 + 2 + 2 + 3 \\ \hline 10 \end{array}$$

$$= \frac{13}{10}$$

$$= 1.3$$

# Children in a Room


Age:

0

1

2

3



$$\frac{0 + 0 + 0 + 1 + 1 + 2 + 2 + 2 + 3}{10}$$

$$= \frac{13}{10} = 1.3$$

$$= \frac{3 \cdot 0 + 2 \cdot 1 + 4 \cdot 2 + 1 \cdot 3}{10}$$

Weighted average

$$= \frac{3}{10} \cdot 0 + \frac{2}{10} \cdot 1 + \frac{4}{10} \cdot 2 + \frac{1}{10} \cdot 3 = 1.3$$

# Expected Value: Motivation Example 1

You play a game with a friend



You win 10 dollars



You win nothing

Game cost:



Do you play the game?

What is the maximum amount of money you would pay to play this game?

# Expected Value: Motivation Example 1

You play a game with a friend



You win 10 dollars



You win nothing


Game cost:

\$5

Long term:  $0.5 \cdot \$10 + 0.5 \cdot \$0 = \$5$  

You expect to win \$5 on average  
 $E[X] = 5$

# Expected Value: Motivation Example 1



# Expected Value: Motivation Example 2

Play another game



Flip three coins. For each heads you win \$1


What is the maximum amount of money you would pay to play this game?

# Expected Value: Motivation Example 2

$X$ : Number of heads

$$\mathbb{E}[X] = 1.5$$

$$\mathbb{E}[X] = \frac{1}{8} \cdot 0 + \frac{3}{8} \cdot 1 + \frac{3}{8} \cdot 2 + \frac{1}{8} \cdot 3 = 1.5$$




# Expected Value: Discrete Case

$X$  a discrete  
random variable


PMF of  $X$   
 $p_X(x) = \mathbf{P}(X = x)$

$$\mathbb{E}[X] = \sum_x x p_X(x)$$



# Expected Value




# Expected Value





# Expected Value - Continuous



# Expected Value - Continuous



# Expected Value - Continuous



# Expected Value - Continuous

Discrete random variables

$$\mathbb{E}[X] = \sum_x x p_X(x)$$

Weighted using PMF

Continuous random variables

$$\int_{-\infty}^{\infty} x \cdot f_X(x) dx$$


Integrals

Weighted using PDF


# Expected Value

Waiting Time
15 min
32 min
58 min
1 min
47 min
14 min
37 min
8 min
29 min
55 min
...


Average = 20.833



Probability




# Expected Value: Uniform Distribution




# Expected Value: Uniform Distribution


$$\text{Average} = \frac{a + b}{2}$$




# Expected Value




# Expected Value: Common Misconception



# Expected Value: Common Misconception



# Expected Value: Common Misconception



# Expected Value

- $\mathbb{E}[X]$
- Mean / Balancing point
- Defined for discrete and continuous random variables
- Weighted average of the PMF / PDF



DeepLearning.AI


# Describing Distributions

---

**Other measures of central  
tendency**


# Median: Motivation

- In the 1980s, the starting salary for a geography graduate at the University of North Carolina was \$250,000.
  - For the rest of the country, the starting salary for a geography graduate was \$22,000.
  - Why?
    1. The program was really good
    2. The university had great connections
    3. One student made lots of money
- 



Michael Jordan


# Outliers




# Outliers




# Median




# Median




# Mode




# Mode




# Mode




# Mode: Multimodal Distribution




# Mean, Median and Mode in Binomial Distribution



# Mean, Median and Mode in Binomial Distribution



# Mean, Median and Mode in Normal Distribution






DeepLearning.AI

# Describing Distributions

---

## Expected value of a function

# Expected Value of a Function




$$\mathbb{E}[X] = x_1p(x_1) + x_2p(x_2) + x_3p(x_3) + x_4p(x_4)$$

$$E[g(X)] = g(x_1)p(x_1) + g(x_2)p(x_2) + g(x_3)p(x_3) + g(x_4)p(x_4)$$

# Expected Value of a Function

Probability:  $1/6$     $1/6$     $1/6$     $1/6$     $1/6$     $1/6$


Roll:    1            2            3            4            5            6



Square:    1            4            9            16            25            36


# Expected Value of a Function

Probability




# Expected Value of a Function

$$\frac{1 + 4 + 9 + 16 + 25 + 36}{6} = \frac{91}{6}$$



# Expected Value of a Function


$$\frac{1 + 4 + 9 + 16 + 25 + 36}{6} = \frac{91}{6} = \frac{1^2}{6} + \frac{2^2}{6} + \frac{3^2}{6} + \frac{4^2}{6} + \frac{5^2}{6} + \frac{6^2}{6}$$
$$= \mathbb{E}[X^2]$$



# Expectation of Linear Function

Probability:  $1/6$     $1/6$     $1/6$     $1/6$     $1/6$     $1/6$

Roll:      1      2      3      4      5      6




Double:      2      4      6      8      10      12

Wins      2 - 5      4 - 5      6 - 5      8 - 5      10 - 5      12 - 5

# Expectation of Linear Function

Probability:  $1/6$     $1/6$     $1/6$     $1/6$     $1/6$     $1/6$

Roll:    1            2            3            4            5            6




Double:    2            4            6            8            10          12

Wins        -3           -2           -1           0           1           2


# Expected Value of a Function

Probability




# Expected Value of a Function

$$\frac{-3 + -2 + -1 + 0 + 1 + 2}{6} = \frac{-3}{6} = -0.5$$




# Expected Value of a Function

$$\frac{(2 \cdot 1 - 5) + (2 \cdot 2 - 5) + (2 \cdot 3 - 5) + (2 \cdot 4 - 5) + (2 \cdot 5 - 5) + (2 \cdot 6 - 5)}{6} = \frac{-3}{6} = -0.5$$




# Expected Value of a Function

$$\frac{(2 \cdot 1) + (2 \cdot 2) + (2 \cdot 3) + (2 \cdot 4) + (2 \cdot 5) + (2 \cdot 6) + 6 \cdot (-5)}{6} = \frac{-3}{6} = -0.5$$



# Expected Value of a Function

$$\frac{(2 \cdot 1) + (2 \cdot 2) + (2 \cdot 3) + (2 \cdot 4) + (2 \cdot 5) + (2 \cdot 6)}{6} + (-5) = \frac{-3}{6} = -0.5$$



# Expected Value of a Function

$$\mathbb{E}[2 \cdot X + (-5)] = \frac{2 \cdot (1 + 2 + 3 + 4 + 5 + 6)}{6} + (-5) = \frac{-3}{6} = -0.5$$

$= 2 \cdot \mathbb{E}[X] + (-5)$



In general:

$$\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$$

$$\mathbb{E}[aX] = a\mathbb{E}[X]$$

$$\mathbb{E}[b] = b$$



DeepLearning.AI

# Describing Distributions

---

## Sum of expectations

# Sum of Expectations

You play a game:

Flip a coin. If heads you win \$ 1, else you win nothing.

Then roll a die. You win the amount you roll.




Win \$1



Win nothing

Win      \$1      \$2      \$3      \$4      \$5      \$6



What are your expected winnings for the game?

# Sum of Expectations




Win \$1



Win nothing

Win      \$1      \$2      \$3      \$4      \$5      \$6



$$\mathbb{E}[X_{coin}] = \$0.5$$

$$\mathbb{E}[X_{dice}] = \$3.5$$

$$\mathbb{E}[X] = \mathbb{E}[X_{coin}] + \mathbb{E}[X_{dice}] = \$0.5 + \$3.5 = \$4$$

In general:  $\mathbb{E}[X_1 + X_2] = \mathbb{E}[X_1] + \mathbb{E}[X_2]$

# Sum of Expectations



Expected number of  
correct assignments?



# Sum of Expectations





1

Expected number of  
correct assignments?





8 billion people


# Sum of Expectations




# Sum of Expectations




# Sum of Expectations



# Sum of Expectations



# Sum of Expectations





Average  
1

Correct


3  
1  
1  
0  
0  
1

6

Aisha	Beto	Cameron
Aisha	Cameron	Beto
Beto	Aisha	Cameron
Beto	Cameron	Aisha
Cameron	Aisha	Beto
Cameron	Beto	Aisha





# Sum of Expectations




$\frac{1}{3}$   
 $\frac{1}{3}$   
 $\frac{1}{3}$

$$\begin{aligned}\mathbb{E}[\text{Matches}] &= \mathbb{E}[A] + \mathbb{E}[B] + \mathbb{E}[C] \\ &= \frac{1}{3} + \frac{1}{3} + \frac{1}{3} \\ &= 1\end{aligned}$$



Average  
1

# Sum of Expectations



Expected number = ?



8 billion people

# Sum of Expectations



$$\mathbb{E} [\text{Matches}] = \mathbb{E} [X_{\text{person}_1}] + \mathbb{E} [X_{\text{person}_2}] + \dots + \mathbb{E} [X_{\text{person}_n}]$$

$n$  people ( $n = 8$  billion)

$$= \frac{1}{n} + \frac{1}{n} + \dots + \frac{1}{n}$$

In general:

$$\mathbb{E} [X_1 + X_2 + \dots + X_n] = \mathbb{E} [X_1] + \mathbb{E} [X_2] + \dots + \mathbb{E} [X_n]$$

$$= n \cdot \frac{1}{n} = 1$$



DeepLearning.AI

# Describing Distributions

---

## Variance

# Variance Motivation: Fair Price To Play the Game

You play a game with a friend



You win 1 dollar



You lose 1 dollar

What is the fair amount of money to pay to play this game?

# Variance Motivation: Fair Price To Play the Game

You play a game with a friend



You win 1 dollar



You lose 1 dollar

Game cost:

\$0

What is the fair amount of money to pay to play this game?

# Variance Motivation: Fair Price To Play the Game

You play a game with a friend



You win 100 dollars



You lose 100 dollars

What is the fair amount of money to pay to play this game?

# Variance Motivation: Fair Price To Play the Game

You play a game with a friend



You win 100 dollars



You lose 10

Game cost:


Variance!

0

What is the fair amount of money to pay to play this game?


# Variance Motivation: Measuring Spread

Probability




# Variance Motivation: Measuring Spread

$X_1$  = amount of money gained in game 1




# Variance Motivation: Measuring Spread


$X_2$  = amount of money gained in game 2



# Variance Motivation: Measuring Spread




# Variance Motivation: Measuring Spread



# Variance Motivation: Measuring Spread

Deviation  
 $x - \mathbb{E}[X]$



Squared  
deviation  
 $(x - \mathbb{E}[X])^2$


$$\mathbb{E}[X_1] = 0$$



Expected  
squared  
deviation  
 $\mathbb{E}[(x - \mathbb{E}[x])^2]$

$$\mathbb{E}[X_2] = 0$$

# Variance Motivation: Measuring Spread



# Variance Formula

$$\text{Variance} = \mathbb{E} [(X - \mathbb{E}[X])^2]$$

1. Find X's mean
2. Find the deviation from that mean for every value of X
3. Square those deviations
4. Average those squared deviations

“Average squared deviation”

# Variance Motivation: Centering With Mean

Game 1



You win 2 dollars



You lose 2 dollars

Game 2



You win 3 dollars



You lose 1 dollar

Which of these games has greater variance?

**Hint:** Think of the spread

# Variance Motivation: Centering With Mean

Game 1



You win 2 dollars



You lose 2 dollars

Game 2



You win 3 dollars



You lose 1 dollar

They have the same variance

# Variance Motivation: Centering With Mean

## Game 1



You win 2 dollars

$$\mathbb{E}[X_1] = \frac{1}{2} \cdot (-2) + \frac{1}{2} \cdot 2 = 0$$

You lose 2 dollars

## Game 2



You win 3 dollars

$$\mathbb{E}[X_2] = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 3 = 1$$

You lose 1 dollar

# Variance Motivation: Centering With Mean

## Game 1



You win 2 dollars

You lose 2 dollars

$$\mathbb{E}[X_1] = 0$$

$$\frac{1}{2}(-2 - \mathbb{E}[X_1])^2 + \frac{1}{2}(2 - \mathbb{E}[X_1])^2 = 4$$

## Game 2



You win 3 dollars

You lose 1 dollar

Different price, but same spread

$$\mathbb{E}[X_2] = 1$$

$$\frac{1}{2}(-1 - \mathbb{E}[X_2])^2 + \frac{1}{2}(3 - \mathbb{E}[X_2])^2 = 4$$

# Variance Formula

$$\begin{aligned}Var(X) &= \mathbb{E}[(X - \mathbb{E}[X])^2] \\&= \mathbb{E}[X^2] - \mathbb{E}[X]^2\end{aligned}$$

# Variance Formula

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2] = \mathbb{E}[X^2 - 2\mathbb{E}[X]X + \mathbb{E}[X]^2]$$

$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$= \mathbb{E}[X^2] - \mathbb{E}[2\mathbb{E}[X]X] + \mathbb{E}[\mathbb{E}[X]^2]$$

$$= \mathbb{E}[X^2] - 2\mathbb{E}[X]\mathbb{E}[X] + \mathbb{E}[X]^2$$

$$= \mathbb{E}[X^2] - 2\mathbb{E}[X]^2 + \mathbb{E}[X]^2$$



$$= \mathbb{E}[X^2] - \mathbb{E}[X]^2$$

$\mathbb{E}[\text{constant} \cdot X] = \text{constant} \cdot \mathbb{E}[X]$

$\mathbb{E}[X]$  is a constant

$\mathbb{E}[\text{constant}] = \text{constant}$

# Properties of the Variance

Probability:	1/6	1/6	1/6	1/6	1/6	1/6
Roll:	1	2	3	4	5	6
						
Win Double:	\$2	\$4	\$6	\$8	\$10	\$12
Net Amount:	-\$3	-\$1	\$1	\$3	\$5	\$7

# Properties of the Variance


changes the spread

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

doesn't change the spread


Variance: "Average squared deviation"

Dice roll is random variable:  $X$



Net winnings is random variable:  $Y = 2X - 5$

$$\text{Var}(Y) = \text{Var}(2X - 5) = 4\text{Var}(X)$$





DeepLearning.AI

# Describing Distributions

---

## Standard deviation

# Standard Deviation

$$\text{Var}(X) = \mathbb{E}[(X - \mu)^2] = \mathbb{E}[X^2] - \mathbb{E}[X]^2$$


Say  $X$  is measured in meters.

Then  $\mathbb{E}[X]$  is measured in meters.

Then  $\text{Var}(X)$  is measured in meters<sup>2</sup>.

Then  $\sqrt{\text{Var}(X)}$  is measured in meters.

Let's call  $std(X) = \sqrt{\text{Var}(X)}$ , the *standard deviation* of  $X$




# Normal Distribution: 68-95-99.7 Rule

Parameters:

- $\mu$ : center of the bell
  - $\sigma$ : spread of the bell
- $$\rightarrow X \sim \mathcal{N}(\mu, \sigma^2)$$

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$




# Normal Distribution: 68-95-99.7 Rule

Parameters:

- $\mu$ : center of the bell
- $\sigma$ : spread of the bell

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$




# Normal Distribution: 68-95-99.7 Rule

Parameters:

- $\mu$ : center of the bell
- $\sigma$ : spread of the bell

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$




# Normal Distribution: 68-95-99.7 Rule

Parameters:

- $\mu$ : center of the bell
- $\sigma$ : spread of the bell

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$






DeepLearning.AI

# Describing Distributions

---

## Sum of Gaussians

# Sum of Gaussians: an Example





$$T \sim \mathcal{N}(10, 2^2)$$

$$L \sim \mathcal{N}(5, 1^2)$$



$$R = T + L$$

# Sum of Gaussians




# Sum of Gaussians

Sample each variable 10000 times



# Sum of Gaussians

$$R = T + L$$



$R$  is still Gaussian!

Expectation is linear

$$\begin{aligned}\mu_R &= \mathbb{E}[R] = \mathbb{E}[T + L] = \mathbb{E}[T] + \mathbb{E}[L] \\ &= \mu_T + \mu_L = 10 + 5 = 15\end{aligned}$$

$$\begin{aligned}\sigma_R^2 &= \text{Var}(R) = \text{Var}(T + L) \\ &= \text{Var}(T) + \text{Var}(L) = \sigma_T^2 + \sigma_L^2\end{aligned}$$


$$= 4 + 1 = 5$$

# Sum of Gaussians

$$R = T + L$$


$R$  is still Gaussian!

$$R = (T + L) \sim \mathcal{N}(10 + 5, 4 + 1)$$



# Sum of Gaussians

$$R = T + L$$



In general:  $W = \textcolor{teal}{a}X + \textcolor{teal}{b}Y$

Independent  $\begin{cases} X \sim \mathcal{N}(\mu_X, \sigma_X^2) \\ Y \sim \mathcal{N}(\mu_Y, \sigma_Y^2) \end{cases}$

$$\rightarrow W \sim \mathcal{N} \left( a\mu_x + b\mu_y, a^2\sigma_x^2 + b^2\sigma_y^2 \right)$$




DeepLearning.AI

# Describing Distributions

---

## Standardizing a Distribution

# Everything Is Nicer When the Mean Is 0



$$X \rightarrow X - \mu$$

# Everything Is Nicer When the Mean Is 0

Why?


$$\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$$

$$\mathbb{E}[X - \mu] = \mathbb{E}[X] - \mathbb{E}[\mu]$$


$$= \mathbb{E}[X] - \mu$$

$$= 0$$

# Everything Is Nicer When the Standard Deviation Is 1



# Everything Is Nicer When the Standard Deviation Is 1



Mean = 0

Standard deviation = 1

$$X \rightarrow \frac{X}{\sigma}$$

# Everything Is Nicer When the Standard Deviation Is 1

Why?

$$Var(cX) = \mathbb{E}[(cX)^2] - \mathbb{E}[cX]^2$$

$$= \mathbb{E}[c^2X^2] - (c\mathbb{E}[X])^2$$

$$= c^2\mathbb{E}[X^2] - c^2\mathbb{E}[X]^2$$

$$= c^2(\mathbb{E}[X^2] - \mathbb{E}[X]^2)$$

$$= c^2Var(X)$$


$$Var\left(\frac{X}{\sigma}\right) = \frac{1}{\sigma^2}Var(X)$$

$$std\left(\frac{X}{\sigma}\right) = \frac{1}{\sigma}std(X)$$

$$= \frac{\sigma}{\sigma}$$

$$= 1$$

# Standardize a Distribution





DeepLearning.AI

# Describing Distributions

---

**Skewness and Kurtosis:  
Moments of a Distribution**

# Moments of a Distribution


$$\mathbb{E}[X] = \frac{1}{3}(-2) + \frac{1}{6}(0) + \frac{1}{2}(1)$$

$$\mathbb{E}[X^2] = \frac{1}{3}(-2)^2 + \frac{1}{6}(0)^2 + \frac{1}{2}(1)^2$$

$$\mathbb{E}[X^3] = \frac{1}{3}(-2)^3 + \frac{1}{6}(0)^3 + \frac{1}{2}(1)^3$$

...

$$\mathbb{E}[X^k] = \frac{1}{3}(-2)^k + \frac{1}{6}(0)^k + \frac{1}{2}(1)^k$$



# Moments of a Distribution

$$\mathbb{E}[X] = p_1x_1 + p_2x_2 + \cdots + p_nx_n$$

$$\mathbb{E}[X^2] = p_1x_1^2 + p_2x_2^2 + \cdots + p_nx_n^2$$


$$\mathbb{E}[X^3] = p_1x_1^3 + p_2x_2^3 + \cdots + p_nx_n^3$$

$$\mathbb{E}[X^4] = p_1x_1^4 + p_2x_2^4 + \cdots + p_nx_n^4$$

...

$$\mathbb{E}[X^k] = p_1x_1^k + p_2x_2^k + \cdots + p_nx_n^k$$

Random variable  $X$





DeepLearning.AI

# Describing Distributions

---

**Skewness and Kurtosis:  
Skewness**

# Lottery vs Insurance



Ticket: \$1  
Jackpot: \$100

Lottery

You **win** \$99 with 1% probability

You **lose** \$1 with 99% probability



Cost: \$1  
Crash Reparation: \$100

Car insurance

You **win** \$1 with 99% probability


You **lose** \$99 with 1% probability

# Lottery vs Insurance



Ticket: \$1  
Jackpot: \$100

Lottery



Cost: \$1  
Crash Reparation: \$100

Car insurance





Ticket: \$1  
Jackpot: \$100

## Lottery

$$\mathbb{E}[X_1] = -1 \cdot 0.99 + 99 \cdot 0.01 = 0$$

Lose 1  
With probability 0.99

Win 99  
With probability 0.01




Cost: \$1  
Crash Reparation: \$100


## Car insurance

$$\mathbb{E}[X_2] = -99 \cdot 0.01 + 1 \cdot 0.99 = 0$$

Lose 99  
With probability 0.01

Win 1  
With probability 0.99






Ticket: \$1  
Jackpot: \$100

Lottery

$$\mathbb{E}[X_1] = -1 \cdot 0.99 + 99 \cdot 0.01 = 0$$

$$Var(X_1) = (-1)^2 \cdot 0.99 + (99)^2 \cdot 0.01 = 99$$




Cost: \$1  
Crash Reparation: \$100

Car insurance

$$\mathbb{E}[X_2] = -99 \cdot 0.01 + 1 \cdot 0.99 = 0$$

$$Var(X_2) = (-99)^2 \cdot 0.01 + (1)^2 \cdot 0.99 = 99$$





Ticket: \$1  
Jackpot: \$100

Lottery

$$\mathbb{E}[X_1] = 0$$

$$Var(X_1) = 99$$



Cost: \$1  
Crash Reparation: \$100



Car insurance

Same expectation  
Same variance

How to tell them apart?

$$\mathbb{E}[X_2] = 0$$

$$Var(X_2) = 99$$






Ticket: \$1  
Jackpot: \$99

Lottery

$$\mathbb{E}[X_1] = 0$$

$$Var(X_1) = 99$$

$$\mathbb{E}[X_1^2] - \mathbb{E}[X_1]^2 = 99$$



Cost: \$1  
Crash Reparation: \$99

Car insurance


Same expectation  
Same variance

How to tell them apart?

$$\mathbb{E}[X_2] = 0$$

$$Var(X_2) = 99$$

$$\mathbb{E}[X_2^2] - \mathbb{E}[X_2]^2 = 99$$





Ticket: \$1  
Jackpot: \$99


## Lottery

$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[X_1^2] = 99$$

$$\mathbb{E}[X_1^3] = (-1)^3 \cdot 0.99 + (99)^3 \cdot 0.01$$

$$= 9702$$



Cost: \$1  
Crash Reparation: \$99

## Car insurance

Same expectation  
Same variance


How to tell them apart?

$$\mathbb{E}[X_2] = 0$$

$$\mathbb{E}[X_2^2] = 99$$

$$\mathbb{E}[X_2^3] = (1)^3 \cdot 0.99 + (-99)^3 \cdot 0.01$$

$$= -9702$$






Ticket: \$1  
Jackpot: \$99

Lottery

$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[X_1^2] = 99$$

$$\mathbb{E}[X_1^3] = \text{Large positive value}$$



Cost: \$1  
Crash Reparation: \$99

Car insurance


Same expectation  
Same variance

How to tell them apart?

$$\mathbb{E}[X_2] = 0$$

$$\mathbb{E}[X_2^2] = 99$$

$$\mathbb{E}[X_2^3] = \text{Large negative value}$$



# Skewness

$$\mathbb{E}[X^3]$$

Almost...

Need to standardize...

# Skewness

$$\text{Skewness} = \mathbb{E} \left[ \left( \frac{X - \mu}{\sigma} \right)^3 \right]$$

# Skewness



Positively  
Skewed



Not  
Skewed



Negatively  
Skewed



$$E \left[ \left( \frac{X - \mu}{\sigma} \right)^3 \right] > 0$$

$$E \left[ \left( \frac{X - \mu}{\sigma} \right)^3 \right] = 0$$

$$E \left[ \left( \frac{X - \mu}{\sigma} \right)^3 \right] < 0$$



DeepLearning.AI

# Describing Distributions

---

**Skewness and Kurtosis:  
Kurtosis**

# Kurtosis: Example

## Game 1

Which one  
is riskier?

**probability**  $\frac{1}{2}$ : You win 1 dollar

**probability**  $\frac{1}{2}$ : You lose 1 dollar

## Game 2


**probability**  $\frac{100}{202}$ : You win 10 cents

**probability**  $\frac{100}{202}$ : You lose 10 cents


**probability**  $\frac{1}{202}$ : You win 10 dollars

**probability**  $\frac{1}{202}$ : You lose 10 dollars

# Kurtosis: Example




Expected value?  
Standard deviation?  
Skewness?



# Kurtosis: Example Expected Value


Game 1

$$\mathbb{E}[X_1] = 0$$




Game 2

$$\mathbb{E}[X_2] = 0$$



# Kurtosis: Example Variance


Game 1



$$Var(X_1) = 1$$

$$std(X_1) = 1$$

Game 2



$$Var(X_2) = 1$$

$$std(X_2) = 1$$


$$\begin{aligned}\mathbb{E}[X_1^2] &= \frac{1}{2}(-1)^2 + \frac{1}{2}(1)^2 \\ &= 1\end{aligned}$$

$$\begin{aligned}\mathbb{E}[X_2^2] &= \frac{100}{202}(-0.1)^2 + \frac{100}{202}(0.1)^2 + \frac{1}{202}(-10)^2 + \frac{1}{202}(10)^2 \\ &= \frac{100}{202} \cdot \frac{1}{100} + \frac{100}{202} \cdot \frac{1}{100} + \frac{1}{202} \cdot 100 + \frac{1}{202} \cdot 100 \\ &= \frac{1}{202} + \frac{1}{202} + \frac{100}{202} + \frac{100}{202} = 1\end{aligned}$$

# Kurtosis: Example Skewness


Game 1

$$Skew(X_1) = 0$$





Game 2

$$Skew(X_2) = 0$$



# Kurtosis



$$E[X_1] = 0$$

$$E[X_1] = 0$$

$$Var(X_1) = 1$$

$$E[X_1^2] = 1$$

$$Skew(X_1) = 0$$

$$E[X_1^3] = 0$$

$$E[X_2] = 0$$

$$E[X_2] = 0$$

$$Var(X_2) = 1$$

$$E[X_2^2] = 1$$


$$Skew(X_2) = 0$$

$$E[X_2^3] = 0$$

Has values way  
farther from 0


# Kurtosis

Game 1



$$\begin{aligned}\mathbb{E}[X_1^4] &= \frac{1}{2}(-1)^4 + \frac{1}{2}(1)^4 \\ &= 1\end{aligned}$$

Game 2



$$\begin{aligned}\mathbb{E}[X_2^4] &= \frac{100}{202}(-0.1)^4 + \frac{100}{202}(0.1)^4 + \frac{1}{202}(-10)^4 + \frac{1}{202}(10)^4 \\ &= 99.01\end{aligned}$$

# Kurtosis

$$\mathbb{E}[X^4]$$


Almost...

Need to standardize...


# Kurtosis

$$\text{Kurtosis} = E \left[ \left( \frac{X - \mu}{\sigma} \right)^4 \right]$$

# Kurtosis: High and Low



$$E \left[ \left( \frac{X - \mu}{\sigma} \right)^4 \right] = \text{small}$$



$$E \left[ \left( \frac{X - \mu}{\sigma} \right)^4 \right] = \text{large}$$

Even if they have the same variance!




DeepLearning.AI

# Describing Distributions

---

## Quantiles and Box-Plots

# Quantiles: Example



Newspaper advertisement

Newspaper Advertisement (X)			
18.3	18.4	23.2	51.2
35.2	29.7	75	8.7
65.9	14.2	54.7	25.9

# Quantiles: Example

What is the median here?

The point that splits your data in half

Newspaper Advertisement (X)			
18.3	18.4	23.2	51.2
35.2	29.7	75	8.7
65.9	14.2	54.7	25.9

# Quantiles: Example


What is the median here?

The point that splits your data in half

$$\text{Median} = \frac{25.9 + 29.7}{2} = 27.8$$

50% quantile

Second quartile




# Quantiles: Example

What about the point that leaves 1/4 of your data to the left and 3/4 to the right?

$$q_{0.25} = Q1 = \frac{18.3 + 18.4}{2} = 18.35$$

25% quantile

First quartile




# Quantiles

In general:


The **k%** quantile ( $q_{k/100}$ ) is the value that leaves k% of your data to the left and  $(100-k)\%$  of your data to the right

Some common quantiles:

- 25% quantile (first quartile - Q1)
- 50% quantile (median - Q2)
- 75% quantile (third quartile - Q3)




# Quantiles



$$\frac{20}{100} = \frac{n_{20}}{n} \approx \mathbf{P}(X \leq q_{0.2})$$

# Quantiles



**k% quantile** ( $q_{k/100}$ ) is the value such that  $\mathbf{P}(X \leq q_{k/100}) = \frac{k}{100}$




DeepLearning.AI

# Describing Distributions

---

**Visualizing data:  
Box-Plots**

# Box-Plots



Newspaper advertisement


Newspaper Advertisement (X)			
8.7	14.2	18.3	18.4
23.2	25.9	29.7	35.2
51.2	54.7	65.9	75

# Box-Plots

$$Q1 = q_{0.25} = \frac{18.3 + 18.4}{2} = 18.35$$

$$Q2 = q_{0.50} = \frac{25.9 + 29.7}{2} = 27.8$$

$$Q3 = q_{0.75} = \frac{51.2 + 54.7}{2} = 52.95$$



# Box-Plots

$$Q1 = q_{0.25} = \frac{18.3 + 18.4}{2} = 18.35$$


$$Q2 = q_{0.50} = \frac{25.9 + 29.7}{2} = 27.8$$

$$Q3 = q_{0.75} = \frac{51.2 + 54.7}{2} = 52.95$$

Interquartile range (IQR)

$$IQR = Q3 - Q1 = 52.95 - 18.35 = 34.6$$

$$\min = 8.7 \quad \max = 75$$



# Box-Plots

$$Q1 = q_{0.25} = \frac{18.3 + 18.4}{2} = 18.35$$


$$Q2 = q_{0.5} = \frac{25.9 + 29.7}{2} = 27.8$$

$$Q3 = q_{0.75} = \frac{51.2 + 54.7}{2} = 52.95$$

Interquartile range (IQR)

$$IQR = Q3 - Q1 = 52.95 - 18.35 = 34.6$$


$$x_{\min} = 8.7 \quad x_{\max} = 75$$



# Box-Plots

What can you tell from this plot?

- Data is skewed
- No outliers (whiskers were cut at max and min value)
- Analyze dispersion




# Box-Plots

Let's see how the box-plot looks for the whole dataset

$$Q_1 = 12.75 \quad Q_2 = 25.75 \quad Q_3 = 45.1$$

$$\text{IQR} = 32.35 \quad x_{\min} = 0.3 \quad x_{\max} = 114$$

Now you can see two outliers






DeepLearning.AI

# Describing Distributions

---

**Visualizing data:  
Kernel density estimation**

# Density Estimation



Newspaper advertisement

Newspaper Advertisement  
(X)

8.7	14.2	18.3	18.4
23.2	25.9	29.7	35.2
51.2	54.7	65.9	75


# Histograms

It represents a density function

- It is positive
- Area under the curve is 1


But...

- PDFs are usually smooth function
- The discontinuities come from the method and not the data




# Kernel Density Estimation

**First:** draw your observations along the x axis




# Kernel Density Estimation

**Second:** draw a gaussian centered at each observation




# Kernel Density Estimation

**Second:** draw a gaussian centered at each observation




# Kernel Density Estimation

**Second:** draw a gaussian centered at each observation




# Kernel Density Estimation

**Third:** multiply everything by  $1/n$  and sum the curves



# Kernel Density Estimation

What if  
you used  
all the  
dataset?







DeepLearning.AI

# Describing Distributions


---

**Visualizing data:  
Violin Plots**

# Violin Plots



# Violin Plots





DeepLearning.AI

# Describing Distributions

---

**Visualizing data:  
QQ plots**



# Assessing Normality of Data

Some models assume normally distributed data

- Linear regression
- Logistic regression
- Gaussian Naive Bayes
- Others

Some tests used in Data Science also assume normality.

# Assessing Normality of Data




# QQ Plots

Quantile-Quantile plots (QQ Plots) compare quantiles



- Standardize your data:

$$\left( \frac{x - \mu}{\sigma} \right)$$

- Compute quantiles
- Compare to gaussian quantiles



# QQ Plots



# W2 Lesson 2



DeepLearning.AI

# Probability Distributions with Multiple Variables

---

**Joint Distribution  
(Discrete)  
Part 1**

# Joint Distributions (Discrete): Example 1


Age (Year)	Count
7	3
8	2
9	4
10	1




**10**

children

# Joint Distributions (Discrete): Example 1



# Joint Distributions (Discrete): Example 1




# Joint Distributions (Discrete): Example 1

Age (years): 7    7    7    8    8    9    9    9    9    10



Height (in): 45    46    46    47    47    49    49    49    50    50

Height (in)	Count
45	1
46	2
47	2
48	0
49	3
50	2

# Joint Distributions (Discrete): Example 1





# Joint Distributions (Discrete): Example 1





What is the probability that a child is **9 years** old and **49 inches** tall?

# Joint Distributions (Discrete): Example 1





What is the probability that a child is **9 years** old and **49 inches** tall?

# Joint Distributions (Discrete): Example 1



What is the probability that a child is **9 years** old and **49 inches** tall?

# Joint Distributions (Discrete): Example 1



What is the probability that a child is 9 years old and 49 inches tall?

$$p_{XY}(9, 49) = \mathbf{P}(X = 9, Y = 49) = \frac{\text{Grey circle}}{10} = \frac{3}{10}$$

# Joint Distributions (Discrete): Example 1

What is the probability that a child is **9 years** old and **49 inches** tall?

$$p_{XY}(9, 49) = \mathbf{P}(X = 9, Y = 49) = \frac{\text{_____}}{10} = \frac{3}{10}$$

$$p_{XY}(x, y) = \mathbf{P}(X = x, Y = y)$$


# Joint Distributions: Example 1

		 	 	  	 	
Age (years):	7	7	8	9	9	10
Height (in):	45	46	47	49	49	50

Height

	45	46	47	48	49	50
7	1	2	0	0	0	0
8	0	0	2	0	0	0
9	0	0	0	0	3	1
10	0	0	0	0	0	1

# Joint Distributions: Example 1



# Joint Distributions: Example 1

## Joint Distribution

All probabilities for all possible combinations of X and Y

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		Probabilities					

# Joint Distributions: Example 1

What is the probability that a child is 8 and 48 inches tall?

$$p_{XY}(x, y) = \mathbf{P}(X = x, Y = y)$$

$$p_{XY}(8, 48) = \mathbf{P}(X = 8, Y = 48)$$

$$p_{XY}(8, 48) = 0$$

$$p_{XY}(7, 46) = \frac{2}{10}$$

	Height (Y)					
	45	46	47	48	49	50
Age (X)	7	1/10	2/10	0	0	0
7	0	0	0	2/10	0	0
8	0	0	0	0	0	0
9	0	0	0	0	3/10	1/10
10	0	0	0	0	0	1/10

Probabilities



DeepLearning.AI

# Probability Distributions with Multiple Variables

---

**Joint Distribution  
(Discrete)  
Part 2**

# Joint Distributions (Discrete): Example 2

$X$

the number rolled on the 1st dice



$$\frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6}$$


$Y$

the number rolled on the 2nd dice



$$\frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6}$$

# Joint Distributions: Example 2



# Joint Distributions: Example 2

$X$ : the number rolled on the 1st dice

$Y$ : the number rolled on the 2nd dice

**$X$  and  $Y$  are independent**



$$P(x) \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6$$

$$P(y) \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6 \quad 1/6$$

$$p_{XY}(\text{dice } 1, \text{dice } 2) = P(\text{dice } 1) \cdot P(\text{dice } 2) = \frac{1}{6} \cdot \frac{1}{6} = \frac{1}{36}$$

$$p_{XY}(x, y) = P(X = x, Y = y) = \frac{1}{36}$$

	1	2	3	4	5	6
1	1/36	1/36	1/36	1/36	1/36	1/36
2	1/36	1/36	1/36	1/36	1/36	1/36
3	1/36	1/36	1/36	1/36	1/36	1/36
4	1/36	1/36	1/36	1/36	1/36	1/36
5	1/36	1/36	1/36	1/36	1/36	1/36
6	1/36	1/36	1/36	1/36	1/36	1/36


# Joint Distributions: Example 2

Thus for independent discrete random variables:

$$p_{XY}(x, y) = \mathbf{P}(X = x, Y = y) = \mathbf{P}(x) \cdot \mathbf{P}(y)$$

# Joint Distributions (Discrete): Example 2

$X$



$Y$

the number rolled on the 1st dice


sum of the two dice

+

$$X = 4$$


$$Y = 4 + 5$$

# Joint Distributions: Example 2



# Joint Distributions - Example 3


$Y$ : Sum of both dice





# Joint Distributions: Example 3




# Joint Distributions: Example 3




# Joint Distributions: Example 3





# Joint Distributions: Example 3





# Joint Distributions: Example 3





# Joint Distributions: Example 3





# Joint Distributions: Example 3





# Joint Distributions: Example 3



# Joint Distributions: Example 3



# Joint Distributions: Example 3





# Joint Distributions: Example 3

Joint Distribution for  
 $X$  and  $Y$


36  
possible  
outcomes

Divide by sum  
(36)



# Joint Distributions: Example 3

Joint Distribution for  
X and Y



$$p_{XY}(3, 7) = \mathbf{P}(X = 3, Y = 7) = \frac{1}{36}$$

$$p_{XY}(1, 1) = \mathbf{P}(X = 1, Y = 1) = 0$$



DeepLearning.AI

# Probability Distributions with Multiple Variables

---

**Joint Distribution  
(Continuous)**

# Joint Continuous Distributions

$X$  : age of a child in year

$Y$  : discrete values of height of child in inches

$X$  : the number rolled on the 1st dice

$Y$  : the number rolled on the 2nd dice

$X$  : the number rolled on the 1st dice

$Y$  : sum of both dice


$X$  and  $Y$  are  
Discrete Random Variables

What about when  $X$  and  $Y$  are  
Continuous Random Variables?

# Joint Continuous Distributions



$X$  variable: Waiting time



# Joint Continuous Distributions

$X$

Waiting time  
before a call is picked up  
[0 - 10 minutes]



2.4 minutes

1.5 minutes

$Y$

Customer  
satisfaction rating  
[0 - 10]



0.0


5.7

Both variables are  
continuous

# Joint Continuous Distributions: Dataset

X variable: Waiting time (mins)  
0 - 10 mins

1000 customers




# Joint Continuous Distributions: Dataset

*Y* variable: Satisfaction rating

0 - 10

1000 customers




# Joint Continuous Distributions: Dataset



$X$  variable: Waiting time (mins)  
0 - 10 mins

$Y$  variable: Satisfaction rating  
0 - 10



1000 customers





# Joint Continuous Distributions: Dataset



# Joint Continuous Distributions: Dataset




# Joint Continuous Distributions: Dataset




# Joint Continuous Distributions: Dataset

3D Probability Density Distribution for Customer Ratings vs Waiting Time




Heatmap of Probability Density Function




# Joint Continuous Distributions: Dataset

3D Probability Density Distribution for Customer Ratings vs Waiting Time



Heatmap of Probability Density Function




# Joint Continuous Distributions: Dataset

$X$  variable: Waiting time (mins)  
0 - 10 mins

$Y$  variable: Satisfaction rating  
0 - 10

1000 customers



# Expected Value

$X$  variable: Waiting time (mins)

0 - 10 mins

$Y$  variable: Satisfaction rating

0 - 10

1000 customers


$$\mathbb{E}[X] = 4.903 \text{ minutes}$$

$$\mathbb{E}[Y] = 5.280$$



# Variances

$$Var(X)$$




# Variances

$$\mathbb{E}[X] = 4.903 \text{ minutes}$$

$$\mathbb{E}[X^2] = 32.561$$

$$\begin{aligned} \text{Var}(X) &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 \\ &= 32.561 - 4.903^2 \\ &= 8.526 \end{aligned}$$




# Variances

$$\mathbb{E}[Y] = 5.280$$


$$\mathbb{E}[Y^2] = 38.037$$

$$\begin{aligned} \text{Var}(Y) &= \mathbb{E}[Y^2] - \mathbb{E}[Y]^2 \\ &= 38.037 - 5.280^2 \\ &= 10.163 \end{aligned}$$



# Variances

$$Var(X)$$
  
$$8.526$$





DeepLearning.AI


# Probability Distributions with Multiple Variables

---

## Marginal and Conditional Distribution

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10



## Marginal Distribution

Distribution of one variable while ignoring the other

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10



To find the marginal distribution of height:

sum the joint probability distribution over all values of age

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		1/10	2/10	2/10	0	3/10	2/10



$$p_Y(y_j) = \sum_i p_{XY}(x_i, y_j)$$

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		1/10	2/10	2/10	0	3/10	2/10



$$p_Y(y_j) = \sum_i p_{XY}(x_i, y_j)$$

$$p_Y(50) =$$

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		1/10	2/10	2/10	0	3/10	2/10



$$p_Y(y_j) = \sum_i p_{XY}(x_i, y_j)$$

$$p_Y(50) = \sum_i p_{XY}(x_i, 50)$$

$$p_Y(50) = \frac{2}{10}$$

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		1/10	2/10	2/10	0	3/10	2/10

Age (years):	7	7	7	8	8	9	9	9	9	10
Height (in):	45	46	46	47	47	49	49	49	50	50

$$p_X(x_i) = \sum_j p_{XY}(x_i, y_j)$$

# Marginal Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10
		1/10	2/10	2/10	0	3/10	2/10



$$p_X(x_i) = \sum_j p_{XY}(x_i, y_j)$$

$$p_X(7) = \sum_j p_{XY}(7, y_j)$$

$$p_X(7) = \frac{3}{10}$$

# Marginal Distribution: Example 1

		Height ( $Y$ )						Age (years): 7 7 7 8 8 9 9 9 9 9 9 9 10									
		45	46	47	48	49	50	Height (in): 45 46 46 47 47 47 49 49 49 49 50 50									
Age ( $X$ )	7	1/10	2/10	0	0	0	0	3/10									
	8	0	0	2/10	0	0	0	2/10									
	9	0	0	0	0	3/10	1/10	4/10									
	10	0	0	0	0	0	1/10	1/10									
			1/10	2/10	2/10	0	3/10	2/10									

Marginal Distribution of Age




Marginal Distribution of Height



# Marginal Distribution: Example 1


Age and Height Dataset  
for 50 children



# Marginal Distribution: Example 1

Marginal Distribution of Age

Age and Height Dataset  
for 50 children



Marginal Distribution of Height

# Marginal Distributions: Example 2

$X$  : the number rolled on the 1st dice



$$\frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6}$$

$Y$  : the number rolled on the 2nd dice



$$\frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{6}$$

# Marginal Distributions: Example 2

		Y							
		1	2	3	4	5	6		
X		1	1/36	1/36	1/36	1/36	1/36	1/36	1/6
2		2	1/36	1/36	1/36	1/36	1/36	1/36	1/6
3		3	1/36	1/36	1/36	1/36	1/36	1/36	1/6
4		4	1/36	1/36	1/36	1/36	1/36	1/36	1/6
5		5	1/36	1/36	1/36	1/36	1/36	1/36	1/6
6		6	1/36	1/36	1/36	1/36	1/36	1/36	1/6
		1/6	1/6	1/6	1/6	1/6	1/6		

$X$ : the number rolled on the 1st dice

$Y$ : the number rolled on the 2nd dice

$p_X(x_i) = \frac{1}{6}$

$p_Y(y_j) = \frac{1}{6}$

# Marginal Distributions: Example 3



$X$  : the number rolled on the 1st dice

$Y$  : sum of the two dice

# Marginal Distributions: Example 3




$X$  : the number rolled on the 1st dice

$Y$  : sum of the two dice


$$p_Y(y_j) = \sum_i p_{XY}(x_i, y_j) = ?$$

$$p_Y(10) = ? = \frac{3}{36}$$

# Marginal Distributions: Example 2





# Marginal Distributions



# Continuous Marginal Distribution

3D Probability Density Distribution for Customer Ratings vs Waiting Time






DeepLearning.AI

# Probability Distributions with Multiple Variables

---


## Conditional Distribution

# Conditional Distribution: Example 1



# Conditional Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10



If age = 9, what is the distribution across the height variable?

## Conditional Distribution

# Conditional Distribution: Example 1

		Height ( $Y$ )					
		45	46	47	48	49	50
Age ( $X$ )	7	1/10	2/10	0	0	0	0
	8	0	0	2/10	0	0	0
	9	0	0	0	0	3/10	1/10
	10	0	0	0	0	0	1/10



If age = 9, what is the distribution across the height variable?

$$p_{Y|X=9}(y) = \mathbf{P}(Y = y | X = 9)$$

# Conditional Distribution: Example 1

Age (X)	Height (Y)					P(X = 9)
	45	46	47	48	49	
9	0	0	0	0	3/10	1/10
	Normalize	Divide by row sum				
9	0	0	0	0	3/4	1/4

$$\mathbf{P}(Y = 49 | X = 9) = \frac{3}{4}$$

$$p_{Y|X=9}(y) = \mathbf{P}(Y = y | X = 9)$$

$$\mathbf{P}(A, B) = \mathbf{P}(A) \cdot \mathbf{P}(B | A)$$

$$\mathbf{P}(X = 9, Y = 49) = P(X = 9) \cdot P(Y = 49 | X = 9)$$

$$P(Y = 49 | X = 9) = \frac{\mathbf{P}(X = 9, Y = 49)}{P(X = 9)}$$

Row sum

# Conditional Distribution: Example 1

(X)	Height (Y)					P(X = 9)	Sum
	45	46	47	48	49		
9	0	0	0	0	3/10	1/10	

$$p_{Y|X=x}(y) = \frac{p_{XY}(x, y)}{p_X(x)}$$

$$p_{Y|X=9}(y) = \mathbf{P}(Y = y | X = 9)$$

$$\mathbf{P}(A, B) = \mathbf{P}(A) \cdot \mathbf{P}(B | A)$$

$$\mathbf{P}(X = 9, Y = 49) = P(X = 9) \cdot P(Y = 49 | X = 9)$$

$$P(Y = 49 | X = 9) = \frac{\mathbf{P}(X = 9, Y = 49)}{P(X = 9)}$$

Row sum

$$\mathbf{P}(Y = 49 | X = 9) = \frac{3/10}{4/10} = \frac{3}{4}$$

# Discrete Conditional Distribution: Formula

$$p_{Y|X=x}(y) = \frac{p_{XY}(x, y)}{p_X(x)}$$

Conditional PDF of  $Y$

Joint PDF of  $X$  and  $Y$

Marginal distribution of  $X$

The diagram illustrates the formula for the conditional probability of  $Y$  given  $X=x$ . It features a central equation  $p_{Y|X=x}(y) = \frac{p_{XY}(x, y)}{p_X(x)}$ . Four curved arrows originate from text labels to specific parts of the equation: one arrow from 'Conditional PDF of  $Y$ ' points to the term  $p_{XY}(x, y)$ ; another from 'Joint PDF of  $X$  and  $Y$ ' points to the numerator  $p_{XY}(x, y)$ ; a third from 'Marginal distribution of  $X$ ' points to the denominator  $p_X(x)$ ; and a fourth arrow originates from the left side of the equation, pointing towards the conditional probability term.

# Conditional Distributions: Example 2




Dice 1: 1/6 1/6 1/6 1/6 1/6 1/6

Dice 2: 1/6 1/6 1/6 1/6 1/6 1/6

$$\begin{aligned} p_{Y|X=4}(y=1) &= \frac{p_{XY}(x=4, y=1)}{p_X(x=4)} \\ &= \frac{1/36}{1/6} \\ &= \frac{1}{6} \end{aligned}$$


	Y							
	1	2	3	4	5	6	Sum	
X	1	1/36	1/36	1/36	1/36	1/36	1/36	1/6
	2	1/36	1/36	1/36	1/36	1/36	1/36	1/6
	3	1/36	1/36	1/36	1/36	1/36	1/36	1/6
	4	1/36	1/36	1/36	1/36	1/36	1/36	1/6
	5	1/36	1/36	1/36	1/36	1/36	1/36	1/6
	6	1/36	1/36	1/36	1/36	1/36	1/36	1/6

# Conditional Distributions: Example 4




# Continuous Conditional Distribution


3D Probability Density Distribution for Customer Ratings vs Waiting Time




Probability distribution for rating given that waiting time was 4 minutes



# Continuous Conditional Distribution



Conditional PDF of  $y$  given  $x = 4$



# Discrete Conditional Distribution

$$p_{Y|X=x}(y) = \frac{p_{XY}(x, y)}{p_X(x)}$$

Conditional PDF of  $Y$

Joint PDF of  $X$  and  $Y$

Marginal distribution of  $X$

The diagram illustrates the derivation of the conditional probability formula. It shows three components: 'Conditional PDF of  $Y$ ' pointing to the numerator  $p_{XY}(x, y)$ ; 'Marginal distribution of  $X$ ' pointing to the denominator  $p_X(x)$ ; and 'Joint PDF of  $X$  and  $Y$ ' pointing to the overall fraction.

# Continuous Conditional Distribution: Formula

$$f_{Y|X=x}(y) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Conditional PDF of  $Y$

Joint PDF of  $X$  and  $Y$

Marginal distribution of  $X$

The diagram illustrates the formula for the conditional probability density function  $f_{Y|X=x}(y)$ . It features three curved teal arrows pointing from labels to parts of the formula. The first arrow points from 'Conditional PDF of  $Y$ ' to the term  $f_{Y|X=x}(y)$ . The second arrow points from 'Joint PDF of  $X$  and  $Y$ ' to the term  $f_{XY}(x, y)$ . The third arrow points from 'Marginal distribution of  $X$ ' to the term  $f_X(x)$ .



DeepLearning.AI

# Probability Distributions with Multiple Variables

---

## Covariance of a Dataset

# Introduction to Covariance

$Y_1$ : height of the child (in)

Age (X)	Height ( $Y_1$ )
6	50
7	52
8	55
9	57
10	60
11	62
12	64
13	65
14	67
15	68

$X$  : age of a child

$Y_2$  : grades in a test

Age (X)	Grades ( $Y_2$ )
6	5
7	7
8	8
9	3
10	1
11	1
12	6
13	10
14	2
15	7

- How is variable  $X$  related to each of the  $Y$  variables?
- How do you compare these relations?


$Y_3$  : number of naps per day

Age (X)	Naps per day ( $Y_3$ )
6	8
7	7
8	6
9	5
10	4
11	3
12	2
13	1
14	1
15	0


# Introduction to Covariance

$X$  : age of a child


$Y_1$  : height of the child (in)






$Y_2$  : grades in a test




$Y_3$  : number of naps per day




# How To Compare These?




# Mean?



$$\mu_x = 10.5 \quad \mu_y = 60$$




$$\mu_x = 10.5 \quad \mu_y = 5$$




$$\mu_x = 10.5 \quad \mu_y = 3.7$$


# Horizontal (X) Variance



$$Var(X) = 9.17$$






$$Var(X) = 9.17$$




$$Var(X) = 9.17$$

# Anything Else?




# Vertical (Y) Variance



$$Var(Y) = 39.56$$






$$Var(Y) = 9.78$$




$$Var(Y) = 7.57$$

# Still no Way To Compare Them



# First Step: Center Them




$$\mu_x = 0$$

$$\mu_y = 0$$

$$Var(X) = 1$$

$$Var(Y) = 1$$




$$\mu_x = 0$$

$$\mu_y = 0$$

$$Var(X) = 1$$

$$Var(Y) = 1$$






$$\mu_x = 0$$

$$\mu_y = 0$$


$$Var(X) = 1$$

$$Var(Y) = 1$$

# Second Step: Notice Trend




# Positives and Negatives




$$\sum xy > 0$$

Positive



$$\sum xy \approx 0$$

Both positive  
and negative



$$\sum xy < 0$$


Negative

# Covariance


$$Cov(X, Y) = \sum xy \quad \text{Almost...}$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$


# Covariance



$$\text{Cov}(X, Y) > 0$$



$$\text{Cov}(X, Y) \approx 0$$



$$\text{Cov}(X, Y) < 0$$

# Covariance Formula

Age vs Height

Covariance > 0

$$\mu_x = 10.5 \quad \mu_y = 60$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{170}{10} = 17 > 0$$

Age (x)	Height (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	50	-4.5	-10	45
7	52	-3.5	-8	28
8	55	-2.5	-5	12.5
9	57	-1.5	-3	4.5
10	60	-0.5	0	0
11	62	0.5	2	1
12	64	1.5	4	6
13	65	2.5	5	12.5
14	67	3.5	7	24.5
15	68	4.5	8	36

$$\sum = 170$$

# Covariance Formula

Age vs Naps per Day

Covariance < 0

$$\mu_x = 10.5 \quad \mu_y = 3.7$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{-74.5}{10} = -7.45 < 0$$

Age (x)	Naps (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	8	-4.5	4.3	-19.35
7	7	-3.5	3.3	-11.55
8	6	-2.5	2.3	-5.75
9	5	-1.5	1.3	-1.95
10	4	-0.5	0.3	-0.15
11	3	0.5	-0.7	-0.35
12	2	1.5	-1.7	-2.55
13	1	2.5	-2.7	-6.75
14	1	3.5	-2.7	-9.45
15	0	4.5	-3.7	-16.65

$$\sum = -74.5$$

# Covariance Formula

Age vs Grades

Covariance  $\approx 0$

$$\mu_x = 10.5 \quad \mu_y = 5$$


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$$Cov(X, Y) = \frac{1}{10} = 0.1 \quad \approx 0$$

Age (x)	Grades (y)	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
6	5	-4.5	0	0
7	7	-3.5	2	-7
8	8	-2.5	3	-7.5
9	3	-1.5	-2	3
10	1	-0.5	-4	2
11	1	0.5	-4	-2
12	6	1.5	1	1.5
13	10	2.5	5	12.5
14	2	3.5	-3	-10.5
15	7	4.5	2	9

$$\sum = 1$$


# Comparing Correlations



Age vs Height

Covariance > 0


$$Cov(x, y) = 17$$



Age vs Grades

Covariance  $\approx 0$

$$Cov(x, y) = 0.1$$



Age vs Naps per Day

Covariance < 0

$$Cov(x, y) = -7.45$$



DeepLearning.AI

# Probability Distributions with Multiple Variables

---


## Covariance of a Probability Distribution

# Covariance of a Probability Distribution: Motivation

X and Y are playing 3 games to either win or lose a dollar


## GAME 1

- $a$ : Both players win \$1 each  
 $b$ : Both players lose \$1 each




## GAME 2

- $c$ : X wins \$1 and Y loses \$1  
 $d$ : X loses \$1 and Y win \$1



## GAME 3


- $a, b, c$  or  $d$




How similar are these 3 games for player X and player Y?

# Covariance of a Probability Distribution: Motivation


GAME 1



GAME 2



GAME 3




How similar are these 3 games for player X and player Y?

$X$  : how much money in dollars player X wins


$Y$  : how much money in dollars player Y wins

# Covariance of a Probability Distribution: Motivation


GAME 1



GAME 2



GAME 3




$$\mathbb{E}[X_1] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

$$\mathbb{E}[Y_1] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

# Covariance of a Probability Distribution: Motivation


GAME 1



$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[Y_1] = 0$$


GAME 2



$$\mathbb{E}[X_2] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$


$$\mathbb{E}[Y_2] = \frac{1}{2}(1) + \frac{1}{2}(-1) = 0$$

GAME 3



# Covariance of a Probability Distribution: Motivation


**GAME 1**



$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[Y_1] = 0$$


**GAME 2**



$$\mathbb{E}[X_2] = 0$$

$$\mathbb{E}[Y_2] = 0$$

**GAME 3**




$$\mathbb{E}[X_3] = 2\left(\frac{1}{4}(1)\right) + 2\left(\frac{1}{4}(-1)\right) = 0$$


$$\mathbb{E}[Y_3] = 2\left(\frac{1}{4}(1)\right) + 2\left(\frac{1}{4}(-1)\right) = 0$$

# Covariance of a Probability Distribution: Motivation


**GAME 1**



**GAME 2**



**GAME 3**



$$Var(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$$

$$\mathbb{E}[X_1] = 0$$

$$\mathbb{E}[X_2] = 0$$

$$\mathbb{E}[X_3] = 0$$


$$\mathbb{E}[Y_1] = 0$$

$$\mathbb{E}[Y_2] = 0$$


$$\mathbb{E}[Y_3] = 0$$

# Covariance of a Probability Distribution: Motivation


**GAME 1**



**GAME 2**



**GAME 3**




$$\mathbb{E}[X_1] = 0 \quad \mathbb{E}[Y_1] = 0$$

$$\mathbb{E}[X_2] = 0 \quad \mathbb{E}[Y_2] = 0$$

$$\mathbb{E}[X_3] = 0 \quad \mathbb{E}[Y_3] = 0$$

$$Var(X_1) = \mathbb{E}[X_1^2] - E[X_1]^2 = \frac{1}{2}(-1)^2 + \frac{1}{2}(1)^2 - 0^2 = 1$$

# Covariance of a Probability Distribution: Motivation




# Covariance of a Probability Distribution: Motivation

Covariance of  $X$  and  $Y$ :  $\text{Cov}(X, Y)$

$$\text{Cov}(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

# Covariance of a Probability Distribution: Motivation




$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
-1	-1	-1	-1	1

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{2}{2} = 1$$

# Covariance of a Probability Distribution: Motivation



$$\mu_x = \mathbb{E}[X_1] = 0$$


$$\mu_y = \mathbb{E}[Y_1] = 0$$

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$x$	$y$	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
-1	-1	-1	-1	1

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{2}{2} = 1$$

# Covariance of a Probability Distribution: Motivation



$$\mu_x = \mathbb{E}[X_2] = 0$$

$$\mu_y = \mathbb{E}[Y_2] = 0$$


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

$x$	$y$	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	-1	1	-1	-1
-1	1	-1	1	-1

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{-2}{2} = -1$$

# Covariance of a Probability Distribution: Motivation

## GAME 3



$$\mu_x = \mathbb{E}[X_3] = 0$$

$$\mu_y = \mathbb{E}[Y_3] = 0$$


$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n}$$

x	y	$(x_i - \mu_x)$	$(y_i - \mu_y)$	$(x_i - \mu_x)(y_i - \mu_y)$
1	1	1	1	1
1	-1	-1	1	-1
-1	1	1	-1	-1
-1	-1	-1	-1	1

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{0}{4} = 0$$


# Covariance of a Probability Distribution: Motivation

**GAME 1**




$$Cov(X, Y) = 1$$

**GAME 2**



$$Cov(X, Y) = -1$$

**GAME 3**



$$Cov(X, Y) = 0$$


# Covariance of a Probability Distribution: Motivation

## GAME 4

a: Both players win \$1 each  $P(a) = 1/2$

b: Both players lose \$1 each  $P(b) = 1/3$

c: Neither players wins nor lose anything  $P(c) = 1/6$



### Unequal Probabilities

$$\mathbb{E}[X_4] = \frac{1}{2}(1) + \frac{1}{6}(0) + \frac{1}{3}(-1) = \frac{1}{6}$$

$$\mathbb{E}[Y_4] = \frac{1}{2}(1) + \frac{1}{6}(0) + \frac{1}{3}(-1) = \frac{1}{6}$$


# Covariance of a Probability Distribution: Motivation

## GAME 4

a: Both players win \$1 each  $\mathbf{P}(a) = 1/2$

b: Both players lose \$1 each  $\mathbf{P}(b) = 1/3$

c: Neither players wins nor lose anything  $\mathbf{P}(c) = 1/6$



### Unequal Probabilities

$$\text{Var}(X_4) = \sum_{i=1}^N (x_i - \mu_x)^2 \cdot \mathbf{P}(x_i)$$

$$\mathbb{E}[X_4] = \frac{1}{6}$$

$$\text{Var}(X_4) = \frac{1}{2}\left(1 - \frac{1}{6}\right)^2 + \frac{1}{6}\left(0 - \frac{1}{6}\right)^2 + \frac{1}{3}\left(-1 - \frac{1}{6}\right)^2$$

$$\mathbb{E}[Y_4] = \frac{1}{6}$$

$$= 0.806$$


# Covariance of a Probability Distribution: Motivation

## GAME 4

a: Both players win \$1 each  $P(a) = 1/2$

b: Both players lose \$1 each  $P(b) = 1/3$

c: Neither players wins nor lose anything  $P(c) = 1/6$



$$\mathbb{E}[X_4] = \frac{1}{6} \quad \text{Var}(X_4) = 0.806$$

$$\mathbb{E}[Y_4] = \frac{1}{6} \quad \text{Var}(Y_4) = 0.806$$

Unequal Probabilities

$\text{Cov}(X, Y) = ?$

# Covariance of Probability Distributions

$$Cov(X, Y) = \frac{\sum (x_i - \mu_x)(y_i - \mu_y)}{n} = \frac{1}{n} \sum (x_i - \mu_x)(y_i - \mu_y)$$


equal probabilities

$$Cov(X, Y) = \sum p_{XY}(x_i, y_i)(x_i - \mu_x)(y_i - \mu_y)$$

unequal probabilities

$$Cov(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

# Covariance of a Probability Distribution: Motivation



**Unequal Probabilities**

$$\begin{aligned} \text{Cov}(X, Y) &= \sum p_{XY}(x_i, y_i)(x_i - \mu_x)(y_i - \mu_y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] \\ &= \frac{1}{2}\left(1 - \frac{1}{6}\right)^2 + \frac{1}{6}\left(0 - \frac{1}{6}\right)^2 + \frac{1}{3}\left(-1 - \frac{1}{6}\right)^2 \end{aligned}$$

$$\mu_x = \mu_y = \mathbb{E}[X_4] = \mathbb{E}[Y_4] = 1/6$$

$$\text{Var}(X_4) = \text{Var}(Y_4) = 0.806$$

$$\text{Cov}(X, Y) = 0.806$$

# Covariance?

$$\mathbb{E}(X) = 4.903$$

$$\mathbb{E}(Y) = 5.280$$


$$\mathbb{E}(XY) = 18.014$$

$$Cov(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$$

$$Cov(X, Y) = 18.014 - (4.903)(5.280)$$

$$Cov(X, Y) = 18.014 - (4.903)(5.280)$$

$$Cov(X, Y) = -7.878$$



# Covariance of a Joint Continuous Distribution

$$Cov(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$$

$$\mathbb{E}(X) = 4.903$$

$$Cov(X, Y) = 18.014 - (4.903)(5.280)$$

$$\mathbb{E}(Y) = 5.280$$

$$Cov(X, Y) = 18.014 - (4.903)(5.280)$$

$$\mathbb{E}(XY) = 18.014$$

$$Cov(X, Y) = -7.878$$




DeepLearning.AI

# Probability Distributions with Multiple Variables

---

## Covariance Matrix

# Covariance Matrix




**Age vs Height**

$$\text{Var}(X) = 9.17$$

$$\text{Var}(Y) = 39.56$$

$$\text{Cov}(X, Y) = 17$$




**Age vs Grades**

$$\text{Var}(X) = 9.17$$

$$\text{Var}(Y) = 9.78$$

$$\text{Cov}(X, Y) = 0.1$$




**Age vs Naps per Day**

$$\text{Var}(X) = 9.17$$

$$\text{Var}(Y) = 7.57$$


$$\text{Cov}(X, Y) = -7.45$$

# Covariance Matrix




$$\text{Var}(X) = \text{Var}(Y) = 1$$

$$\text{Cov}(X, Y) = 1$$



$$\text{Var}(X) = \text{Var}(Y) = 1$$


$$\text{Cov}(X, Y) = -1$$



$$\text{Var}(X) = \text{Var}(Y) = 1$$

$$\text{Cov}(X, Y) = 0$$

# Covariance Matrix




$$\text{Var}(X) = 8.526$$


$$\text{Var}(Y) = 10.163$$

$$\text{Cov}(X, Y) = -7.878$$

# Covariance Matrix



# Covariance Matrix




$$\text{Var}(X) = 8.526$$

$$\text{Var}(Y) = 10.163$$

$$\text{Cov}(X, Y) = -7.878$$

# Covariance Matrix


# Covariance Matrix




$$Var(X) = 8.526$$

$$Var(Y) = 10.163$$

$$Cov(X, Y) = -7.878$$



# Covariance Matrix



$$Var(X) = 8.526$$

$$Var(Y) = 10.163$$

$$Cov(X, Y) = -7.878$$

	$X$	$Y$
$X$	$Var(X)$	$Cov(X, Y)$
$Y$	$Cov(X, Y)$	$Var(Y)$


$$\begin{bmatrix} Var(X) & Cov(X, Y) \\ Cov(X, Y) & Var(Y) \end{bmatrix}$$

8.526	-7.878
-7.878	10.163

$$\begin{bmatrix} 8.534 & -7.878 \\ -7.878 & 10.173 \end{bmatrix}$$

**Covariance Matrix**

# Covariance Matrix



Age vs Height

$$Var(X) = 9.17$$


$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$

$$\begin{bmatrix} Var(X) & Cov(X, Y) \\ Cov(X, Y) & Var(Y) \end{bmatrix}$$

$$\begin{bmatrix} 9.17 & 17 \\ 17 & 39.56 \end{bmatrix}$$

# Covariance Matrix




$$\begin{bmatrix} \mathbf{Var}(X) & \mathbf{Cov}(X, Y) \\ \mathbf{Cov}(X, Y) & \mathbf{Var}(Y) \end{bmatrix}$$

$$\mathbf{Var}(X) = 1$$

$$\mathbf{Var}(Y) = 1$$

$$\mathbf{Cov}(X, Y) = 1$$

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$



$$\mathbf{Var}(X) = 1$$

$$\mathbf{Var}(Y) = 1$$

$$\mathbf{Cov}(X, Y) = -1$$

$$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$$

# Covariance of a Joint Continuous Distribution

Dataset with 3 Variables:

X	Y	Z	$Var(X)$	$Cov(X, Y)$
X	Y	Z	$Var(Y)$	$Cov(X, Z)$
X	Y	Z	$Var(Z)$	$Cov(Y, Z)$

X	$Var(X)$	$Cov(X, Y)$	$Cov(X, Z)$
Y	$Cov(X, Y)$	$Var(Y)$	$Cov(Y, Z)$
Z	$Cov(X, Z)$	$Cov(Y, Z)$	$Var(Z)$

# Covariance of a Joint Continuous Distribution

Dataset with 5 Variables:

V	W	X	Y	Z
			$Var(V)$	$Cov(V, W)$
			$Var(W)$	$Cov(V, X)$
			$Var(X)$	$Cov(V, Y)$
			$Var(Y)$	$Cov(V, Z)$
			$Var(Z)$	$Cov(W, X)$
				$\vdots$
				$Cov(Y, Z)$

$Var(V) \quad Cov(V, W)$   
 $Var(W) \quad Cov(V, X)$   
 $Var(X) \quad Cov(V, Y)$   
 $Var(Y) \quad Cov(V, Z)$   
 $Var(Z) \quad Cov(W, X)$   
 $\vdots$   
 $Cov(Y, Z)$

	V	W	X	Y	Z
V	$Var(V)$	$Cov(V, W)$	$Cov(V, X)$	$Cov(V, Y)$	$Cov(V, Z)$
W	$Cov(V, W)$	$Var(W)$	$Cov(W, X)$	$Cov(W, Y)$	$Cov(W, Z)$
X	$Cov(V, X)$	$Cov(W, X)$	$Var(X)$	$Cov(X, Y)$	$Cov(X, Z)$
Y	$Cov(V, Y)$	$Cov(W, Y)$	$Cov(X, Y)$	$Var(Y)$	$Cov(Y, Z)$
Z	$Cov(V, Z)$	$Cov(W, Z)$	$Cov(X, Z)$	$Cov(Y, Z)$	$Var(Z)$

# Covariance of a Joint Continuous Distribution

$\sum =$

## Covariance Matrix

	$V$	$W$	$X$	$Y$	$Z$
$V$	$Var(V)$	$Cov(V, W)$	$Cov(V, X)$	$Cov(V, Y)$	$Cov(V, Z)$
$W$	$Cov(V, W)$	$Var(W)$	$Cov(W, X)$	$Cov(W, Y)$	$Cov(W, Z)$
$X$	$Cov(V, X)$	$Cov(W, X)$	$Var(X)$	$Cov(X, Y)$	$Cov(X, Z)$
$Y$	$Cov(V, Y)$	$Cov(W, Y)$	$Cov(X, Y)$	$Var(Y)$	$Cov(Y, Z)$
$Z$	$Cov(V, Z)$	$Cov(W, Z)$	$Cov(X, Z)$	$Cov(Y, Z)$	$Var(Z)$




DeepLearning.AI

# Probability Distributions with Multiple Variables

---

## Correlation Coefficient

# Correlation Coefficient




Age vs Naps per Day

$$Var(X) = 9.17$$

$$Var(Y) = 7.57$$

$$Cov(X, Y) = -7.45$$

Is the correlation here strong?



Age vs Height

$$Var(X) = 9.17$$

$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$

# Correlation Coefficient

Age vs Naps per Day

$$Var(X) = 9.17$$

$$Var(Y) = 7.57$$

$$Cov(X, Y) = -7.45$$

-1



0

1




## Correlation Coefficient

Age vs Height

$$Var(X) = 9.17$$

$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$



# Correlation Coefficient

Age vs Naps per Day

$$Var(X) = 9.17$$

$$Var(Y) = 7.57$$

$$Cov(X, Y) = -7.45$$

-1

0

1

Age vs Height

$$Var(X) = 9.17$$

$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$

$$\text{Correlation Coefficient} = \frac{Cov(X, Y)}{\sigma_x \cdot \sigma_y} = \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}}$$


# Correlation Coefficient

Age vs Naps per Day

$$Var(X) = 9.17$$

$$Var(Y) = 7.57$$

$$Cov(X, Y) = -7.45$$



$$\begin{aligned}\text{Correlation Coefficient} &= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}} \\ &= \frac{-7.45}{\sqrt{9.17} \cdot \sqrt{7.57}} \\ &\approx -0.894\end{aligned}$$


# Correlation Coefficient

Age vs Height

$$Var(X) = 9.17$$

$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$



Correlation  
Coefficient

$$= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}}$$

$$= \frac{17}{\sqrt{9.17} \cdot \sqrt{39.56}}$$

$\approx 0.893$

# Correlation Coefficient

Age vs Naps per Day

$$Var(X) = 9.17$$

$$Var(Y) = 7.57$$

$$Cov(X, Y) = -7.45$$




Age vs Height


$$Var(X) = 9.17$$

$$Var(Y) = 39.56$$

$$Cov(X, Y) = 17$$



# Correlation Coefficient



$$Var(X) = 9.17$$


$$Var(Y) = 9.78$$

$$Cov(X, Y) = 0.1$$

$$\begin{aligned} \text{Correlation Coefficient} &= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}} \\ &= \frac{0.1}{\sqrt{9.17} \cdot \sqrt{9.78}} \end{aligned}$$

$$\approx 0.01$$

# Correlation Coefficient



$$Var(X) = 8.526$$

$$Var(Y) = 10.163$$


$$Cov(X, Y) = -7.878$$

$$\begin{aligned}\text{Correlation Coefficient} &= \frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}} \\ &= \frac{-7.878}{\sqrt{8.562} \cdot \sqrt{10.163}}\end{aligned}$$

$\approx -0.845$

# Correlation Coefficient

**Correlation Coefficient** =  $\frac{Cov(X, Y)}{\sigma_x \cdot \sigma_y}$  =  $\frac{Cov(X, Y)}{\sqrt{Var(X)} \cdot \sqrt{Var(Y)}}$





DeepLearning.AI

# Probability Distributions with Multiple Variables

---

## Multivariate Gaussian Distribution


# Multivariate Gaussian Distribution

For a single variable,  $X$

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$

Parameters:

- $\mu$ : center of the bell
- $\sigma$ : spread of the bell



# Multivariate Gaussian Distribution: an Example


Two variables

$H$  : Height of an adult in inches


$$H \sim \mathcal{N}(\mu_H, \sigma_H)$$

$W$  : Weight of an adult in pounds

$$W \sim \mathcal{N}(\mu_W, \sigma_W)$$



# Multivariate Gaussian Distribution: an Example





If  $W, H$  were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$



$$= \frac{1}{\sqrt{2\pi}\sigma_H} e^{-\frac{1}{2}\frac{(h-\mu_H)^2}{\sigma_H^2}} \cdot \frac{1}{\sqrt{2\pi}\sigma_W} e^{-\frac{1}{2}\frac{(w-\mu_W)^2}{\sigma_W^2}}$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

# Multivariate Gaussian Distribution: an Example



# Multivariate Gaussian Distribution: an Example



# Multivariate Gaussian Distribution: an Example

If  $W, H$  were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$
$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$\left\| \begin{bmatrix} \frac{h-\mu_H}{\sigma_H} \\ \frac{w-\mu_W}{\sigma_W} \end{bmatrix} \right\|^2 = \begin{bmatrix} h-\mu_H & w-\mu_W \\ \sigma_H & \sigma_W \end{bmatrix} \begin{bmatrix} h-\mu_H \\ w-\mu_W \end{bmatrix}$$

$\left[ \begin{array}{c} h - \mu_h \\ w - \mu_w \end{array} \right] = \left[ \begin{array}{c} h \\ w \end{array} \right] - \left[ \begin{array}{c} \mu_h \\ \mu_w \end{array} \right]$

Multiply by diagonal matrix

# Multivariate Gaussian Distribution: an Example

If  $W, H$  were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$
$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

det ( $\Sigma$ )<sup>1/2</sup>

$$\left\| \begin{bmatrix} \frac{h-\mu_H}{\sigma_H} \\ \frac{w-\mu_W}{\sigma_W} \end{bmatrix} \right\|_2^2 = \begin{bmatrix} h-\mu_H & w-\mu_W \\ \sigma_H & \sigma_W \end{bmatrix} \begin{bmatrix} \frac{1}{\sigma_H^2} & 0 \\ 0 & \frac{1}{\sigma_W^2} \end{bmatrix} \begin{bmatrix} h-\mu_H \\ w-\mu_W \\ \sigma_W \end{bmatrix}$$
$$= ([h \ w] - [\mu_H \ \mu_W]) \begin{bmatrix} \frac{1}{\sigma_H^2} & 0 \\ 0 & \frac{1}{\sigma_W^2} \end{bmatrix} \left( [h \ w] - [\mu_H \ \mu_W] \right)$$

Covariance matrix!  
( $\Sigma$ )

$$= \left( [h \ w] - [\mu_H \ \mu_W] \right)^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} \left( [h \ w] - [\mu_H \ \mu_W] \right)$$

$\mu$

# Multivariate Gaussian Distribution: an Example

If  $W, H$  were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} \exp\left(-\frac{1}{2} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \begin{bmatrix} \mu_H \\ \mu_W \end{bmatrix} \right)^T \begin{bmatrix} \sigma_H^2 & 0 \\ 0 & \sigma_W^2 \end{bmatrix}^{-1} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \begin{bmatrix} \mu_H \\ \mu_W \end{bmatrix} \right)\right)$$

# Multivariate Gaussian Distribution: an Example

If  $W, H$  were independent

$$f_{HW}(h, w) = f_H(h)f_W(w)$$

$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$= \frac{1}{2\pi \det \Sigma^{1/2}} \exp \left( -\frac{1}{2} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu} \right)^T \boldsymbol{\Sigma^{-1}} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu} \right) \right)$$

# Multivariate Gaussian Distribution: an Example

Dependent case:

$$f_{HW}(h, w) = \cancel{f_H(h)f_W(w)}$$
$$= \frac{1}{2\pi\sigma_H\sigma_W} e^{-\frac{1}{2}\left(\frac{(h-\mu_H)^2}{\sigma_H^2} + \frac{(w-\mu_W)^2}{\sigma_W^2}\right)}$$

$$\Sigma = \begin{bmatrix} \sigma_H^2 & Cov(H, W) \\ Cov(H, W) & \sigma_W^2 \end{bmatrix}$$

$$= \frac{1}{2\pi \det \Sigma^{1/2}} \exp \left( -\frac{1}{2} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu} \right)^T \Sigma^{-1} \left( \begin{bmatrix} h \\ w \end{bmatrix} - \boldsymbol{\mu} \right) \right)$$

# Multivariate Gaussian Distribution: General Definition

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\frac{(x-\mu)^2}{\sigma^2}}$$

$x = [x_1 \ x_2 \ \dots \ x_n]^T$

$\mu = [\mu_1, \mu_2, \dots, \mu_n]^T$

Mean vector

$f_X(x_1, x_2, \dots, x_n)$

random variables

$X = [X_1 \ X_2 \ \dots \ X_n]$

• For univariate, we work with scalar values and variances

• For multivariate, we work with vectors and the covariance matrix

covariance matrix / spread of the bell

$|\Sigma|$  determinant of the covariance matrix



DeepLearning.AI

# Probability Distributions with Multiple Variables

---

## Conclusion