

## Laboratory 09

### Finite Element method for the Stokes problem

#### Exercise 1.

Let  $\Omega \subset \mathbb{R}^3$  be the domain shown in Figure 1. Let us consider the stationary Stokes problem:


$$\begin{cases} -\nu \Delta \mathbf{u} + \nabla p = \mathbf{f} & \text{in } \Omega, & (1a) \\ \nabla \cdot \mathbf{u} = 0 & \text{in } \Omega, & (1b) \\ \mathbf{u} = \mathbf{u}_{\text{in}} & \text{on } \Gamma_{\text{in}}, & (1c) \\ \nu(\nabla \mathbf{u})\mathbf{n} - p\mathbf{n} = -p_{\text{out}}\mathbf{n} & \text{on } \Gamma_{\text{out}}, & (1d) \\ \mathbf{u} = \mathbf{0} & \text{on } \Gamma_{\text{wall}}, & (1e) \end{cases}$$

where  $\mathbf{u} : \Omega \rightarrow \mathbb{R}^3$  and  $p : \Omega \rightarrow \mathbb{R}$  are the velocity and pressure fields of a viscous, incompressible fluid,  $\nu = 1 \text{ m}^2/\text{s}$ ,  $\mathbf{f} = \mathbf{0} \text{ m/s}^2$ ,  $\mathbf{u}_{\text{in}} = (-\alpha y(2-y)(1-z)(2-z) \text{ m/s}, 0 \text{ m/s}, 0 \text{ m/s})^T$ ,  $\alpha = 1/(\text{m}^3 \cdot \text{s})$ ,  $p_{\text{out}} = 10 \text{ Pa}$ .


**1.1.** Derive the weak formulation of the problem.

**1.2.** Derive the finite element formulation to problem (1).


**1.3.** Implement a finite element solver for (1) and compute its numerical solution using the mesh `mesh/mesh-step-5.msh` (whose boundary tags are described in Figure 1).



$\Gamma_{\text{in}}$  (mesh tag 0).



$\Gamma_{\text{wall}}$  (mesh tag 1).



$\Gamma_{\text{out}}$  (mesh tag 2).

Figure 1: Domain and partition of its boundary.