
## Module 3: Assignment 3

## Ellen Bledsoe

## 2022-11-03

```
3.
## # A tibble: 4 x 3
                mean_fish sd_fish
##
     <chr>
                     <dbl>
                             <dbl>
## 1 Emporer
                             10.9
                      24.9
## 2 Hope
                              9.73
                      24.3
                              9.94
## 3 Iceberg
                      25.8
## 4 Sulzberger
                              9.98
                      24.9
```




```
5.
## Df Sum Sq Mean Sq F value Pr(>F)
## bay 3 542 180.6 1.751 0.155
## Residuals 1916 197625 103.1
8.
```

## Tukey multiple comparisons of means

```
95% family-wise confidence level
##
##
## Fit: aov(formula = fish_kg ~ bay, data = fish)
##
## $bay
##
                              diff
                                          lwr
                                                    upr
## Hope-Emporer
                      -0.53796408 -2.2236167 1.1476886 0.8447466
## Iceberg-Emporer
                       0.94195591 -0.7436968 2.6276086 0.4764075
## Sulzberger-Emporer 0.04106038 -1.6445923 1.7267130 0.9999118
## Iceberg-Hope
                       1.47991999 -0.2057327 3.1655727 0.1085321
## Sulzberger-Hope
                        0.57902446 \ -1.1066282 \ 2.2646771 \ 0.8135811 
## Sulzberger-Iceberg -0.90089553 -2.5865482 0.7847571 0.5157262
 11.
## # A tibble: 4 x 3
##
     bay
                mean_fish sd_fish
##
     <chr>
                    <dbl>
                             <dbl>
                     9.96
                             1.19
## 1 Emporer
                             1.02
## 2 Hope
                     1.96
## 3 Iceberg
                     6.02
                             2.13
## 4 Sulzberger
                     2.08
                             0.958
```

12 -

12.



13. ## Df Sum Sq Mean Sq F value Pr(>F) ## bay 8682 2894 1467 <2e-16 \*\*\* ## Residuals 796 1570

```
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
 15.
##
    Tukey multiple comparisons of means
      95% family-wise confidence level
##
## Fit: aov(formula = num_seals ~ bay, data = seals)
##
## $bay
##
                     diff
                                lwr
                                          upr
                                                  p adj
## Hope-Emporer
                    -8.00 -8.3615601 -7.6384399 0.0000000
## Iceberg-Emporer
                    -3.94 -4.3015601 -3.5784399 0.0000000
## Sulzberger-Emporer -7.88 -8.2415601 -7.5184399 0.0000000
## Iceberg-Hope
                     4.06 3.6984399 4.4215601 0.0000000
                     ## Sulzberger-Hope
## Sulzberger-Iceberg -3.94 -4.3015601 -3.5784399 0.0000000
```