


The Mutual Information Diagram for Uncertainty Visualization

Carlos D. Correa and Peter Lindstrom



Center for Applied Scientific Computing - CASC
Lawrence Livermore National Lab

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.





Visualization should help compare models with observations




Average annual temperature 1900-2000 as predicted by various climate models.

Which model is more **similar** to a reference model or observations?

Trend plots often do not expose these aspects.



Visualization should help find correlations of similar outputs – important for uncertainty quantification




Divide ensembles in 6 latitude zones and 3 temporal averages


- Are there **correlations** across seasons or latitudes?
- Are there large discrepancies in the different outputs?





Visualization should be important





A scatterplot matrix becomes **impractical** for many outputs







Visual Summaries





- Represent directly summary quantities, e.g., mean, standard deviation, entropy.

- Box-plots and their many variants

- One plot per ensemble may result in clutter

- Visualizing several statistics simultaneously in **a metric space**: Taylor diagram






The Taylor Diagram

Simultaneously plots

Standard deviation,
Root Mean Square Error and
Correlation R.


$$RMS^2 = \sigma_X^2 + \sigma_Y^2 - 2\sigma_X\sigma_Y R_{XY}$$

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$





Applications of the Taylor Diagram






Taylor, 2005
(Taylor Diagram Primer)





Anscombe's Trio

Variables B,C,D: **same** standard deviation and **same** correlation w.r.t. A




Correlation
0.00
0.10
0.20
0.30
0.40

Information Theory to
the Rescue!





Information Theory Primer




$$I(X;Y) = H(X) + H(Y) - H(X, Y)$$

- **Entropy** $H(X)$
 - Measure of information uncertainty of X
- **Joint Entropy** $H(X, Y)$
 - Uncertainty of X, Y
- **Conditional Entropy** $H(X|Y)$
 - Uncertainty of X given that I know Y
- **Mutual Information** $I(X;Y)$
 - How much knowing X reduces the uncertainty of Y



Variation of Information $VI = H(X | Y) + H(Y | X)$





The Variation of Information VI: a measure of distance in information theory

$$VI(X, Y) = H(X) + H(Y) - 2I(X; Y)$$

$$RVI = \sqrt{VI}$$

$$h_X = \sqrt{H(X)}$$

$$h_Y = \sqrt{H(Y)}$$

$$RVI(X, Y)^2 = h_X^2 + h_Y^2 - 2I(X; Y)$$

$$RVI(X, Y)^2 = h_X^2 + h_Y^2 - 2h_X h_Y \frac{I(X; Y)}{h_X h_Y}$$



The Variation of Information VI: a measure of distance in information theory

$$VI(X, Y) = H(X) + H(Y) - 2I(X; Y)$$

$$RVI = \sqrt{VI}$$

$$h_X = \sqrt{H(X)}$$

$$h_Y = \sqrt{H(Y)}$$

$$RVI(X, Y)^2 = h_X^2 + h_Y^2 - 2I(X; Y)$$



$$RVI(X, Y)^2 = h_X^2 + h_Y^2 - 2h_X h_Y \frac{I(X; Y)}{h_X h_Y}$$

Normalized Mutual
Information (NMI)

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$



RVI Diagram



Equivalences



Statistics \iff Information Theory

$$\text{RMS}(X, Y) \iff \text{RVI} \sqrt{VI(X, Y)}$$

$$\text{Variance } \sigma_X^2 \iff \text{entropy } H(X)$$


$$\text{Covariance } cov(X, Y) \iff \text{mutual information } I(X; Y)$$

$$\text{Correlation } R_{XY} = \frac{cov(X, Y)}{\sqrt{\sigma_X^2 \sigma_Y^2}} \iff NMI_{XY} = \frac{I(X; Y)}{\sqrt{H(X)H(Y)}}$$






VI Diagram



$$VI(X, Y)^2 = H(X)^2 + H(Y)^2 - 2H(X)H(Y)c_{XY}$$




Experiment of 2D distributions with outliers






MI diagram is more resilient to outliers

Outliers have a significant impact on correlation





The information in both the “clean” and “dirty” distributions is essentially the same





Computing Entropy and Mutual Information *may* require estimation of underlying probability functions





Although there are differences,
relative distances are **consistent** for
each choice of kernel



Uncertainty Quantification in Climate Simulations




Precipitation average
7 Zonal averages (color)
3 Temporal averages (shape)
3 different ensemble sets (size)





Intercomparison Studies




Annual mean temperature 1900-2000





Intercomparison Studies




Annual mean temperature 1900-2000






Intercomparison Studies

Annual mean temperature 1900-2000





MID applies to discrete data: useful when comparing Clustering Results




- Summarize study in clustering [Filippone et al. 2009]
- 8 different methods
- 4 classification problems

Concluding Remarks

- Taylor diagram:
 - easy to compute.
 - Well understood in geophysical sciences, climate.
- MI diagram:
 - Counterpart using information theory.
 - requires an estimation step that may introduce additional uncertainties.
 - extends nicely to categorical data, multi-variate distributions.
 - exposes non-linearities, difficult to see via (linear) correlation.
- More informed decisions when combining both diagrams.

Thanks!



Questions?