

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Week 10](#) > [Practic...](#) > Q4: Val...

Q4: Value Functions

Problem 4: Value Functions

Consider a general search problem defined by:

- A set of states, \mathcal{S} .
- A start state s_0 .
- A set of goal states \mathcal{G} , with $\mathcal{G} \subset \mathcal{S}$.
- A successor function $\text{Succ}(s)$ that gives the set of states s' that you can go to from the current state s .
- For each successor s' of s , the cost (weight) $W(s, s')$ of that action.

As usual, the search problem is to find a lowest-cost path from the start state s_0 to a goal $g \in \mathcal{G}$. You may assume that each non-goal state has at least one successor, that the weights are all positive, and that all states can reach a goal state in a finite number of steps.

Define $C(s)$ to be the *optimal cost* of the state s ; that is, the cost of the lowest-cost path from s to any goal state. For $g \in \mathcal{G}$, clearly $C(g) = 0$.

Part 1

4/4 points (ungraded)

Select the Bellman-style (one-step lookahead) equation that expresses $C(s)$ for a non-goal state s in terms of the optimal costs of other states.

☐ $C(s) = \sum_{s' \in \text{Succ}(s)} W(s, s')$

☐ $C(s) = \min_{s' \in \text{Succ}(s)} W(s, s')$

☐ $C(s) = \max_{s' \in \text{Succ}(s)} W(s, s')$

☐ $C(s) = \sum_{s' \in \text{Succ}(s)} [W(s, s') + C(s')]$

☒ $C(s) = \min_{s' \in \text{Succ}(s)} [W(s, s') + C(s')] \checkmark$

☐ $C(s) = \max_{s' \in \text{Succ}(s)} [W(s, s') + C(s')]$

✓ Correct (4/4 points)

Part 2

2/2 points (ungraded)

Consider a heuristic function $h(s)$ with $h(s) \geq 0$. What relation must hold between $h(s)$ and $C(s)$ for $h(s)$ to be an admissible heuristic?

☐ $h(s) = C(s), \forall s \in S$

☒ $h(s) \leq C(s), \forall s \in S \checkmark$

☐ $h(s) \geq C(s), \forall s \in S$

Submit

✓ Correct (2/2 points)

Part 3

By analogy to value iteration, define $C_k(s)$ to be the minimum cost of any plan starting from s that is either of length k or reaches a goal state in at most k actions. Define $C_0(s) = 0$ for all s . $C(s)$ is defined as before to be the cost of the lowest-cost path from s to any goal state, without any restriction on the number of actions starting from s .

Suppose we use C and C_k as heuristic functions.

Part 3.1

2/2 points (ungraded)

Which of the following statements regarding the admissibility of C and C_k are true? Check all that apply.

☐ C_k might be inadmissible for any given value of k .

☒ C_k is admissible for all k .

☐ Let L be the length of the longest path from a state to a goal. C_k is only guaranteed to be admissible if $k \geq L$.

☒ C is admissible.



Submit

✓ Correct (2/2 points)

Part 3.2

2/2 points (ungraded)

Which of the following statements regarding the consistency of C and C_k are true? Check all that apply.

☐ C_k might be inconsistent for any given value of k .

☒ C_k is consistent for all k .

☐ Let L be the length of the longest path from a state to a goal. C_k is only guaranteed to be consistent if $k \geq L$.

☒ C is consistent.



Submit

✓ Correct (2/2 points)

© All Rights Reserved